



Reference Tables for Quantity of F-100 to Give an Individual Child per Feed

Weight of child (kg)	Range of volumes per 3-hourly feed of F-100 (8 feeds daily) *		Range of volumes per 4-hourly feed of F-100 (6 feeds daily) *		Range of daily volumes of F-100		Weight of child (kg)	Range of volumes per 3-hourly feed of F-100 (8 feeds daily) *		Range of volumes per 4-hourly feed of F-100 (6 feeds daily) *		Range of daily volumes of F-100	
	Minimum ml	Maximum ml	Minimum ml	Maximum ml	Minimum (150 ml/kg/day)	Maximum (220 ml/kg/day)		Minimum ml	Maximum ml	Minimum ml	Maximum ml	Minimum (150 ml/kg/day)	Maximum (220 ml/kg/day)
2.0	40	55	50	75	300	440	11.0	205	305	275	405	1650	2420
2.2	40	60	55	80	330	484	11.2	210	310	280	410	1680	2464
2.4	45	65	60	90	360	528	11.4	215	315	285	420	1710	2508
2.6	50	70	65	95	390	572	11.6	220	320	290	425	1740	2552
2.8	55	75	70	105	420	616	11.8	220	325	295	435	1770	2596
3.0	55	85	75	110	450	660	12.0	225	330	300	440	1800	2640
3.2	60	90	80	115	480	704	12.2	230	335	305	445	1830	2684
3.4	65	95	85	125	510	748	12.4	235	340	310	455	1860	2728
3.6	70	100	90	130	540	792	12.6	235	345	315	460	1890	2772
3.8	70	105	95	140	570	836	12.8	240	350	320	470	1920	2816
4.0	75	110	100	145	600	880	13.0	245	360	325	475	1950	2860
4.2	80	115	105	155	630	924	13.2	250	365	330	485	1980	2904
4.4	85	120	110	160	660	968	13.4	250	370	335	490	2010	2948
4.6	85	125	115	170	690	1012	13.6	255	375	340	500	2040	2992
4.8	90	130	120	175	720	1056	13.8	260	380	345	505	2070	3036
5.0	95	140	125	185	750	1100	14.0	265	385	350	515	2100	3080
5.2	100	145	130	190	780	1144	14.2	265	390	355	520	2130	3124
5.4	100	150	135	200	810	1188	14.4	270	395	360	530	2160	3168
5.6	105	155	140	205	840	1232	14.6	275	400	365	535	2190	3212
5.8	110	160	145	215	870	1276	14.8	280	405	370	545	2220	3256
6.0	115	165	150	220	900	1320	15.0	280	415	375	550	2250	3300
6.2	115	170	155	225	930	1364	15.2	285	420	380	555	2280	3344
6.4	120	175	160	235	960	1408	15.4	290	425	385	565	2310	3388
6.6	125	180	165	240	990	1452	15.6	295	430	390	570	2340	3432
6.8	130	185	170	250	1020	1496	15.8	295	435	395	580	2370	3476
7.0	130	195	175	255	1050	1540	16.0	300	440	400	585	2400	3520
7.2	135	200	180	265	1080	1584	16.2	305	445	405	595	2430	3564
7.4	140	205	185	270	1110	1628	16.4	310	450	410	600	2460	3608
7.6	145	210	190	280	1140	1672	16.6	310	455	415	610	2490	3652
7.8	145	215	195	285	1170	1716	16.8	315	460	420	615	2520	3696
8.0	150	220	200	295	1200	1760	17.0	320	470	425	625	2550	3740
8.2	155	225	205	300	1230	1804	17.2	325	475	430	630	2580	3784
8.4	160	230	210	310	1260	1848	17.4	325	480	435	640	2610	3828
8.6	160	235	215	315	1290	1892	17.6	330	485	440	645	2640	3872
8.8	165	240	220	325	1320	1936	17.8	335	490	445	655	2670	3916
9.0	170	250	225	330	1350	1980	18.0	340	495	450	660	2700	3960
9.2	175	255	230	335	1380	2024	18.2	340	500	455	665	2730	4004
9.4	175	260	235	345	1410	2068	18.4	345	505	460	675	2760	4048
9.6	180	265	240	350	1440	2112	18.6	350	510	465	680	2790	4092
9.8	185	270	245	360	1470	2156	18.8	355	515	470	690	2820	4136
10.0	190	275	250	365	1500	2200	19.0	355	525	475	695	2850	4180
10.2	190	280	255	375	1530	2244	19.2	360	530	480	705	2880	4224
10.4	195	285	260	380	1560	2288	19.4	365	535	485	710	2910	4268
10.6	200	290	265	390	1590	2332	19.6	370	540	490	720	2940	4312
10.8	205	295	270	395	1620	2376	19.8	370	545	495	725	2970	4356
							20.0	375	550	500	735	3000	4400

* Volumes per feed are rounded to the nearest 5 ml.



Reference Tables for Infants 0-6 Months Old and Older Infants weighing less than 3.0kg

Reference Table for Maintenance Amounts of F-100-Diluted to Give to an Individual Infant per Feed

Bodyweight (kg)	F-100-Diluted (ml per feed) (assumes 8 feeds per day, given 3-hourly)
≥ 1.2	25
1.3–1.5	30
1.6–1.7	35
1.8–2.1	40
2.2–2.4	45
2.5–2.7	50
2.8–2.9	55
3.0–3.4	60
3.5–3.9	65
4.0–4.4	70

Reference Table for Amounts of F-100-Diluted (Marasmus) or F-75 (Kwashiorkor) to Give to Non-breastfed Infants in the Stabilisation Phase

Bodyweight (kg)	F-100-Diluted or F-75 (ml per feed), 8 feeds per day, no breastfeeding (3-hourly feeds)
≤ 1.5	30
1.6–1.8	35
1.9–2.1	40
2.2–2.4	45
2.5–2.7	50
2.8–2.9	55
3.0–3.4	60
3.5–3.9	65
4.0–4.4	70

Reference Table for Amounts of F-100-Diluted to Give to Non-breastfed Infants 0–6 Months or Older Infants Weighing Less Than 3.0 kg in the Transition Phase

Bodyweight (kg)	F-100-Diluted (ml per feed), 8 feeds per day, no breastfeeding (3-hourly feeds)
≤ 1.5	45
1.6–1.8	53
1.9–2.1	60
2.2–2.4	68
2.5–2.7	75
2.8–2.9	83
3.0–3.4	90
3.5–3.9	96
4.0–4.4	105

Reference Table for Amounts of F-100-Diluted to Give to Non-breastfed Infants 0–6 Months or Older Infants Weighing Less Than 3.0 kg in the Rehabilitation Phase

Bodyweight (kg)	F-100-Diluted (ml per feed), 6 to 8 feeds per day, no breastfeeding
≤ 1.5	60
1.6–1.8	70
1.9–2.1	80
2.2–2.4	90
2.5–2.7	100
2.8–2.9	110
3.0–3.4	120
3.5–3.9	130
4.0–4.4	140



Stabilisation Phase Reference Tables for F-75 for Children with Severe Wasting (Marasmus)

Weight of child (kg)	Volume of F-75 per feed (ml) ^a			Daily total (130 ml/kg)	80% of daily total ^a (minimum)
	Every 2 hours ^b (12 feeds)	Every 3 hours ^c (8 feeds)	Every 4 hours (6 feeds)		
2.0	20	35	45	130	260
2.2	25	35	50	130	286
2.4	25	40	50	130	312
2.6	30	40	55	130	338
2.8	30	45	60	130	364
3.0	35	50	65	130	390
3.2	35	50	70	130	416
3.4	35	55	75	130	442
3.6	40	60	80	130	468
3.8	40	60	80	130	494
4.0	45	65	85	130	520
4.2	45	70	90	130	546
4.4	50	70	95	130	572
4.6	50	75	100	130	598
4.8	50	80	105	130	624
5.0	55	80	110	130	650
5.2	55	85	115	130	676
5.4	60	90	115	130	702
5.6	60	90	120	130	728
5.8	65	95	125	130	754
6.0	65	100	130	130	780
6.2	65	100	135	130	806
6.4	70	105	140	130	832
6.6	70	105	145	130	858
6.8	75	110	145	130	884
7.0	75	115	150	130	910
7.2	80	115	155	130	936
7.4	80	120	160	130	962
7.6	80	125	165	130	988
7.8	85	125	170	130	1014
8.0	85	130	175	130	1040
8.2	90	135	180	130	1066
8.4	90	135	180	130	1092
8.6	95	140	185	130	1118
8.8	95	145	190	130	1144
9.0	100	145	195	130	1170
9.2	100	150	200	130	1196
9.4	100	155	205	130	1222
9.6	105	155	210	130	1248
9.8	105	160	210	130	1274
10.0	110	165	215	130	1300
10.2	110	165	220	130	1326
10.4	115	170	225	130	1352
10.6	115	170	230	130	1378
10.8	115	175	235	130	1404
11.0	120	180	240	130	1430

Weight of child (kg)	Volume of F-75 per feed (ml) ^a			Daily total (130 ml/kg)	80% of daily total ^a (minimum)
	Every 2 hours ^b (12 feeds)	Every 3 hours ^c (8 feeds)	Every 4 hours (6 feeds)		
11.2	120	180	245	130	1456
11.4	125	185	245	130	1482
11.6	125	190	250	130	1508
11.8	130	190	255	130	1534
12.0	130	195	260	130	1560
12.2	130	200	265	130	1586
12.4	135	200	270	130	1612
12.6	135	205	275	130	1638
12.8	140	210	275	130	1664
13.0	140	210	280	130	1690
13.2	145	215	285	130	1716
13.4	145	220	290	130	1742
13.6	145	220	295	130	1768
13.8	150	225	300	130	1794
14.0	150	230	305	130	1820
14.2	155	230	310	130	1846
14.4	155	235	310	130	1872
14.6	160	235	315	130	1898
14.8	160	240	320	130	1924
15.0	165	245	325	130	1950
15.2	165	245	330	130	1976
15.4	165	250	335	130	2002
15.6	170	255	340	130	2028
15.8	170	255	340	130	2054
16.0	175	260	345	130	2080
16.2	175	265	350	130	2106
16.4	180	265	355	130	2132
16.6	180	270	360	130	2158
16.8	180	275	365	130	2184
17.0	185	275	370	130	2210
17.2	185	280	375	130	2236
17.4	190	285	375	130	2262
17.6	190	285	380	130	2288
17.8	195	290	385	130	2314
18.0	195	295	390	130	2340
18.2	195	295	395	130	2366
18.4	200	300	400	130	2392
18.6	200	300	405	130	2418
18.8	205	305	405	130	2444
19.0	205	310	410	130	2470
19.2	210	310	415	130	2496
19.4	210	315	420	130	2522
19.6	210	320	425	130	2548
19.8	215	320	430	130	2574
20.0	215	325	435	130	2600

^a Volumes in these columns are rounded to the nearest 5 ml.

^b Give 2-hourly feeds for at least the first day. When there is little or no vomiting, moderate diarrhoea (< 5 watery stools per day), and the child finishes most feeds, change to 3-hourly feeds.

^c After a day on 3-hourly feeds, if there is no vomiting, less diarrhoea, and the child finishes most feeds, change to 4-hourly feeds.



Stabilisation Phase Reference Tables for F-75 for Children with Bilateral Pitting Oedema (+++)

Weight with +++ oedema (kg)	Volume of F-75 per feed (ml) ^a			Daily total (100 ml/kg)	80% of daily total (minimum)
	Every 2 hours ^b (12 feeds)	Every 3 hours ^c (8 feeds)	Every 4 hours (6 feeds)		
3.0	25	40	50	100	300
3.2	25	40	55	100	320
3.4	30	45	55	100	340
3.6	30	45	60	100	360
3.8	30	50	65	100	380
4.0	35	50	65	100	400
4.2	35	55	70	100	420
4.4	35	55	75	100	440
4.6	40	60	75	100	460
4.8	40	60	80	100	480
5.0	40	65	85	100	500
5.2	45	65	85	100	520
5.4	45	70	90	100	540
5.6	45	70	95	100	560
5.8	50	75	95	100	580
6.0	50	75	100	100	600
6.2	50	80	105	100	620
6.4	55	80	105	100	640
6.6	55	85	110	100	660
6.8	55	85	115	100	680
7.0	60	90	115	100	700
7.2	60	90	120	100	720
7.4	60	95	125	100	740
7.6	65	95	125	100	760
7.8	65	100	130	100	780
8.0	65	100	135	100	800
8.2	70	105	135	100	820
8.4	70	105	140	100	840
8.6	70	110	145	100	860
8.8	75	110	145	100	880
9.0	75	115	150	100	900
9.2	75	115	155	100	920
9.4	80	120	155	100	940
9.6	80	120	160	100	960
9.8	80	125	165	100	980
10.0	85	125	165	100	1000
10.2	85	130	170	100	1020
10.4	85	130	175	100	1040
10.6	90	135	175	100	1060
10.8	90	135	180	100	1080

Weight with +++ oedema (kg)	Volume of F-75 per feed (ml) ^a			Daily total (100 ml/kg)	80% of daily total (minimum)
	Every 2 hours ^b (12 feeds)	Every 3 hours ^c (8 feeds)	Every 4 hours (6 feeds)		
12.0	100	150	200	100	1200
12.2	100	155	205	100	1220
12.4	105	155	205	100	1240
12.6	105	160	210	100	1260
12.8	105	160	215	100	1280
13.0	110	165	215	100	1300
13.2	110	165	220	100	1320
13.4	110	170	225	100	1340
13.6	115	170	225	100	1360
13.8	115	175	230	100	1380
14.0	115	175	235	100	1400
14.2	120	180	235	100	1420
14.4	120	180	240	100	1440
14.6	120	185	245	100	1460
14.8	125	185	245	100	1480
15.0	125	190	250	100	1500
15.2	125	190	255	100	1520
15.4	130	195	255	100	1540
15.6	130	195	260	100	1560
15.8	130	200	265	100	1580
16.0	135	200	265	100	1600
16.2	135	205	270	100	1620
16.4	135	205	275	100	1640
16.6	140	210	275	100	1660
16.8	140	210	280	100	1680
17.0	140	215	285	100	1700
17.2	145	215	285	100	1720
17.4	145	215	290	100	1740
17.6	145	220	295	100	1760
17.8	150	225	295	100	1780
18.0	150	225	300	100	1800
18.2	150	230	305	100	1820
18.4	155	230	305	100	1840
18.6	155	235	310	100	1860
18.8	155	235	315	100	1880
19.0	160	240	315	100	1900
19.2	160	240	320	100	1920
19.4	160	240	325	100	1940
19.6	165	245	325	100	1960
19.8	165	250	330	100	1980
20.0	165	250	335	100	2000

^a Volumes in these columns are rounded to the nearest 5 ml.

^b Give 2-hourly feeds for at least the first day. When there is little or no vomiting, moderate diarrhoea (< 5 watery stools per day), and the child finishes most feeds, change to 3-hourly feeds.

^c After a day on 3-hourly feeds, if there is no vomiting, less diarrhoea, and the child finishes most feeds, change to 4-hourly feeds.