

FINAL

PREVENTION IS BETTER THAN CURE.

Final Report of the Evaluation:

Prevention or Cure?

**Comparing Preventive and Recuperative Approaches to Targeting Maternal
and Child Health and Nutrition Programs in Rural Haiti**

International Food Policy Research Institute - Cornell University – World Vision Haiti Team

Written by:

Purnima Menon, Cornell University
Marie T. Ruel, International Food Policy Research Institute (IFPRI)

With contributions from (in alphabetical order)

Mary Arimond, IFPRI
Jean-Pierre Habicht, Cornell University
Bekele Hankebo, WV-Haiti
Cornelia Loechl, International Potato Center
John Maluccio, Middlebury College
Mduduzi Mbuya, Cornell University
Lesly Michaud, WV-Haiti
Gretel Pelto, Cornell University

Submitted to:
The Food and Nutrition Technical Assistance (FANTA) Project

July 31, 2007

This report was made possible through the generous support of the American people through the support of the Office of Health, Infectious Disease and Nutrition, Bureau for Global Health, and the Office of Food for Peace, Bureau for Democracy, Conflict and Humanitarian Assistance, United States Agency for International Development (USAID), under terms of Cooperative Agreement No. HRN-A-00-98-00046-00, through the Food and Nutrition Technical Assistance (FANTA) project operated by the Academy for Educational Development (AED). The contents are the responsibility of AED and do not necessarily reflect the views of the USAID or the United States Government.

Financial support for this research is also provided by the Government of Germany, WV-Haiti, and the World Food Programme.

RECOMMENDED CITATION:

Menon, P., and M. T. Ruel, with (*in alphabetical order*) M. Arimond, J.-P. Habicht, B. Hankebo, C. Loechl, J. Maluccio, M. N. Mbuya, and G. Pelto. *Prevention Is Better than Cure. Final Report of the Evaluation: Prevention or Cure? Comparing Preventive and Recuperative Approaches to Targeting Maternal and Child Health and Nutrition Programs in Rural Haiti*. Submitted to the Food and Nutrition Technical Assistance Project, Academy for Educational Development, Washington, D.C. July 2007.



ACKNOWLEDGMENTS

This three-year evaluation has been the culmination of the efforts and commitment of numerous people and organizations. We begin by expressing our gratitude to key individuals who played critical roles in this evaluation, and then acknowledge all the other individuals and groups who have contributed to the success of this study.

First, we acknowledge the critical support from **Gilles Bergeron** at the Food and Nutrition Technical Assistance (FANTA) Project, Academy for Educational Development (AED), who conceived of this study and saw it through from the early stages to the very end. In addition to his roles at FANTA for coordinating this work, Gilles has made substantial intellectual contributions to the evaluation throughout the study. In addition, he was instrumental in connecting the IFPRI-Cornell evaluation group with numerous other individuals and organizations in Haiti, all of which have strengthened this study.

Second, we are grateful to the senior management team at WV-Haiti, particularly **Wesley Charles, Bekele Hankebo, and Lesly Michaud**. All of them were involved in this study from its inception. Their willingness to engage with the research team at all stages of this evaluation, to fund the fieldwork for the study, as well as to invest in the development of the preventive approach are all indicative of a commitment to improving nutrition among Haitian children. In addition, their commitment provides an example of an agency that aims to improve staff performance and program effectiveness by developing and supporting a motivational work context for staff at all levels.

Third, we thank **Eunyong Chung** of USAID/Washington for her continued support and engagement with this evaluation and for the critical insights and inputs she provided at various junctures. We also are grateful to **Carell Laurent**, who was based at the Food for Peace Office at USAID/Port-Au-Prince at the inception of this study. Carell galvanized support for this study at USAID/ Port-Au-Prince and among other stakeholders in Haiti and was a driving force in seeing this study materialize in Haiti. Florence Cadet – also from USAID/Port-au-Prince – also contributed significantly and helped ensure continued financial and logistical support from USAID/Port-au-Prince.

Finally, we are grateful for the many contributions of all those who were involved with this study, whether in the form of intellectual, financial, in-kind, logistical, and personal support. Here, we acknowledge these many forms of contribution by naming those people and organizations who we engaged with through the course of this evaluation.

Any omissions are sincerely regretted.

- ❖ **FANTA:** Gilles Bergeron, Bruce Cogill, Anne Swindale, Megan Deitchler
- ❖ **WV-Haiti:** Lesly Michaud, Bekele Hankebo, Wesley Charles, Frank Williams, John Dorman, Jean-Marie Boisrond, Rony Destine, Rulisa Kimenyi, Makayi Chinyam, Kassa Selfago, Nixon Desir, Emmanuel Tinord, Rafael Cambroner, Cedrelle Jean Louis, Harry Francois, Lionel Isaac, Maude St. Victor

- ❖ **WV-Haiti Regional office at Hinche:** Widy Laroque, Jean Louis Pierre, Dorlean Delince, Wagnes Desir, Ruth Calvaire, Marie-Carmel Joseph, Charlot Willelmyne, Delide Jean Baptiste, Elisabeth Elysée, Jean Robert Francoeur, Wisler Nicolas, Nuclair Guillaume, as well as all nurses, health promoters, assistant health promoters, and food monitors at the regional office
- ❖ **USAID/Port-au-Prince:** Florence Cadet, Michael Kerst, Carell Laurent
- ❖ **Ministry of Health (Haiti):** Dr. Francesca Joceline Marhone Pierre, Head of the National Coordination Office of the Program on Nutrition
- ❖ **Haiti-based research staff:** Elisabeth Metellus, Arsene Ferrus, Remy Lafalaise, Pierre Lenz Dominique, Remy Lafalaise, Mathieu Honoré, Louise Dumay, Josianne Loredan, Slande Celestre, Yves Francois-Pierre, as well the survey fieldworkers and team chauffeurs
- ❖ **Collaborators in Haiti:** Edouine Francois (Freedom from Hunger), DidacArts, Paul Thomas, Paul Brea, and Michelle Cayemittes (Insitut Haitien de l'Enfance,), Colette Vilgrain (IntellConsult), Agathes Pellerin (CRS), Sonia Stines (CRS), Jude-Marie Banatte (CRS), Marjolein Moreaux (Save the Children, US), Mireille Sylvain (CARE), Bette Gebrian (Haitian Health Foundation)
- ❖ **International Food Policy Research Institute (IFPRI):** Wahid Quabili, Ali Subandoro, Jay Willis, Suneetha Kadiyala, Ginette Mignot, Lynette Aspillera, Celine Castillo-Macy
- ❖ **Cornell University:** Kate Dickin, Edward Frongillo, Francoise Vermeylen, Amanda Zongrone

In closing, we acknowledge the communities, families, and women who were interviewed over the course of this study, from the formative research for the development of the intervention to the large surveys that provided the impact data. Our research teams were always welcomed by these people of the Central Plateau of Haiti, and for this we express our sincere thanks.

ACRONYMS

ADP	Area Development Program
AED	Academy for Educational Development
AHPs	Assistant Health Promoters
ARI	Acute Respiratory Infection
BCC	Behavior Change Communication
BCG	Bacille Calmette Guerin (vaccine)
BF	Breastfeeding
BMI	Body Mass Index
CBR	Cost-benefit ratio
CF	Complementary Feeding
CER	Cost-effectiveness ratio
CIF	Cost, Insurance, and Freight (prices)
CRSP	Collaborative Research Support Program
CS	Cooperating Sponsors
CSB	Corn-Soy Blend
CTS	Commodity Tracking System
CU	Cornell University
DAP	Development Activity Program
DD	Dietary Diversity
DTP	Diphtheria, Tetanus, and Pertussis (vaccine)
EBF	Exclusive Breastfeeding
EMMUS	<i>Enquête Mortalité Morbidité et Utilisation des Services</i>
FANTA	Food and Nutrition Technical Assistance
FDPs	Food Distribution Points
FES	Focused Ethnographic Study
FFH	Freedom from Hunger
FOB	Free on Board (prices)

FY	Fiscal Year
GDP	Gross Domestic Product
GMP	Growth Monitoring and Promotion
HAZ	Height-for-Age Z-score
HH	Household
HPs	Health Promoters
IHE	<i>Institut Haïtien de l'Enfance</i>
IFPRI	International Food Policy Research Institute
LAC	Latin American and the Caribbean
LAM	Lactational Amenorrhea Method
LSMS	Living Standards Measurement Survey
MCs	Mothers' Clubs
MCHN	Maternal and Child Health and Nutrition
OR	Operations Research
ORS	Oral Rehydration Salts
PVOs	Private Voluntary Organizations
RP	Rally Posts
SD	Standard Deviation
SFB	Soy-Fortified Bulgur
SSS	Salt-Sugar Solution
TIPS	Trials of Improved Practices
USAID	United States Agency for International Development
USD	U.S. dollar
WAZ	Weight-for-Age Z-score
WHZ	Weight-for-Height Z-score
WSB	Wheat-Soy Blend
WV	WV-Haiti

TABLE OF CONTENTS

ACKNOWLEDGMENTS	i
ACRONYMS.....	iii
EXECUTIVE SUMMARY	xvii
1. INTRODUCTION AND SCIENTIFIC RATIONALE FOR THE STUDY	1
1.1 Haiti: The Poorest Country in the Western Hemisphere	2
1.2 Title II-MCHN Programs.....	2
1.3 Rationale for a Preventive Approach to Reducing Undernutrition.....	3
1.3.1 The recuperative versus the preventive approach.....	3
1.3.2 Biological rationale for targeting food supplementation to children under 24 months	4
1.3.3 Behavioral rationale for combining food supplementation and behavior change communication strategies.....	7
1.3.4 Conclusions.....	10
1.4 Organization of the Report.....	11
2. OBJECTIVES AND DESIGN OF THE EVALUATION.....	13
2.1 Introduction.....	13
2.2 Objectives of the Evaluation.....	13
2.2.1 Additional objectives	13
2.3 Design of the Overall Evaluation.....	14
2.3.1 Program theory and pathways of influence	14
2.3.2 Designing evaluation activities based on program theory	17
2.4 Project Activities.....	19
2.4.1 Evaluation activities.....	21
2.5 Conclusions.....	26
3. WV-HAITI'S MCHN PROGRAM	27
3.1 Introduction.....	27
3.2 Services Provided.....	27
3.3 Program Service Delivery Points.....	29
3.3.1 Rally Post services	29
3.3.2 Mothers' Club services	30
3.3.3 Food Distribution Point (FDP) services.....	31
3.3.4 Pre-and postnatal consultations.....	32
3.3.5 Home visits	32
3.4 Program Organization/Staffing.....	32
3.5 Conclusions.....	34

4. GETTING IT RIGHT FROM THE START: DESIGNING THE PREVENTIVE AND RECUPERATIVE APPROACHES.....	35
4.1 Introduction.....	35
4.2 The Food Assistance Component	35
4.3 The BCC Component.....	36
4.3.1 Research phase.....	36
4.3.2 Development phase.....	39
4.4 The Implementation Plan for the Preventive and Recuperative Program Approaches	44
4.4.1 At Rally Posts (RPs)	44
4.4.2 At the Mothers' Clubs (MCs)	46
4.4.3 Food Distribution Points (FDP).....	49
4.5 Conclusions.....	49
5. DELIVERING IT WELL: THE OPERATIONS RESEARCH PROCESS	51
5.1 Introduction.....	51
5.1.1 Integration of operations research into implementation and evaluation processes.....	51
5.2 Objectives	53
5.3 Methods.....	54
5.3.1 Program context and stakeholders	54
5.3.2 Data collection methods.....	54
5.3.3 Consultative workshop.....	56
5.4 Results.....	57
5.4.1 First round of operations research (OR1)	57
5.4.2 Outcomes of consultative workshop.....	61
5.4.3 Second round of operations research (OR2).....	63
5.5 Conclusions.....	73
6. PROGRAM PARTICIPATION AND HOUSEHOLD AND CAREGIVER CHARACTERISTICS	75
6.1 Introduction.....	75
6.2 Data and Analysis	75
6.2.1 Data and variables.....	75
6.2.2 Analysis.....	76
6.3 Results.....	76
6.3.1 Program participation.....	76
6.3.2 Household and caregiver characteristics.....	82
6.4 Summary of Results.....	86

7. NUTRITIONAL IMPACT OF THE PREVENTIVE APPROACH COMPARED TO THE RECUPERATIVE APPROACH	87
7.1 Introduction.....	87
7.2 Objective of the Impact Evaluation	87
7.3 Intervention and Comparison Groups	87
7.4 Methods.....	88
7.4.1 Evaluation design.....	88
7.4.2 Data analysis	88
7.5 Results.....	89
7.5.1 Baseline characteristics.....	89
7.5.2 Intervention impact: Results from probability design analysis.....	90
7.5.3 Results of plausibility analysis.....	91
7.5.4 Differences in morbidity between the two program groups	96
7.6 Summary of Results.....	98
7.6.1 Probability results	98
7.6.2 Plausibility results.....	99
7.7 Conclusions.....	99
8. IMPACT OF THE PROGRAM ON KNOWLEDGE AND PRACTICES	101
8.1 Introduction.....	101
8.2 Data and Variable Creation.....	101
8.2.1 Maternal knowledge.....	101
8.2.2 Trial and adoption of key recommended practices	104
8.2.3 Child-feeding and care practices.....	105
8.2.4 Other care practices: Preventive care, hygiene practices, and care during illness.....	108
8.3 Analysis.....	109
8.4 Results.....	109
8.4.1 Impact of the program on maternal knowledge	109
8.4.2 Trial and adoption of key recommended practices	114
8.4.3 Impact of intervention on infant and young child-feeding and care practices	121
8.4.4 Impact of interventions on preventive care, care during illness, and hygiene.....	133
8.4.5 Comparisons between participants and nonparticipants on child-feeding and care practices.....	136
8.5 Conclusions.....	141
9. IMPACT OF THE PROGRAM ON FOOD SECURITY	143
9.1 Introduction.....	143
9.2 Data and Analysis	143
9.2.1 Data and variables.....	143
9.2.2 Analysis.....	144

9.3 Results.....	145
9.3.1 Impact of the Program on Household Food Insecurity	145
9.4 Conclusions.....	151
10. INDIRECT BENEFITS OF THE PROGRAM.....	153
10.1 Introduction.....	153
10.2 Data and Analysis	153
10.2.1 Data and variables.....	153
10.2.2 Analysis.....	155
10.3 Results.....	156
10.3.1 Impact of the program on household and respondent asset ownership	156
10.3.2 Indirect impact of the program on women’s well-being.....	158
10.4 Conclusions.....	162
11. COST-EFFECTIVENESS OF THE PREVENTIVE APPROACH RELATIVE TO THE RECUPERATIVE APPROACH	163
11.1 Introduction.....	163
11.2 Methodology	164
11.2.1 The incremental cost-effectiveness between the two program approaches.....	164
11.2.2 Measuring the costs.....	164
11.3 Data: Costs in the Program Areas.....	169
11.3.1 Direct and outside program costs in the study areas.....	169
11.3.2 Opportunity costs in the study areas	174
11.4 Cost-Effectiveness of Preventive versus Recuperative.....	175
11.4.1 Incremental cost-effectiveness estimates	175
11.4.2 Interpreting the incremental cost-effectiveness estimates	177
11.5 Conclusions.....	178
12. KEY FINDINGS, AND PROGRAM AND RESEARCH IMPLICATIONS	181
12.1 Relative Impact and Cost-Effectiveness	181
12.1.1 Is the preventive approach more effective than the recuperative approach at reducing childhood undernutrition?	181
12.1.2 What is the relative cost-effectiveness of the preventive and recuperative approaches?.....	182
12.2 Limitations of the Study.....	184
12.3 Pathways of Impact.....	185
12.3.1 Factors pertaining to program design	185
12.3.2 Factors pertaining to program delivery.....	187
12.3.3 Factors pertaining to program access and utilization.....	189
12.3.4 Conclusions on pathways of impact	195
12.4 Implications for Programs, Policies, and Future Research	196
12.4.1 Implications for programs and policies.....	196
12.4.2 Research implications	198

12.5 Lessons for Program Evaluations	199
12.6 Conclusions.....	200
REFERENCES	201
Annex 2.1 List of household survey modules.....	209
Annex 3.1 Organizational structure of the Health component of the program.....	211
Annex 3.2 Organizational structure of the Commodities component of the program.....	213
Annex 4.1 Infant and child feeding practices in Haiti compared to best practices, and constraints and opportunities for behavior change in Central Plateau	215
Annex 4.2 Identification of programmatic options to address the constraints to infant feeding, and to support facilitating factors	219
Annex 4.3 BCC strategy matrix for a BCC program to prevent malnutrition among children between 0-24 months	223
Annex 4.4 Schedules of learning sessions and topics at Mothers' Clubs.....	227
Annex 5.1 Summary matrix from Consultative Workshop, February 2003.....	231
Annex 5.2 Mothers' Clubs observation checklist items	237
Annex 5.3 Questions for staff related to motivation.....	239
Annex 5.4 Questions for staff related to supervision.....	241
Annex 8.1 Scoring of questions in Infant and Young Child Feeding Knowledge Test.....	243
Annex 8.2 Scoring of questions in General Nutrition and Health knowledge Test (responses that were considered appropriate)	245
Annex 8.3a Guiding principles for complementary feeding of the breastfed child.....	249
Annex 8.3b Guiding Principles for feeding non-breastfed children 6-24 months of age	251
Annex 8.4 Awareness, trial, and adoption of key recommended practices, by program group and participation (ever).....	255
Annex 8.5 Percent of infants fed according to recommendations during early postnatal period, by intervention area and by participation	257
Annex 8.6 Age of introduction of selected liquids and foods, by intervention area and by participation	259

Annex.8.7 Percent of children who were fed at least the minimum recommended number of times, and mean number of meals and snacks, by age group, current program participation, and breastfeeding status	261
Annex 8.8 Percent of children who consumed selected food groups in the previous 24 hours, by intervention area and participation.....	263
Annex 8.9 Percent of children who consumed selected animal source foods in the previous 24 hours, by intervention area and participation.....	265
Annex 8.10 Frequency of consumption of nutrient-rich foods, including recipes promoted in mothers' clubs, by intervention area and participation (mother ever participated)	267
Annex 8.11 Frequency of consumption of nutrient-rich foods, including recipes promoted in mothers' clubs, by intervention area and current participation (children 6-42 months)	269
Annex 8.12 Mean dietary diversity, by age group, intervention area, and participation (children 6-42 months).....	271
Annex 8.13 Receipt of vitamin and mineral supplements, by index children (12-41 months) and their mothers	273
Annex 8.14 Feeding during and after diarrhea: Index children	275
Annex 8.15 Immunization status, by intervention group and by participation (among children 12-41 months of age)	277
Annex 9.1 Scoring of variables for the food insecurity scale	279
Annex 10.1 Scoring of women's well-being scales.....	281
Annex 12.1 Summary of evaluation results	283

LIST OF TABLES

2.1 Summary of study activities.....	20
3.1 Services provided by the WV MCHN program in Central Plateau	28
3.2 Composition of direct and indirect food rations, per beneficiary category	31
4.1 Package of interventions provided in the recuperative and preventive approaches	45
5.1 Data collection methods used in the OR process.....	55

5.2	Types of information communicated to mothers during growth monitoring at Rally Posts, by year, by program area, and by research method	66
5.3	Types of advice given to mothers at Rally Posts in 2004, by child's progress.....	67
5.4	Advice given at Rally Posts in 2004, by child's progress and program area.....	67
5.5	Quality of information and facilitation at Mothers' Clubs, by year and by program area	68
5.6	Summary of program changes in WV-Haiti MCHN program, and differences between preventive and recuperative program areas in 2004	69
6.1	Program participation at the <i>household</i> level (includes participation by all children in the household, as well as pregnant/lactating women).....	77
6.2	Enrollment of index children in the food assistance program, by program group	78
6.3	Timing of exposure for index children, by program group	78
6.4	Receipt of food assistance among index children, by child age and program participation.....	79
6.5	Participation in Mothers' Clubs and use of Rally Post services	80
6.6	RP participation, by child age and program group	81
6.7	Program uptake and targeting, by program group	81
6.8	Household characteristics, by program and by program participation (ever participated)....	83
6.9	Caregiver/respondent characteristics, by program group and by program participation (ever participated)	85
7.1	Key differences between recuperative and preventive approach intervention packages.....	87
7.2	Comparison of child, maternal, and household characteristics between program groups at baseline.....	90
7.3	Mean anthropometric outcomes at final survey, by program group	91
8.1	Guiding principles and recall data collected on recommended practices	106
8.2	Maternal nutrition and health knowledge, by program group	111
8.3	Maternal nutrition and health knowledge, by program group and participation (ever participated)	112
8.4	Program impact on maternal nutrition and health knowledge (regression analysis)	113

8.5 Differences in trial of recommended practices between preventive and recuperative groups and reasons for non-trial of practices, by program exposure	115
8.6 Child-feeding and care practices, by program group and time of survey	122
8.7 Preventive care, care during illness, and hygiene, by program group and time of survey, for index children 12 to 42 months of age	134
8.8 Summary of findings on behavior change outcomes	137
9.1 Household food insecurity, by program and time of survey.....	145
9.2 Selected household food insecurity experiences, by program and time of survey	146
9.3 All household food insecurity measures at final survey, by participation (ever and current)	146
9.4 Program impact on household food insecurity experiences at final survey (regression analysis).....	148
9.5 Program impact on months of inadequate household food provisioning (MIHFP) at final survey (regression analysis).....	149
9.6 Program impact on severity of months of inadequate household food provisioning (MIHFP) at final survey	149
10.1 Household and women’s asset ownership, by program group, participation, and time of the survey.....	157
10.2 Respondent mental well-being, by program group, participation, and time of survey	159
10.3 Program impact on self-rated health (random effects regression)	160
10.4 Program impact on mental stress scale (presence/absence of symptoms) (random effects regression)	161
11.1 Variables and data sources for incremental cost-effectiveness analysis.....	165
11.2 Number of program beneficiary-months in Central Plateau and study areas	170
11.3 Number of program child beneficiaries per month in the Central Plateau and study areas (FY 2005).....	172
11.4 Direct program and outside program costs in the study areas (\$000).....	173
11.5 Direct program and outside program costs per beneficiary-month in the study areas (\$)	174
11.6 Incremental cost-effectiveness	176

LIST OF FIGURES

1.1 Mean weight-for-height (WHZ), weight-for-age (WAZ), and height-for-age (HAZ) of rural children in Haiti (EMMUS-III 2001)	5
2.1 Program impact pathways.....	16
2.2 An “engaged” model of evaluation and technical assistance.....	18
3.1 Flow of activities at the Rally Post	30
4.1 Steps of the design process of the BCC component	37
5.1 An “engaged” model of evaluation and program development.....	52
7.1 Prevalence of undernutrition among children 12-41 months at baseline (groups combined) and end line, by program group.....	92
7.2 Mean anthropometric outcomes by child age and program group at final survey.....	93
7.3 Prevalence of stunting, underweight and wasting by child age and program group at final survey.....	93
7.4 Weight-for-age Z-scores of index children (12-41 mo) and their younger siblings (0-11), by age and program group at final survey	95
7.5 Percentage of children who had fever in the past two weeks, by age and program group (final survey).....	96
7.6 Percentage of children with cold/cough in the past two weeks, by age and program group (final survey).....	97
7.7 Percentage of children with diarrhea in the past two weeks, by age and program group (final survey).....	97
7.8 Percentage of children with fast breathing in the past two weeks, by age and program group (final survey).....	98
8.1a From awareness to adoption: Leaving expressed breast milk behind when going out.....	116
8.1b From awareness to adoption: Adding breast milk to gruel.....	116
8.1c From awareness to adoption: Adding an egg to child’s gruel.....	117
8.1d From awareness to adoption: Preparing gruel with beans and nuts.....	117
8.1e From awareness to adoption: Feeding enriched gruel in the evening.....	118

8.1f From awareness to adoption: Preparing mashed plantain with pumpkin.....	118
8.1g From awareness to adoption: Feeding an extra meal after illness	119
8.2 Percent of mothers who reported optimal early feeding practices during first day of life, by program group and baseline and final survey—Index child.....	124
8.3 Percent of mothers who reported optimal early feeding practices during first day of life, by program group and baseline and final survey—Younger siblings	125
8.4 Comparing baseline and final sample exclusive breastfeeding in past 24 hours for infants < 6 months: Kaplan-Meier survival functions (preventive/recuperative groups combined).....	125
8.5 Maternal recall of timing of introduction of liquids and foods, at baseline and final survey (preventive/recuperative groups combined)	127
8.6 Percent of breastfed children having received complementary foods minimum recommended times yesterday, by age, program group, and time of survey.....	129
8.7 Percent of non-breastfed index children fed complementary foods a minimum recommended times, by age, program group, and time of survey	130
8.8 Percent of index children who consumed selected food groups in previous 24 hours, by program group, and baseline and final survey	131
9.1 Household food insecurity experiences score, by program group and current participation at final survey	147
9.2 Prevalence of severe food insecurity (based on terciles of the household food insecurity experiences score), by program group and current participation at final survey.....	147
9.3 Months of inadequate household food provisioning (MIHFP), by program group and current participating at final survey	148
9.4 Proportion of households with insufficient food in each month of the calendar year, by program group (derived from the months of inadequate household food provisioning (MIHFP) measure).....	150
12.1 Program impact pathways.....	186

LIST OF BOXES

3.1 Five service delivery venues for the WV-Haiti food-assisted MCHN program.....	27
4.1 Example from decision matrix to organize formative research results on exclusive breastfeeding.....	39
4.2 Example from program-planning matrix to address constraints to exclusive breastfeeding	40
4.3 Example of BCC strategy planning matrix on expression of breast milk.....	40
4.4 Example of message adaptation for expression of breast milk.....	42
4.5 Example of learning session on expression of breast milk	42
4.6 Mothers' Club learning sessions.....	48
5.1 Example of prioritization matrix used at the consultative workshop	57
8.1 Knowledge test questions on topics covered at MCs and RPs	103
8.2 From awareness to trial and adoption: Key practices included in the survey.....	105

EXECUTIVE SUMMARY

Background

Food-assisted maternal and child health and nutrition (MCHN) programs have traditionally targeted children less than 5 years of age who were identified as undernourished through growth monitoring activities. Scientific evidence, however, shows that children under two are more at risk of becoming undernourished, and more responsive to nutrition interventions than older children. Research findings consistently show that the earlier and the longer food supplementation is provided before the child reaches two years of age, the greater the benefits not only on growth in early life, but also on long-term physical, cognitive, and reproductive performance. Thus, investing in the first two years of life provides benefits way beyond childhood and is an essential element of development strategies and human capital formation interventions.

Although the benefits of intervening early in life are unequivocal, nutrition-focused programs globally have been hesitant to adopt the universal targeting of children under two because of the lack of evidence of the effectiveness of the approach in large-scale programs. In order to fill this knowledge gap, researchers from the International Food Policy Research Institute and Cornell University undertook a study commissioned by the Food and Nutrition Technical Assistance (FANTA) Project at the Academy for Educational Development (AED) to compare the effectiveness and cost-effectiveness of a “preventive” approach, which targets all children under two, with the traditional “recuperative” approach, which targets children under five years of age once they have become undernourished. The project was carried out in collaboration with World Vision-Haiti (WV-Haiti) in the context of their PL 480 Title II food-assisted MCHN program in the Central Plateau.

Study Objectives

The main objective of the study was to compare, under programmatic conditions, the impact on childhood undernutrition of targeting food assistance and behavior change communication (BCC) in the context of a Title II MCHN program to all children 6-24 months of age (preventive approach), versus to undernourished children less than 5 years of age (recuperative approach). We also compared the cost and cost-effectiveness of the two program approaches, as well as their impact on a variety of intermediate outcomes, such as food security, maternal nutrition knowledge, and infant feeding practices.

Intervention Packages

The preventive and recuperative packages of nutrition services were designed based on the best available biological evidence about the efficacy of supplementation as well as formative research to develop the BCC strategy that would accompany a monthly food assistance ration. The composition and size of the monthly food ration was identical for both program approaches, but the programs differed in the following three aspects: (1) the eligibility criterion (undernutrition among under-five children for recuperative; and age 6-24 months for preventive); (2) the focus, timing, sequencing, and number of sessions of BCC (see below); and (3) the timing and duration of eligibility to receive the food and BCC intervention (nine months from time child

is identified as being undernourished for recuperative – as decided by WV-Haiti for this approach; the whole 18-months period when children are between 6 and 24 months for preventive). Severely malnourished children (weight-for-age Z-scores < -3) 24 months or older in the preventive approach were also eligible to participate in the program for nine months. Both program approaches targeted pregnant and lactating women, who were eligible to receive a monthly food ration until the sixth month of lactation.

Receipt of the food ration was conditional on regular monthly attendance at other program health services, including the Rally Posts (which provide preventive health-care services), pre- and postnatal consultations (for pregnant and lactating women only), and Mothers' Clubs (MCs), which are small peer group education sessions. The MCs were the mainstay of the BCC strategy used by both program approaches. Although the content of the MC sessions was largely similar in both program approaches, the approaches differed in the sequencing of the sessions and the number of sessions offered. The sequencing of MC sessions was age-based in the preventive approach and mothers attended a maximum of 18 monthly sessions, while in the recuperative approach, the session topics were chosen to be relevant for mothers of malnourished children, and 9 monthly sessions were offered (as per original design of WV-Haiti's recuperative approach). A total of 12 MC sessions were offered for pregnant and lactating women, 6 in pregnancy and 6 during the first six months of lactation. Health staff who facilitated the MCs were trained in the technical content of the MC sessions as well as in adult education-based communication and facilitation skills.

Evaluation Design

The evaluation used a community-level, cluster-randomized pre-post design, whereby 10 paired clusters of communities were randomly assigned to either the preventive or the recuperative program group. Cross-sectional surveys were conducted to assess the prevalence and severity of undernutrition among children 12-41 months of age at baseline and three years later, and statistical methods for analyzing group randomized designs were applied to evaluate impact.

The baseline survey was conducted between May and September 2002 and the post-evaluation survey was conducted exactly three years later, between May and September 2005, to minimize seasonal variations. All components of the intervention packages, except the newly developed BCC strategy, were implemented immediately following the baseline survey, i.e., in August-September 2002. The full BCC package, however, was implemented several months later (in May 2003) due to delays in material development, staff training, and field implementation.

In addition to data on the anthropometric outcomes, information was also gathered on a variety of intermediate outcomes relating to the use of the program services, household food security and assets, caregiver nutrition knowledge, physical and mental well-being, and infant feeding and care practices. The surveys were designed with the UNICEF conceptual framework in mind, so as to be comprehensive in their assessment of the resources available for care at the household and caregiver level.

The evaluation activities included a two-stage operations research process in addition to the surveys. The first stage, conducted a few months after the full intervention package was implemented in 2003, assessed the quality of program implementation and discussed the results with WV-Haiti to improve the quality of implementation. The second stage re-assessed the quality of implementation in 2004, and compared the quality of implementation between the two program approaches. No major implementation differences were seen, and implementation was of high quality, which suggested that both program approaches were operating largely as designed.

Results

At baseline, the two groups of communities (preventive and recuperative) were similar in all the main outcomes of the study as well as other major determinants of infant feeding and nutritional outcomes, such as food security, household socioeconomic status, maternal education, work patterns, social status, and physical and mental well-being.

Impact on child nutritional status

The key finding of the study is that in communities randomly allocated to receive a *preventive* approach of Title II-MCHN program, the prevalence of stunting, underweight, and wasting was 4, 6, and 4 percentage points lower after 3 years of operation compared to communities exposed to the *recuperative* program approach. The adjusted prevalence of stunting, underweight, and wasting among children 12-41 months in preventive areas was 33.9%, 14.8%, and 3.7%, respectively, whereas in recuperative communities, it was 38.2%, 20.8%, and 7.4%, respectively (using WHO reference standards (WHO 2006)). Mean height-for-age, weight-for-age, and weight-for-height Z-scores were also significantly higher in the preventive compared to the recuperative program communities. The differences (adjusted for child age and gender and clustering effect) in favor of the preventive group were + 0.14 for HAZ, + 0.24 for WAZ, and + 0.24 for WHZ. At baseline, there were no differences between program groups in any of the anthropometric indicators.

The magnitude of differences in favor of the preventive group for mean anthropometric indicators is comparable to other effectiveness trials aimed at reducing undernutrition through improved complementary feeding (Caulfield, Huffman, and Piwoz 1999), and to the average impact of USAID Title II MCHN programs documented by Swindale and collaborators (2004). Although the studies included in these reviews, which used before/after or post-intervention designs with a control group are not directly comparable to our study design (which compared two food-assisted MCHN program approaches), they are indicative of a range of effect that may be expected from this type of intervention. If we assume that our recuperative approach had some impact on reducing undernutrition (as suggested by the review of USAID food-assisted MCHN programs), then the larger impact of the preventive approach must be viewed as additional to that of the recuperative approach.

Plausibility of impact: The results of the main impact analysis are also supported by the fact that children who were exposed to the preventive program for the whole period of greatest

nutritional vulnerability (i.e., from 6-23 months of age) benefited more from the intervention than children who exposed only partially during this time period.

When compared to baseline, children's nutritional status appears to have deteriorated among the recuperative group, especially with regards to the prevalence of underweight and wasting, which increased from 18.0% to 20.8% (underweight) and from 3.4% to 7.4% (wasting). Stunting increased by 0.2 percentage point in the recuperative group, while it decreased by 3.5 percentage points in preventive areas. Underweight also declined by 2.8 percentage points in the preventive areas, while it increased by the same amount in the recuperative group. Wasting increased by 4 percentage points in the recuperative group, and decreased marginally in the preventive group. These results suggest that the preventive approach may have helped mitigate the deleterious effects on childhood malnutrition of the economic and political crisis that occurred in Haiti during the study period.

Relative costs and cost-effectiveness

The *total* direct costs for the two program approaches are the same, but the total variable/outside costs of the preventive approach, as designed and implemented in the evaluation area, are much higher than the costs of the recuperative approach. When examining costs *per beneficiary month*, rather than total costs, we find that the direct program costs per beneficiary-month are higher in the recuperative than in the preventive approach (\$21 USD versus \$15 USD). The outside program costs (which include the costs of the food commodities and health-care supplies) are the same in both groups – \$12 per beneficiary month.

The differences in costs (both total and per beneficiary month) is due to the much larger number of beneficiary-months in the preventive compared to the recuperative approach, owing to the design features of the preventive approach, as well as the differential participation rates for the two program approaches. Specifically, (1) *There is a larger number of eligible children in preventive compared to recuperative areas* because the prevalence of underweight children among the under-5s is less than the proportion of children 6-24 months among children less than 59 months); (2) *The duration of eligibility is longer, by design, in the preventive compared to the recuperative approach*: children in the recuperative approach were enrolled in the program for 9 months – as determined in WV-Haiti's development assistance program plan, while children in the preventive approach would be eligible to remain in the program for the entire period between 6 and 24 months, i.e., for up to 18 months. Thus the duration of intervention, by design, was double in the preventive approach; (3) *The rates of program uptake are higher in the preventive compared to the recuperative areas*: our analysis of program uptake (see below) suggests that the preventive approach seemed to elicit higher participation rates among eligible children than the recuperative approach, thereby further increasing the number of beneficiary-months in the preventive approach.

In sum, the larger cost of the preventive compared to the recuperative approach is due to the larger number of beneficiary-months, which, in turn, is due to a combination of factors including design, undernutrition prevalence, and program uptake patterns. The relative cost and relative cost-effectiveness of the preventive approach over the recuperative approach can, therefore, differ from context to context.

Pathways of impact

The impact of the program is dependent on good service delivery, as well as good participation in the program services by eligible beneficiaries, and finally, by appropriate use of the program inputs (food and BCC) by program beneficiaries that then lead to the intended intermediate impacts on food security, maternal knowledge, and childcare and feeding practices.

As noted above, there were no implementation differences between the program approaches, and the program was implemented in all evaluation communities. Thus, differences between program approaches on child nutrition outcomes may have been achieved through differences in program participation and take-up. Other potential explanations for differences between program approaches include differences in impacts on the care environment at the household and caregiver levels, and/or differences in biological vulnerability (under two years vs. older children). In addition, the substantially higher proportion of children exposed to the fortified foods in the preventive group also likely contributed to the differences in outcome.

We briefly review the different steps in the pathways of impact below.

Program participation and uptake

The enrollment rates in the food assistance and BCC package among pregnant and lactating women were similar in both program communities. Enrollment among children was different and was true to the design (as intended). Overall, more children in the preventive communities had ever been enrolled in the program (73%) than in the recuperative communities (28%).

Around 75% of eligible children in the preventive program (6-23 months of age) were receiving food assistance at the time of the final survey. In the recuperative group, only 29% of children 6-59 months who were underweight at the time of the final survey were enrolled, most likely because a smaller proportion of eligible children were brought to the Rally Posts - the main entry point into the program. Targeting was excellent in preventive communities, with 93% of enrolled children meeting the age eligibility criteria; in the recuperative group, around 57% of currently enrolled children were underweight. Thus, program uptake in the two approaches was quite different, most likely because of differences in population understanding of the targeting mechanisms in the two approaches.

Impact on household care context and resources

At final survey, households in preventive communities had statistically significantly lower food insecurity than households in recuperative areas, but differences were small. More meaningful differences were observed between current participants and nonparticipants in both program groups, suggesting a positive short-term impact of food assistance on household food security. Compared to baseline, overall food insecurity did not improve over the three-year duration of the study, and continues to be severe in the program area. There was thus no evidence of a long-term effect of the program on food security; the same was true for asset ownership, which had not improved since baseline, and was not different between program communities or between participants and nonparticipants. Thus, overall, the program seems to

have had a positive impact on food security in the short term, suggesting that the slightly greater impact on households in preventive compared to recuperative communities is due to their longer eligibility to receive food assistance. There was, however, no evidence that the program had a longer-term impact on food insecurity (i.e., beyond the period of receipt of food aid), which is most likely due to the severe economic constraints faced by poor household in Haiti at the time of the study. It is possible that more sustained impacts of such a program could be obtained in less constrained times.

Impact on maternal context and resources

We did not find any differences between program groups on maternal resources such as education, social support, and autonomy in decisionmaking either at baseline or at the final survey. Impacts on other resources are discussed below.

Maternal knowledge: Maternal knowledge of several topics related to infant and young child feeding and general health and nutrition topics was higher among the preventive compared to the recuperative group at final survey, but differences were generally small. Larger differences were found between mothers who had ever participated in the program, compared to those who had never participated. Maternal knowledge also significantly improved from baseline to final survey in both program groups, suggesting that overall, the BCC strategy was successful in improving overall maternal knowledge of health and nutrition in the program communities. The longer duration of exposure to the BCC in the preventive compared to the recuperative group and the more age-specific and timely delivery of messages, however, resulted in only small differences in knowledge between groups.

Physical and mental well-being: At baseline, we found that food insecurity was strongly associated with all measures of women's well-being, particularly mental well-being. At final survey, respondents in the preventive communities were better off than those in the recuperative communities on four of the women's well-being measures – self-rated health, mental stress, and two measures of life satisfaction. *Current participants* in preventive communities also had better self-rated health, lower mental stress, and lower time stress than nonparticipants. Thus, it appears that the program's short-term impact on food security had positive benefits on women's mental well-being; and given that women in preventive communities received food assistance for longer periods of time (for up to 30 months if they were enrolled in the program during pregnancy and the first 6 months of lactation), they benefited more in terms of improved mental well-being than women in recuperative communities. Again, there was no evidence of long-term benefits of the program on these aspects, as reflected by the lack of difference in well-being between mothers who had *ever participated* and those who had not.

Impact on child feeding and care practices

Awareness, trial, and adoption of recommended practices

The final survey included an assessment of the awareness, trial, and adoption of seven key practices recommended by the program. For most key practices, respondents in preventive program areas were more likely to report awareness, trial, and adoption than were respondents in recuperative areas. In most cases, however, differences between program areas were of

relatively small magnitude, while differences between those *ever* exposed to the program and those *never* exposed were large. Thus participation in either program approach had a beneficial effect on awareness, trial, and adoption of recommended child feeding and care practices, with mothers in preventive communities doing slightly better on these aspects than women from recuperative communities, probably as a result of their longer exposure to the BCC intervention and the more timely delivery of the messages (i.e., at the age when knowledge and adoption of specific practices was most relevant). However, the reported reasons for non-trial and adoption of practices suggests that overall, practices that required few material resources to try and adopt were more likely to have improved with exposure to the program.

Infant and young child feeding practices

Breast feeding: There were no differences between program groups in early feeding practices or in the timing of introduction of liquids and complementary foods. This was expected since the two program approaches offered exactly the same services until the child reached 6 months of age. Large improvements in these practices were seen since baseline, however, and equally large differences were observed between participants and nonparticipants for both program approaches. Breastfeeding duration was the same in both program groups at baseline and at final survey and ranged from 18 to 24 months of age.

Complementary feeding practices, including consumption of the donated commodities: There were few differences between program groups in complementary feeding practices at final survey, with the exception of diet quality and the consumption of animal source foods, which were slightly higher in the preventive group. The infant and young child feeding indicator – which combines information on breastfeeding, number of meals/day and dietary diversity – also showed a statistically significant, but modest difference in favor of the preventive compared to the recuperative communities. As with maternal knowledge, differences in many of the feeding practices were larger between ever participants and never-participants than between program groups. These include meal frequency, use of baby bottles, vitamin A supplement consumption, and consumption of fortified donated commodities (WSB) (the latter being expected). Several practices had markedly improved since baseline; these included a reduction in the use of baby bottles (which was halved since baseline), increases in vitamin A supplementation, and in appropriate feeding during and after diarrhea. The overall consumption of animal source foods, however, was lower at final survey than at baseline, probably a reflection of the economic crisis and related price increases in Haiti over the study period.

The proportion of children consuming recipes made with WSB was not different between current participants in the program groups. However, since there is a larger *number* of program beneficiaries receiving WSB and other fortified foods in the preventive group, particularly among the younger children, it can be assumed that the fortified commodities contribute more to nutrient intake among children in the preventive approach than in the recuperative approach.

Preventive and curative care-seeking and hygiene

Immunization rates were not different between groups at baseline or final surveys, but they had improved from a very low 11% of children fully immunized at baseline to approximately 30% at final survey. Patterns of attendance at Rally Posts were similar between

groups for children during the first year, which explains the lack of difference in immunization rates between groups. The low immunization coverage seemed to be largely due to poor supply as opposed to low demand, as suggested by our operations research results. Care-seeking for fever, cough, fast breathing, and diarrhea were not different between program groups at baseline or at the final survey and were not different between participants and nonparticipants. However, care-seeking rates were lower at the final survey when compared to the baseline survey, possibly due to decreased severity of illness and/or better home management of illness. Use of oral rehydration (ORS) and sugar-salt solutions (SSS) were higher since baseline, suggesting improved home care for diarrhea as a result of the program. There were no differences in markers of hygiene practices between program groups either at baseline or final surveys, but there was a slight decrease in hygiene scores since baseline. There were also no meaningful differences in markers of hygiene practices between participants and nonparticipants.

Conclusions on the pathways of impact

Our findings suggest that the pathways of impact, which led to better child nutritional outcomes among preventive communities, operated mainly through the changes in the childcare context resulting from participation in the program and through greater availability of the fortified commodities to children in the preventive approach. More specifically, participants in preventive communities had greater food security while in the program, which positively affected the caregivers' mental well-being; this, combined with better knowledge, awareness, trial, and adoption of several recommended feeding and care practices, likely resulted in a generally more supportive care environment, which, in turn, could have had a greater impact of the preventive compared to the recuperative approach on child nutrition outcomes. Furthermore, a larger number of children in the preventive approach received fortified commodities in preventive approach than in the recuperative approach. Since consumption of the fortified commodities is high among those who receive them, this is also a potential pathway of impact.

It is important to note that in general, differences between the preventive and recuperative approaches in the different aspects of the childcare context were relatively small. This was somewhat surprising, given the much longer duration of exposure to program inputs among participants in the preventive approach, the higher participation rates, and the explicit effort to deliver the BCC intervention in the most timely fashion in the preventive compared to the recuperative approach. The results thus suggest that it may be the cumulative effect of relatively small differences in the multiple aspects that comprise the care environment, which are responsible for the larger benefits of the preventive compared to the recuperative approach in improving child nutritional status in our study.

Our results also showed that the two program approaches were operating equally well and that none of the program implementation and staff-related factors differed between the two approaches. This allows us to conclude with certainty that the greater nutritional impact observed in preventive communities was truly due to a more effective program approach.

Implications for Programs and Policies

This section briefly summarizes the overall lessons learned from this evaluation and their implications for programs and policies.

A preventive approach to addressing childhood undernutrition is more effective than a curative approach. The direct implications of our results are that in order to improve effectiveness, food-assisted MCHN programs should target all children under the age of two years, as opposed to malnourished children under-five, and continue to target pregnant and lactating women. Severely malnourished children up to 5 years of age should continue to be screened and receive appropriate care.

Focusing on the under-twos is both feasible and successful in a programmatic context. There is renewed global attention around this critical age group, but few examples exist of feasible, successful, and effective programs. This evaluation provides an example of the feasibility and effectiveness of this approach, in a programmatic context, as well as an example of *how* such programs can be developed, strengthened, and monitored under real programmatic conditions. Our work thus provides programmatic evidence for current policy to focus nutrition interventions on the under-twos.

A well-designed and well-implemented behavior change strategy can improve infant feeding practices regardless of whether a preventive or recuperative approach is used for targeting. This evaluation provides an example of an approach and a specific set of tools that were used for developing and implementing a locally relevant, programmatically feasible BCC strategy for improving child feeding and care practices among children under two years of age.

Variable cost per beneficiary-month is approximately the same for the two programs but total costs are higher for preventive approach as designed in this study. The implications of these findings are that programs should carefully review their program design, geographic priority areas, and targeting mechanisms based on their resources and target number of beneficiary-months. Programs can also attempt to balance or reduce the costs of the preventive approach by changing the age range for targeting, the duration for which children are enrolled in the program, and/or even the amount of the food assistance provided. The impact of making any of these changes to the preventive approach, however, should be rigorously tested to ensure that these modifications do not result in losses in effectiveness and nutritional impact of the program.

Investing in formative and operations research is important for program success. This evaluation provides strong evidence that investing in formative research can help design effective BCC programs that are grounded in the sociocultural context, locally relevant, and programmatically appropriate. The study also provides evidence that operations research provides critical insights regarding the quality of implementation and service delivery for evaluators and program implementers.

Recommendations for Future Research

Evaluation of the preventive approach in other settings, preferably using a control group. Future evaluations of the preventive approach should include a control group, so that the magnitude of the absolute effect of the preventive approach could be assessed. Evaluations in other contexts should also pay adequate attention to program theory so as to generate knowledge and consensus on developing and implementing programmatic approaches for delivering food-assisted MCHN services to the critical under-two age group.

Separating contributions of food assistance and BCC components. Since the food component contributes the majority of the cost of the program approach, it would be useful to conduct evaluations that allow separating the contributions and cost of food assistance from the BCC component. At the same time, the role of limited food and economic resources in dampening the potential impact of BCC programs in impoverished contexts should be kept in mind. Research should also assess the long-term impact of both program approaches and determine whether short-term benefits in nutritional status are maintained over time.

Testing alternative designs for a preventive approach. Future evaluations should test different delivery systems for the preventive approach and compare the cost of alternative designs such as modifying the age of eligibility, reducing the duration of eligibility or the size of the food assistance package. These evaluations should include a control group so that the cost-effectiveness of the different approaches could be assessed.

Lessons for Program Evaluations

Using program theory to develop evaluations. The approach to this evaluation was based on program theory that considered the full pathway of expected impacts. Developing the evaluation activities based on this framework was useful not only for the measurement of the different program inputs and outcomes but also to help identify bottlenecks that could influence the program's impact.

We paid careful attention to the design of the preventive approach in particular because it was important to ensure that the approach was truly preventive in nature. This attention to design is important to consider in all evaluations, because evaluating a package that is not designed to be true to the concept behind it is inefficient and does not allow appropriate interpretation of the results of an evaluation.

Documenting implementation quality and program utilization. Attention to the quality of implementation of the intervention package as well as utilization of the intervention by intended users is critical in any impact evaluation. As with the design, a poorly implemented and poorly utilized intervention is not likely to yield the expected impact. Thus, attention to design and to implementation quality, as well as patterns of utilization, are both essential to draw conclusions about the effectiveness of interventions.

Age of intervention versus age of expected impact. Finally, we suggest that it is extremely important that evaluations focusing on child growth outcomes pay attention to the age at which an intervention to improve growth is delivered versus the age at which impact is expected. Outcome assessments should be conducted within the age group that is most likely to show impact, which is not necessarily the same as the age of targeting of interventions.

Conclusions

This evaluation shows that a preventive approach to a Title-II MCHN program is more effective than the traditional, recuperative approach at reducing childhood undernutrition among children aged 12-41 months. The variable costs of both the program approaches was approximately the same on a beneficiary-per-month basis but the preventive approach was more

expensive in the Haiti setting because of differences in design and participation rates between the two approaches. Moreover, the relatively low levels of undernutrition in this population (compared to other equally poor countries) results in a larger number of children under two years of age, compared to underweight children, and thus more beneficiary-months in the preventive approach. Both programs also had a significant impact on improving maternal knowledge and feeding practices compared to baseline, showing that a well-designed and well-implemented BCC strategy integrated within a Title-II MCHN program can be highly effective. These benefits were obtained with two carefully designed and implemented program approaches operating under particularly difficult field conditions in rural Haiti.

We believe the findings of this study are generalizable, given the remarkably similar patterns of child growth globally, and the similar prevalence of undernutrition in Haiti and other countries in the world. We conclude by suggesting that there are four conditions for the results of this study to be replicable: (1) good program design based on sound formative research; (2) effective implementation and service delivery monitored with regular operations research; (3) good incentive structure and high staff motivation monitored and fostered by effective supervision; and (4) similar or higher levels of undernutrition than in Haiti (e.g., 25% - 30%). Although a similar preventive program could be effective in a population with lower levels of undernutrition than in Haiti, the cost per case of undernutrition prevented would be higher. However, most poor countries in Africa and Asia, and even some countries of Latin America, currently have a higher prevalence of undernutrition than Haiti. For these countries, a preventive approach is strongly justified for MCHN programs.

1. INTRODUCTION AND SCIENTIFIC RATIONALE FOR THE STUDY

This report presents the results of a three-year project evaluating a food-assisted (Title II) Maternal and Child Health and Nutrition (MCHN) program implemented by WV-Haiti in the Central Plateau of Haiti. The evaluation compared two different approaches of targeting food assistance and a behavior change and communication strategy to improve child-feeding and care practices: (1) the *recuperative*¹ approach, which targets children with mild or moderate undernutrition (weight-for-age < -2 Z-scores); and (2) the *preventive* approach, which targets all children 6-24 months of age, in an effort to prevent, rather than cure, childhood undernutrition.

The project was motivated by a critical programmatic question for Title II program implementers around the world, i.e., whether blanket targeting of Title-II MCHN programs to children less than two years of age is more effective than targeting undernourished children. Scientific evidence from controlled intervention trials suggests that children under two years of age are more responsive to nutrition interventions than older children (Schroeder et al. 1995). Research also shows that the earlier and the longer food supplementation is provided before the child reaches 2 years of age, the greater the benefits on growth and on a series of long-term physical, cognitive, and reproductive outcomes (Rivera and Habicht 1996, 2002; Martorell 1995). Based on this, and other, evidence, the USAID Food for Peace office has recommended applying a preventive approach to human capital formation in Title II MCHN programs, including investing in early childhood nutrition.² A recent World Bank report also urges governments to invest in preventing undernutrition among children under two years of age as a means of promoting national development and human capital formation (World Bank 2006). The focus is on the prevention of undernutrition, rather than cure; and on achieving cost-effectiveness by allocating resources to children who have the greatest potential for response.

Program implementers are thus increasingly encouraged to reallocate their resources to focus on prevention among children under two, rather than on the recuperation of under-five children with mild to moderate undernutrition. Without programmatic evidence showing the feasibility and effectiveness of the preventive approach, however, program implementers have been hesitant to move in that direction.

To address this knowledge gap, researchers from the International Food Policy Research Institute and Cornell University undertook a study commissioned by the Food and Nutrition Technical Assistance (FANTA) project at the Academy for Educational Development (AED) to compare the cost-effectiveness of the preventive and recuperative approaches, as implemented by WV-Haiti in the context of their Title II MCHN program in the Central Plateau. This report summarizes the research and program development activities that were undertaken by the team to design, implement, and evaluate the impact and cost-effectiveness of the two program approaches on alleviating childhood undernutrition in rural Haiti.

¹ The term “recuperative” is used in this report to refer to mild and moderate undernutrition, i.e., weight-for-age or height-for-age less than -1 (mild) or -2 (moderate) Z-scores. The report does not directly address severe malnutrition defined as weight-for-age or height-for-age < -3 Z-scores and the community therapeutic care approach used for the treatment of severe acute malnutrition.

² From the Food for Peace Expanded Conceptual Framework for Food Aid and Food Security, 2002.

The remainder of this chapter presents some background information on Haiti and on Title II MCHN programs, and describes the scientific rationale that motivated the study. It concludes by describing the organization of the report.

1.1 Haiti: The Poorest Country in the Western Hemisphere

Haiti is the poorest country of the Latin American and Caribbean (LAC) region and is ranked 153rd out of 177 countries by the Human Development Index (UNDP 2005). Three-quarters of the Haitian population are poor (living on less than \$2 USD per day), and over half are extremely poor (living on less than \$1 USD) (Sletten and Egset 2004). Extreme poverty is largely a rural problem, with 77% of the extreme poor living in rural areas. The country's economy has been deteriorating steadily since the 1980s and real per capita GDP has since been falling at an average of 2.5% per year (<http://devdata.worldbank.org>). Not only is Haiti the poorest country of the region, it also has the most unequal income distribution, with a Ginni coefficient of 0.65, next to Brazil at 0.59 (World Bank 2003).

Not surprisingly, this impoverished country has the worst health indicators of the region, with an under-five mortality rate of 118 per 1,000, a prevalence of chronic undernutrition of 23% (reaching 30% in the poorest regions), an excessively high maternal mortality rate (680 per 100,000 live births), and the highest prevalence of HIV/AIDS outside Sub-Saharan Africa (2.5%) (<http://web.worldbank.org>). Life expectancy is a mere 51 years. Access to services is a key constraint: only half of the population has access to clean drinking water and 28% to sanitary facilities; childhood measles immunization coverage is only around 50%, and less than one-quarter of all births are attended by skilled health staff (<http://devdata.worldbank.org>). Public spending on health was 3% of GDP in 2002, compared to about 6.5% of GDP in Mexico (in 2005).³ Education indicators are equally alarming: half of the population is illiterate (UNESCO 2002) and primary school attendance rates were 52% for men and 56% for women in 2000 for the country as a whole, and as low as 34% in some of the poorest rural areas (EMMUS-III 2001).

Haiti's current economic, social, and development profile is the result of years of internal conflicts and climatic shocks that have devastated its economy, depleted its natural and human capital, weakened its institutions, and inflicted severe hardship on its population. The years 2002-2006, when the study took place, were no exception. In fact, the period between 2003 and 2005 brought greater hardships than the preceding years, with political unrest in 2003 leading up to the exile of the president, Jean-Bertrand Aristide, in early 2004, and continued political strife and violence following the exile. Although much of the violence was restricted to the capital city, the political climate had serious impacts on the economy of Haiti and on food and fuel prices in the country (International Crisis Group 2005).

1.2 Title II-MCHN Programs

USAID spends around \$90 million/year on Title II-funded MCHN programs (Swindale et al. 2004). These programs either use donated food commodities as direct food assistance to

³ From the OECD Health Data 2007 (accessed July 27, 2007)
http://www.oecd.org/document/16/0,3343,en_2649_34631_2085200_1_1_1_1,00.html

program beneficiaries and/or monetize the donated commodities to support the MCHN services offered through the programs. Most Title II MCHN programs are intended to be community-based, and to offer or support a variety of interventions with proven impacts on child nutrition and child survival, i.e., immunizations, breastfeeding promotion, complementary feeding promotion, vitamin A capsule distribution, etc. (Swindale et al. 2004).

A recent review of Title II-funded MCHN programs (Swindale et al. 2004) suggests that overall, MCHN programs implemented using the Title II mechanism have yielded reductions in stunting that average 2.4 percentage points per year and reductions in underweight that average 1.9 percentage points per year. Although few of the studies reviewed used rigorous evaluation designs, overall the results are indicative of positive, albeit relatively small, benefits on childhood undernutrition.

In Haiti, USAID provides Title II resources to four major private voluntary organizations (PVO): CARE, Catholic Relief Services, Save the Children USA, and WV-Haiti (referred to as Cooperating Sponsors (CS)). All four CS in Haiti use the Title II resources to provide MCHN services directly or through collaboration with the government services, and Title II support is usually provided to program beneficiaries as direct food assistance. All four CS use a recuperative approach that provides food assistance to malnourished children under the age of five years.

In its 2002-2006 Development Activity Program (DAP) proposal to receive Title II resources, WV-Haiti noted their intention to pilot test the targeting of food assistance to children under two in one of their priority areas (WV-Haiti 2001). This was a unique opportunity to test the approach under programmatic conditions and compare it with the recuperative approach. Thus FANTA/AED, with support from USAID, commissioned an evaluation, which is described in this report.

1.3 Rationale for a Preventive Approach to Reducing Undernutrition

This section presents the scientific rationale that motivated this study; it describes what was known at the onset of the study regarding the potential usefulness of a preventive approach to tackle childhood undernutrition. The focus is on food supplementation studies - as opposed to nutritional interventions to improve child-feeding and care practices - because this is where most of the scientific evidence from controlled trials is available. The section also presents the behavioral rationale for the specific set of child-feeding and care practices that the study focused on in its behavior change and communication strategy (see Chapter 4 for a description of the strategy).

1.3.1 The recuperative versus the preventive approach

Food distribution programs have often failed to improve the nutritional status of their beneficiaries (Beaton and Ghassemi 1982) and their lack of impact has been attributed to the fact that they often failed to target children who were most likely to benefit. The traditional recuperative approach has the major drawback of intervening too late, i.e., once children's growth has fallen below a certain cutoff point. This often happens after months of growth

faltering and at ages when children have limited potential to respond to nutrition interventions, particularly in such a way as to improve stunting.

The preventive approach, on the other hand, is designed specifically to address this concern by targeting children *before* their growth falters **and** by intervening during their period of highest growth velocity, and of maximum potential to benefit from nutrition interventions (0-24 months of age) (Schroeder et al. 1995; Lutter et al. 1990; Allen 1994; Rivera and Habicht 1996, 2002). The preventive approach thus aims at preventing children from getting malnourished and thus needing to be recuperated. By maintaining children's growth at a higher level throughout their most vulnerable period (0-24 months of age), it is expected that not only short-term, but also long-term, benefits on growth will be achieved (Martorell 1995).

1.3.2 Biological rationale for targeting food supplementation to children under 24 months

The rationale for targeting food supplements (and other nutritional interventions) to children under 24 months of age is based primarily on current knowledge regarding the patterns of growth of young children and their response to food supplementation interventions. A brief overview of this scientific evidence is presented below.

1.3.2.1 Patterns of child growth in Haiti and other developing countries


Recent national-level data from Haiti show that approximately one-third of children less than three years of age in Haiti have low height-for-age (< -2 Z-scores) (EMMUS-III 2001). Growth curves using data from the Demographic and Health Surveys data from 2000/2001 for Haiti (see Figure 1.1) show the typical pattern found in most developing countries around the world where mean height-for-age (and weight-for-age) decline almost linearly from birth to approximately 18 months, after which they tend to stabilize at a low level (Ruel 2001; Shrimpton et al. 2001).

The data on patterns of child growth in Haiti (and in most of the developing world) show that children under two are clearly the most vulnerable to growth faltering and this period is, therefore, a time when interventions to prevent the rapid decline in nutritional status are most needed. Note that the similarity in growth patterns between children from Haiti and from other contexts provides reassurance that the information currently available on optimal timing and duration of supplementation (presented in the following sections) applies to all developing country contexts.

1.3.2.2 Patterns of response to food supplementation

Evidence from randomized controlled trials suggests that the impact of supplementary feeding interventions on child growth is determined by two key factors: (1) the timing of the intervention (child age at the time of the supplementation), and (2) the duration of supplementation. A brief overview of relevant findings for the present evaluation is presented below.

Figure 1.1 Mean weight-for-height (WHZ), weight-for-age (WAZ), and height-for-age (HAZ) of rural children in Haiti (EMMUS-III 2001)



Effect of timing of supplementation interventions on overall growth impact

Schroeder et al. (1995) showed that in rural Guatemala, the greatest impact of food supplementation was achieved among children in their first and second years of life, and that no impact was found from three to seven years of age. In a different study setting, urban Colombia, Lutter et al. (1990) demonstrated that within the first 24 months, the greatest response to supplementation was seen in infants between 9 and 12 months of age, the peak period of diarrheal morbidity in this population. Finally, observational research from the Nutrition Collaborative CRSP in Mexico, Kenya, and Egypt showed that improved diets after the age of 18 months were not associated with better nutritional status (Allen 1994). Based on some of this research, Beaton provides convincing arguments in his review of targeting for supplementary feeding more than 10 years ago, that the “effective age range for substantive impacts on linear growth seems to be six months to two or perhaps three years” (Beaton 1993, p. 47). He clarifies that the implications are not that no benefit at all can be expected beyond this age, but that the magnitude of benefits may not justify the additional costs of providing food supplements to children after three years of age.

Thus, evidence suggests that maximum benefits from improving dietary intake, including through programs that provide food supplements, will be most effective in preventing malnutrition in the period of approximately 6-24 months of age. This is not surprising because this is the period of maximum expected growth velocity and also the period of greatest risks of growth faltering due to inappropriate complementary feeding practices and increased risks of infectious diseases rates, especially diarrhea.

Effect of timing of supplementation interventions on faltering and recovery rates

While it is important to examine the overall impact of supplementation at different child ages, it is also useful to understand through longitudinal analysis how supplementation affects the rates of growth faltering and the rates of recovery from faltering. Only two studies, both using the Guatemala longitudinal supplementation study conducted in the 1970s in four rural communities, have examined the differential impact of supplementation on faltering and recovery rates in weight-for-height (WHZ) (Rivera and Habicht 1996, 2002).

The analyses confirm that the impact of supplementation on the prevention of faltering (maintaining a weight/length category during a specific supplementation period) is age-dependent. The authors found a much larger impact on the *prevention* of faltering in WHZ among children who were 6 to 24 months old at the time of the intervention (Rivera and Habicht 2002). In this age group, the faltering rate among those receiving the food supplementation intervention was 0.19 in contrast to 0.45 among nonsupplemented children, a difference of 0.26, which was due to the supplementation. The much smaller difference of 0.08 for the same comparison among children between 24 and 48 months of age was not significant.

Recovery from faltering was also found to be age dependent. Among 6-24-month-old children who had received the supplementation for 12 months, the rates of recovery from faltering was 0.78 for supplemented children and only 0.41 for those without the supplement, a difference of 0.37, which was due to the supplementation. Again there was no effect among the 24-48 month old children.

Effect of the duration of supplementation

There is limited research on the optimal duration of food supplementation needed for maximal impact. The only information available that we are aware of comes from analyses of the Guatemala longitudinal trial (Rivera and Habicht 1996). In this context, although 59% of infants had recovered from faltering in WHZ within 3 months of supplementation, greater impacts were achieved with 12 months of supplementation, reaching almost 80% of children. These data suggest that longer durations of supplementation (6-12 months) are likely to have more impact than shorter durations (3 months).

Overall, scientific evidence regarding patterns of growth faltering in early childhood and the impact of food supplementation on child growth in countries like Haiti, suggests that:

- ❖ The age of “active” growth faltering is between 6 and 18-24 months of age;

- ❖ The greatest growth response is achieved when food supplementation is provided during the first 24 months of age and for a duration of at least 6-12 months.

These insights suggest that a preventive approach of food supplementation targeted to all children between 6 and 24 months is likely to contribute to reducing childhood undernutrition in poor communities. Food supplements without appropriate care and feeding practices, however, are unlikely to make a significant dent in undernutrition prevalence. Thus, food supplementation interventions must be combined with a strong behavior change and communication strategy to help caregivers use the donated food commodities optimally and adopt recommended child-feeding and care practices. The behavioral rationale for combining food supplementation and BCC interventions is presented below.

1.3.3 Behavioral rationale for combining food supplementation and behavior change communication strategies

In children, positive changes in health and nutrition manifest themselves in many forms, including improvements in growth and development and reductions in morbidity and mortality. Each of these outcomes is the result of complex interactions between familial caregiving practices and the biological underpinnings of health and nutrition. For example, in order to protect a child from a vaccine-preventable disease, such as measles, the family must know when and where to take the child for the vaccination, must have the resources to carry out these actions, and the vaccine itself must be safe and effective. The availability of the vaccine is powerless to prevent the disease in the child without the parental actions.

Similarly, to enable children to grow normally, there are many parental caregiving practices related to food that are essential to ensuring adequate nutritional intake. These practices include obtaining and selecting foods that meet nutritional requirements, preparing them safely and in a form that is appropriate for the child's age, and feeding them in a manner that encourages adequate intake. In order to engage in these critical caregiving practices, parents need access to the foods their children require; they need access to fuel, water, and other materials to prepare and preserve these foods, and time and physical energy to carry out the activities. They also need knowledge. These are essential underpinnings of nutrition and health-giving practices, which in turn are the prerequisites for child health and well-being. Because caregiving practices are the links between resources and knowledge, on one hand, and child health on the other, programs that seek to improve child health and nutrition must, by definition, change caregiving practices.

Thus, it is important that both the preventive and recuperative approaches implemented by WV-Haiti ensure that in addition to providing *food* to children under the age of 24 months, adequate information is provided to caregivers to ensure that these foods are *fed* appropriately to these young children. Along with this, it is also crucially important to ensure that other aspects of feeding and care that are important for the preventive age group are also addressed. The key aspects of care and feeding to address in the vulnerable period of 0-24 months of age are breastfeeding, complementary feeding, and other preventive and curative health-related practices like good hygiene, timely immunization, appropriate home health care, and care-seeking during illness. This section presents a summary of current recommendations for the feeding of infants and young children under the age of 24 months and also briefly summarizes current evidence

regarding the impact of BCC programs on influencing these practices. The technical basis for the feeding recommendations is not described here. However, they are described in detail in an article in the *Food and Nutrition Bulletin* (Dewey and Brown 2003).

1.3.3.1 Current infant and young child-feeding recommendations

Current infant and young child-feeding recommendations are derived from “Guiding Principles for Complementary Feeding of the Breastfed Child” (PAHO/WHO 2003), and provide guidelines for appropriate feeding of breastfed infants from 0-23 months of age in developing countries. The following specific dimensions of infant feeding are covered in these Guidelines:

- *Duration of exclusive breastfeeding and age of introduction of complementary foods:* Practice exclusive breastfeeding from birth to 6 months of age, and introduce complementary foods at 6 months of age while continuing to breastfeed.
- *Maintenance of breastfeeding:* Continue frequent, on-demand breastfeeding until 2 years of age or beyond.
- *Responsive feeding:* Practice responsive feeding, applying the principles of psychosocial care.
- *Safe preparation and storage of complementary foods:* Practice good hygiene and proper food handling.
- *Amount of complementary foods needed:* Start at 6 months with small amounts of food and increase the quantity as the child gets older, while maintaining frequent breastfeeding.
- *Food consistency:* Gradually increase food consistency and variety, as the infant gets older, adapting to the infant’s requirements and abilities.
- *Meal frequency and energy density:* For the average healthy breastfed infant, meals of complementary foods should be provided 2-3 times per day at 6-8 months of age and 3-4 times per day at 9-11 and 12-23 months of age.
- *Nutrient content of complementary foods:* Feed a variety of foods to ensure that nutrient needs are met. Meat, poultry, fish, or eggs should be eaten daily, or as often as possible. Vitamin A-rich fruits and vegetables should be eaten daily.
- *Use of vitamin-mineral supplements or fortified products for infant and mother:* Use fortified complementary foods or vitamin-mineral supplements for the infant, as needed.
- *Feeding during and after illness:* Increase fluid intake during illness, including more frequent breastfeeding, and encourage the child to eat soft, varied, appetizing, favorite foods.

Although Guiding Principles do currently exist for non-breastfed children (Dewey, Cohen, and Rollins 2004), they were not available at the time our BCC strategy was designed, and were therefore not considered. Other special situations that were not addressed in our BCC strategy include the feeding of children recuperating from severe malnutrition and the feeding of infants born to HIV-positive mothers.

1.3.3.2 Evidence of the impact of behavior change communication interventions on feeding practices and growth outcomes

A review of complementary feeding interventions provides evidence that BCC programs can be effective in reducing child undernutrition in a variety of contexts (Caulfield, Huffman, and Piwoz 1999). Studies published after the review by Caulfield, Huffman, and Piwoz (1999) also continue to provide evidence that targeting of behavior change communication (BCC) to caregivers of children in the 6-24 month age range can have beneficial impacts on child growth. Studies from settings such as India (Bhandari et al. 2004, 2005) as well as Peru (Penny et al. 2005) show that different approaches of targeting BCC programs to mothers of children 0-24 months of age can yield benefits in terms of improved breastfeeding practices as well as improved complementary feeding practices, and thus better growth outcomes.

A few studies also point to the impact of targeting mothers of undernourished children with nutrition education. In particular, the success of the Hearth approach and the Positive Deviance approach in settings such as Haiti and Vietnam are notable (Pachon et al. 2002; Mackintosh, Marsh, and Schroeder 2002; Schroeder et al. 2002). A recent study from Bangladesh also points to the added benefit of combining nutrition education with provision of supplementary food (Roy et al. 2006) rather than either intervention alone on improving growth outcomes for undernourished children. At the same time, although this evidence suggests that undernourished children can recuperate through intensive BCC interventions, there is no evidence to suggest that targeting BCC interventions to mothers of undernourished children has impacts on overall community-wide undernutrition. Furthermore, as with studies on supplementation of young children, there are no studies that compare the behavioral and biological impact of targeting behavior change communication to caregivers of children in the critical age group of 0-24 months to the impact of targeting caregivers of undernourished children.

1.3.3.2.1 Factors that influence the success of BCC programs

The review by Caulfield, Huffman, and Piwoz (1999) also provides a basis for understanding the various factors that influence the success of behavior change communications programs to achieve their goals. The authors found that the success of BCC programs depends not only on the design, targeting, and outreach of the program, but also on contextual factors, such as community involvement and political commitment. Specifically, the successful programs reviewed by Caulfield and collaborators used very similar approaches to program design. These approaches all included a number of stages of formative research, such as a review of existing materials related to infant feeding in the program areas, ethnographic research to understand current infant feeding practices and their motivations, an assessment of current complementary foods, and recipe trials to develop enriched complementary foods. Additionally the development of program strategies in all cases used a comprehensive approach, which took

into account contextual facilitating factors and the findings from the formative research. These programs showed substantial improvements in caregiver knowledge and recall of program messages, increased intakes of complementary foods, and, furthermore, improvements in child nutritional status (height-for-age and weight-for-age) that were similar to improvements seen from food supplementation studies. Even taking the problematic design of some of the program evaluations into account, the authors estimated that BCC programs could improve child nutritional status by as much as 0.1 to 0.4 Z-scores.

Some of the features of the programs that could have contributed to these successes are likely to be their attention to the local context within which the program was to operate, an in-depth understanding of infant feeding practices that was based on solid formative research, and a program strategy that used a comprehensive approach in its design and implementation. Almost all the programs reviewed, for instance, used very clear key messages that were age-appropriate and action-oriented and that would allow caregivers to make easy changes in infant feeding that were adapted to the child's stage of development. The programs also used multiple approaches to reach caregivers, usually combinations of mass media and individual advice and counseling. In addition, they used a variety of communications methods, such as radio spots, cooking demonstrations, story telling and drama, and all of these were accompanied by appropriate visual communications materials like posters, counseling cards, and take-home cards on infant feeding.

1.3.3.2.2 Implications for the design of the preventive and recuperative approaches

The implications of the evidence on the success of BCC programs for this evaluation are that the preventive approach of targeting food assistance could be expected to benefit from a BCC program that is also designed to be preventive in nature, and that addresses the special needs of children in the 6-24 month age group. This is particularly important because the changing nature and complexity of recommended infant and young child-feeding practices in the 6-24 month age range present a daunting challenge both to caregivers and to program implementers. To be effective, a BCC program must address the range of practices recommended at different ages, and must do so in ways that are culturally appropriate and timely. In addition, current evidence suggests that successful BCC programs are those that are grounded in formative research and locally relevant, and that use multiple channels to reach program clients. These principles have been applied in developing the preventive and recuperative approaches, and the process for doing so is described in Chapter 4 of this report.

1.3.4 Conclusions

This section presented the biological and behavioral rationale for designing nutrition interventions that have a preventive focus and that target children less than two years of age. In addition, it laid the basis for the discussion on infant and young child-feeding practices, which is further discussed in Chapter 4. This section also provided evidence that the growth patterns of Haitian children are similar to those of children in other developing countries worldwide. In doing so, it provides justification for the use of global evidence on child growth and nutritional interventions to design this particular evaluation in Haiti. It also provides some reassurance that the results of this evaluation can be used to inform decisions about the potential impact of a prevention-oriented nutrition intervention in other settings.

This evaluation is the first study to compare two approaches of targeting a Title II-MCHN program, and to do so with a full consideration of both the biological and behavioral issues around targeting the food assistance and BCC packages. The next two chapters present an overview of the context within which the program is implemented and the process used to design the full preventive approach and the strengthened recuperative approach. In doing so, they address some of the issues raised in this section about the optimal timing and duration of targeting under-twos as well as the behavioral issues to be considered in developing a BCC program targeting this age group.

1.4 Organization of the Report

This report is organized into 12 chapters that lead from the objectives and design of the study to the programmatic implications of the findings of the evaluation. **Chapter 2** presents the objectives and design of the evaluation, and discusses the role of different research activities within the overall evaluation. **Chapter 3** describes WV-Haiti's Haiti MCHN program and the services provided through the program. **Chapter 4** describes the process of designing the preventive and recuperative approaches and provides an overview of formative research and program development activities. **Chapter 5** provides an overview of the operations research process used to strengthen program delivery. **Chapter 6** describes the program participation patterns in the two program approaches as well as the characteristics of participants and nonparticipants. **Chapter 7** describes the primary nutritional impact of the preventive approach relative to the recuperative approach, while **Chapter 8** describes the impact of the program on behavior change outcomes, including maternal knowledge, trial, and adoption of key recommended practices as well as infant and child-feeding practices. **Chapter 9** presents results that demonstrate the impact of the program on household food security while **Chapter 10** describes indirect benefits of the program on household assets and maternal well-being measures. **Chapter 11** presents the relative cost-effectiveness of the two program approaches. The report concludes with **Chapter 12**, where key results are summarized, and their program, policy, and research implications are discussed.

2. OBJECTIVES AND DESIGN OF THE EVALUATION

2.1 Introduction

This chapter presents the objectives of the evaluation and an overview of the program theory behind this evaluation. It also describes how the various project activities were designed to address the intended objectives and how they fit with the theoretical framework that guided this evaluation. Finally, the chapter describes the collaborative engagement of the research teams from the International Food Policy Research Institute (IFPRI) and Cornell University with the implementation team from WV-Haiti.

2.2 Objectives of the Evaluation

The overall objective of the evaluation was to compare the impact and cost-effectiveness of the preventive and recuperative approaches of integrated nutrition and health interventions including a food supplementation component. The *specific objectives* were to:

- 1) Compare the impact of the preventive and recuperative approaches on the following outcomes among children aged 12-41 months:
 - a. Attained growth (mean WAZ, HAZ, WHZ, and their distributions);
 - b. Prevalence of undernutrition (stunting, wasting, underweight).
- 2) Compare the cost of the two approaches with respect to financial and human resources such as the amount of food required, staff training, and time.
- 3) Compare the cost-effectiveness of the two approaches, combining information from 1 and 2.

2.2.1 Additional objectives

In addition to the objectives mentioned above, the overall study also had the following additional objectives:

- 4) Assess differences between the two interventions in coverage of their respective targeted age groups (preventive: 6-24 months; curative: 6-60 months).
- 5) Document, with the use of operations research methods, differences between the two intervention groups in (1) the effectiveness of delivery of the various components of the two intervention packages; (2) the quality of the services provided; and (3) the institutional setup that appears to facilitate successful implementation.
- 6) Document, using qualitative research methods, the intrahousehold utilization and consumption of the food commodities, particularly consumption by the target individual.

- 7) Assist WV-Haiti in the design and implementation of a fully developed preventive approach to be compared with the recuperative approach. This will include designing new education messages that emphasize prevention of growth faltering, and designing a delivery mechanism to ensure the timely delivery of the messages to the targeted audience.
- 8) Assist WV-Haiti in reviewing and improving (if necessary) the set of education messages currently used in the recuperative approach.

This report presents detailed results that pertain to the main three objectives (impact, cost, and cost-effectiveness) as well the additional objective pertaining to the coverage of the program approaches. Objectives pertaining to the operations research, the design and implementation of the preventive and the strengthening of the recuperative approach are briefly described here, and have been elaborated in previous reports (Menon et al. 2002a, 2002b, 2005; Loechl et al. 2003a, 2003b, 2004). The objective pertaining to the use of the donated food commodities is partially discussed in the first round of operations research (Loechl et al. 2004). It is also addressed briefly in the chapter on child-feeding practices in this report.

2.3 Design of the Overall Evaluation

2.3.1 Program theory and pathways of influence

Specification and assessment of “program theory” pathways (Rossi, Lipsey, and Freeman 2004) is important for understanding how and why a set of program activities achieved (or did not achieve) their intended impact. This is particularly true for complex programs that can involve multiple intervention components, each of which place different demands both on program implementers and on program clients.

According to Rossi, Lipsey, and Freeman, program theory can be expressed as impact theory as well as process theory. Impact theory postulates the pathways through which an intervention is expected to achieve stated objectives. Process theory, on the other hand, is more concerned with the pathways that ensure that an intervention is implemented and utilized as designed. In the case of the WV-MCHN program, the impact theory is based on the tenets of the Title-II programming environment, where it is recognized that it is important to provide food assistance to improve family food security and increase access to fortified foods in the household. At the same time, it is also recognized that to reduce child undernutrition, provision of other inputs such as behavior change communication (BCC) are necessary to ensure a home environment conducive to good childcare practices and child growth. Thus, the impact theory for the WV-MCHN program in Haiti is that provision of food and nutrition education inputs can ensure good care practices and consumption of fortified food commodities, which in turn should ensure good nutrition.

The process theory for the WV MCHN program is based on the institutional demands and needs for implementing a Title II program providing both food assistance and BCC inputs. Working backwards from the most immediate program inputs, it is apparent that distributing food assistance requires capacity in monitoring beneficiary eligibility, ensuring smooth logistics, preventing leakage and corruption. On the other hand, ensuring good behavior change

communication requires staff who are trained in technical content as well as counseling and facilitation skills. In both cases, it is essential that other managerial and organizational processes are in place to ensure that workers are supported, motivated, and supervised appropriately.

Process theory can also be used to describe expectations regarding program uptake and utilization by intended beneficiaries. In the program approaches being evaluated, there are some common expectations about uptake and program use. Specifically, in both groups, the expectation is that all families with children under 60 months of age will use the RPs so as to receive general preventive health-care services. Then, for children identified as program beneficiaries, the expectation is that caregivers will fulfill their responsibilities to attend MCs and continue to bring their children to RPs.


Figure 2.1 provides a broad view of the “flow” of programmatic inputs from WV-Haiti to the child. This broad framework is intended to identify the contributing role of each step in this flow of activities to the achievement of the programmatic goals of improved child health and nutrition. It recognizes the variety of factors that influence program *delivery* as well as those that influence program *uptake* by caregivers and households in this context. In doing so, it demonstrates the complexity of the actions and actors that need to come together to lead to the ultimate intended outcome of the program, i.e., child health and nutrition.

(1) *WV-Haiti program management*: Factors that are important at this level are those aspects of the WV management that facilitated the design and implementation of the preventive and recuperative approaches, and the changes to the program implementation based on the operations research carried out in 2003. In particular, the management’s commitment and investment in ensuring high quality program implementation are important considerations at this level, as are general organizational structures, principles, and resources.

(2) *WV-Haiti program implementation level*: Staff-related factors that are likely to influence the efficiency and quality of program operations are important to consider at this level. Some of these factors include technical expertise, job motivation, supervision, etc. In addition to these, factors external to the frontline staff but internal to the program, such as resources needed for smooth operations (transport, medical supplies, fuel, interruptions in the food pipeline, etc.), can influence program operations. Similarly, factors external to the program, for instance, infrastructure and political disruptions, can also exert an influence on program implementation.

(3) *WV program operations and inputs*: At this level, we consider the Maternal and Child Health and Nutrition (MCHN) program inputs provided to the program beneficiaries. These include the food assistance package, as well as the health and nutrition knowledge provided by the BCC strategy and other WV MCHN program services. Finally, other WV programs in the evaluation areas, such as water, agriculture or sanitation, to name a few, can modify the influence of the MCHN “intervention package.” The program inputs are considered separately from the contextual issues described above that influence the delivery of these inputs.

Figure 2.1 Program impact pathways



(4) *Care context*: The program inputs provided by the WV MCHN program are expected to influence the context within which childcare occurs. Specifically, the program inputs can have an influence on community, household, and caregiver resources for the care of young children, including resources such as food availability, access and utilization, as well as knowledge, access to health care, etc. In addition, quality of program services and the rapport

between program and the communities can influence how communities and families attend and use program services. Consequently, they are expected to have an impact on how the program inputs are translated into care practices for children, and ultimately into child outcomes. The program inputs can change the resources available for childcare practices, as well as the nature of the relationship between the care resources and care practices.

(5) *Child impacts*: The impacts at this level are the final impacts of interest of the program evaluation. Indicators of child nutritional status (HAZ, WAZ, and WHZ) are the primary impact indicators of this evaluation.

2.3.2 Designing evaluation activities based on program theory

The nutrition intervention delivered through the WV MCHN program includes two major components: the food assistance component and the BCC component. Both of these components required different skills and capacities from program implementation staff. They also required different kinds of “compliance” from program beneficiaries. In short, the full implementation and uptake of the program requires a diverse set of actors to act in concert.

Recognizing this complexity, the entire evaluation process was designed to ensure that the key program components were *designed* from a “best practices” perspective and that they were implemented as best as possible. This was accomplished by engaging in a set of activities that informed the design and implementation of the program approaches. However, since the research team was not engaged in any implementation per se, all activities that pertained to program development and implementation were conducted through active engagement with WV-Haiti (see Figure 2.2). This resulted in a program that was designed and implemented as well as possible under the extant field conditions in Haiti.


From an evaluation perspective, the complexity of the program theory was captured through attention to assessment of each step in the program pathway. This was accomplished through the careful documentation of program implementation via operations research studies. In addition, the baseline and final surveys gathered detailed data on care resources and program utilization patterns from the survey communities. This ensured that all steps along the program pathway were documented well to aid in the interpretation and understanding of evaluation results.

Ensuring program components were designed well: The evaluation paid close attention to the design and implementation of the BCC strategy used in both program approaches by engaging in a solid formative research study as well as a consultative BCC strategy development process with WV-Haiti. Since WV-Haiti’s commodities tracking system was already designed and was operating well, the design process only involved discussions that resulted in the definition of the preventive and recuperative food ration packages.

Assessing and enhancing the implementation of program components: The implementation of the MCHN program, including the implementation of the food assistance and BCC components, was examined through a two-stage operations research process. Furthermore, with the recognition that program management and supervision have a strong influence on how

program activities are implemented in the field by frontline staff, the evaluation also examined the management structures of the program and the work context of frontline program staff.

Figure 2.2 An “engaged” model of evaluation and technical assistance



The information gathered on program implementation in the first operations research study was used to discuss implementation problems and concerns with the WV-Haiti management and field staff, which led to an internal program improvement process implemented by WV-Haiti. In a subsequent round of operations research, the research team assessed these

program improvements and also compared implementation between the two program approaches.

Assessing program uptake: The program evaluation activities were also grounded in the understanding that accessing and translating the WV-Haiti MCHN program inputs (food assistance and BCC) into child health outcomes requires substantial investments and attention by caregivers and households. This understanding was used to design the BCC strategy in ways that could encourage the best use of program resources for the child's well-being. Also, the *research* activities were designed to ensure that caregiver and household use of the program resources could be assessed to allow for a better and more nuanced understanding of how program inputs were translated into child health. Specifically, a variety of household and caregiver resources that could influence the use of program inputs were measured both at baseline and final surveys. Since some of these household and caregiver resources (particularly household food security and caregiver knowledge) could have been modified by the program activities, the evaluation process paid close attention to assessing them. In addition, the second round of operations research gathered qualitative data on participant adoption of the BCC recommendations. Finally, at the time of the final survey, a variety of data was gathered to evaluate the use of general program services as well as the use of the specific nutrition inputs, i.e., the food assistance and the BCC recommendations.

An overview of the program design and research activities that were employed in this evaluation is described in subsequent sections. Details about specific activities are presented in Chapters 3 through 11.

2.4 Project Activities

The study objectives were met through an engaged process with WV-Haiti that included two major components: *a program design* component that designed and developed the two program approaches in collaboration with WV-Haiti, and an *evaluation* component that compared the impact and cost-effectiveness of the two intervention approaches.

The *program and implementation plan design* component included a research stage that preceded the program development stage, and included the following activities:

A) Research

- i. A desk review of best practices for child nutrition and care, and current approaches and materials used in BCC programs in Haiti
- ii. An exploratory pre-survey qualitative study
- iii. A formative research study including a focused ethnographic study (FES) and household trials of improved practices (TIPs)
- iv. Two rounds of operations research to inform program implementation⁴

⁴ Note that the operations research studies were done as a sequence. However, the first operations research study was intended to assess the overall implementation of the newly developed program activities and use that information to correct any implementation problems. The second round, on the other hand, was conducted more to examine potential differences in implementation and utilization between the two program models.

- B) Development of the behavior change and communication (BCC) strategy and preventive approaches
- i. Identification of priority programmatic actions for the BCC component
 - ii. Development of the BCC strategy
 - iii. Development of BCC materials and training plans
 - iv. Training of field staff
 - v. Developing an implementation plan

The *evaluation* included four main components:

- A) An impact evaluation;
- B) A qualitative study, to guide the design of the baseline survey instrument;
- C) A round of operations research that assessed and compared program implementation and uptake between the two intervention groups;
- D) A cost-effectiveness study.

The research phase of the program design component was carried out largely by the IFPRI-Cornell team, while the program development phase was carried out with continuous engagement with the WV-Haiti team. Furthermore, although the research team from IFPRI and Cornell were highly involved in the design of the preventive approach and the development of the behavior change and communication strategy, all program implementation was done and supervised by WV-Haiti.

A broad timetable of the study activities is presented in Table 2.1.

Table 2.1 Summary of study activities

Year	2002	2003	2004	2005/2006
Design of program and implementation plan	Research <ul style="list-style-type: none"> ➤ Desk review ➤ Exploratory qualitative study ➤ Formative research study Development <ul style="list-style-type: none"> ➤ Identification of priority actions for BCC ➤ Development of BCC strategy ➤ Development of materials and training plan ➤ Training ➤ Development of implementation plan 	<ul style="list-style-type: none"> ➤ Implementation of both the preventive and recuperative approaches ➤ Operations research (Round 1) 	<ul style="list-style-type: none"> ➤ Program improvement consultation (with WV) 	
Evaluation activities	<ul style="list-style-type: none"> ➤ Pre-survey qualitative research ➤ Baseline survey 		<ul style="list-style-type: none"> ➤ Operations research (Round 2) ➤ Cost study (mid-term) 	<ul style="list-style-type: none"> ➤ Cost study ➤ Final survey

The remainder of this chapter briefly summarizes the evaluation activities. Specific objectives, methods, and results that pertain to each of the components are described in Chapter 5 (operations research), Chapters 6 through 10 (results of the impact evaluation), and Chapter 11 (cost-effectiveness study results).

2.4.1 Evaluation activities

The four evaluation components are described below.

2.4.1.1 Impact evaluation

The evaluation used a community-level randomized pre-post design, whereby 10 paired clusters of communities were randomly assigned to either the preventive or the recuperative program group. It was designed to use a probability design while providing enough evidence on the plausibility of the impact as to inform future evaluations of this nature, as well as future programs that use this approach (Habicht, Victora, and Vaughn 1999).

The main outcomes of the evaluation were mean height-for-age Z-scores (HAZ), weight-for-age Z-scores (WAZ), and weight-for-height Z-scores (WHZ), and the prevalence of childhood stunting, underweight, and wasting.

The baseline survey was conducted between May and September 2002 and the post-evaluation survey was conducted exactly three years later, between May and September 2005, to minimize seasonal variations. All components of the intervention packages, except the newly developed BCC strategy, were implemented immediately following the baseline survey, i.e., in August-September 2002. The full BCC package, however, was implemented several months later (in May 2003) due to delays in design, staff training, and implementation.

Twenty clusters of communities, each attended by one Health Agent (WV-Haiti local staff in charge of program activities), were selected for the evaluation from WV-Haiti's program areas. Each cluster was paired with another one selected to be similar in location (e.g., distance to the main highway and/or main town), geographic and ecologic conditions (e.g., whether located in the plains or the mountains), access to a health-care center, and the existence of a WV-Haiti private sponsorship program (Area Development Program) (Menon and Ruel 2003). Within each pair of clusters, one was randomly assigned to the preventive approach and the other one to the recuperative approach. Thus, the unit of randomization was the cluster of communities covered by one Health Agent.

The number of clusters (20) was determined by balancing the cost of conducting the surveys with the need to have enough clusters to be able to achieve the desired sample size of 750 children per program approach. A larger number of clusters, with fewer second-stage-sampling units (in this case, households) within each cluster is usually preferable, but the cost of including more clusters is higher than that of including households within clusters.

The baseline survey was used primarily to examine the differences between the program communities at baseline, and by doing so, to assess the success of the randomization. The impact assessment was based on comparisons between the two approaches at the final survey.

Further details about the target age groups for the surveys, sample sizes, and other design issues are provided below.

2.4.1.1.1 Sample size estimation

Sample sizes were estimated using an equation for estimating sample sizes for differences in proportions between the preventive and recuperative groups (Cohen 1988). The necessary sample sizes to examine differences in prevalence rates are the same as those necessary to examine differences between distributions, and they are larger than those necessary to examine differences in means (Brownie, Habicht, and Cogill 1986). The effect size (magnitude of differences between intervention groups at post-intervention) used for the calculation was based on previous studies of the effect of supplementation on child growth, which ranges from 0.25 to 0.46 Z-scores for WAZ and 0.04 to 0.35 Z-scores for HAZ (Caulfield, Huffman, and Piwoz 1999). An improvement of +0.35 Z-scores of nutritional status in a population with average Z-scores around -2.0 will result in a decline in prevalence of undernutrition from 50% to 38% (a decrease of 12 percentage points).

Since the intervention in this case was randomized at the cluster level rather than at the level of the individual child, it was necessary to account for the clustering of characteristics within a cluster (called the design effect⁵). This must be taken into account when calculating sample sizes because it increases the sample size needed when the intervention is randomized at the cluster, rather than the individual level. Since information on design effect was not available at the time of sample size calculations, we assumed a design effect of 1.5 based on discussions with WV-Haiti and others working in Haiti about the homogeneity of communities in the evaluation area.

We estimated a sample size of 75 children per cluster, for a total of 1,500 children. This sample size provided the ability to detect differences between groups in the final survey of 7.5 percentage points or larger in the prevalence of stunting, assuming an average design effect size of 1.5 (clustering of characteristics within cluster), an alpha of 0.05, and power of 0.90. This sample size also provided the ability to detect differences larger than 7.5 percentage points in underweight, 5 percentage points in wasting, and differences larger than -0.2 in mean Z-scores for HAZ, WAZ, and WHZ.

2.4.1.1.2 Age group selected for impact assessment

Children 12-41 months of age were selected for the impact assessment, based on available scientific evidence on the age of greatest nutritional vulnerability and largest potential for response to nutritional interventions. This estimation was based on research showing that nutrition interventions (such as protein energy supplementation) are more effective in improving growth if provided as early as possible before the child reaches 24 months of age (i.e., from 6 months on) (Schroeder et al. 1995), and if sustained for at least 12 months (Rivera and Habicht

⁵ The design effect is the ratio of the variance for the cluster sample divided by the expected variance of a simple random sample of the same size. Since the design effect is dependent on the variance between cluster, it will be smaller if the number of clusters is large and the number within each cluster is small (Foreman 1991). For complex nutrition surveys, it has been shown that clusters with 30 children in each cluster lead to design effects for stunting (HAZ %<-2) that range from 0.44 to 2.13 and 1 to 1.62 for underweight (defined as W/A < 60%) (Katz 1995).

2002). Moreover, growth patterns of Haitian children, which are similar to worldwide patterns documented for children living in impoverished environments, show the greatest rates of growth faltering during the period between 6 and 17 months of age (EMMUS-III 2001; Shrimpton et al. 2001).

Children considered most likely to benefit from the preventive approach were those who were first exposed to the supplementation between 6-11 months of age, and for the whole duration of their period of greatest vulnerability (i.e., up to 24 months of age). These children would be 24-41 months old at the final survey. In addition, we also included children 12-23 months, who were only partially exposed (i.e., had not yet reached 24 months at final survey). For the recuperative approach, the 12-41 months age range was also expected to include mostly children who had already been exposed to the program, with some possible truncation among children in the younger age range, given that the peak prevalence of underweight children in Haiti is 12-17 months (EMMUS-III 2001).

As noted above, program implementation started immediately after the baseline survey, except for the new BCC strategy, which was implemented 9 months later. This meant that children who were 36-41 months at final survey were not fully exposed to all program components in either program model. Thus, the sample at final survey includes children 24-35 months of age who were fully exposed to the preventive program, and two groups of partially exposed children (12-23 months; and 36-41 months).

In 2002, the survey sample included 801 children in the preventive group and 801 children in the recuperative group, for a total of 1,602 children between 12 and 41 months of age. In 2005, data were collected on 749 children in preventive and 751 children in the recuperative group, yielding a total of 1,500 children between 12 and 41 months of age in both groups.

2.4.1.1.3 Survey design

The baseline and final surveys included both a household and community questionnaire. The household questionnaire was administered to the mother of the index child⁶ (referred to as the caregiver), whereas the community questionnaire was administered using a group interview methodology with key community members, such as religious leaders, medical staff, or schoolteachers.

The community questionnaire provided data at the community level that allowed a comparison of the different clusters after the randomization process was completed. Community-level data were gathered on the smaller community units (called *localités*) within each cluster, because these were more meaningful and finite geographic entities than the clusters themselves. Information was gathered on access to the nearest major town, the main activity of the residents, key geographic characteristics, access to public services such as schools and markets, and access to health services such as hospitals, dispensaries, and the services provided by the WV-Haiti program.

⁶ The “index child” is defined as the 12-41 month-old child of interest for assessment of impact on nutritional status.

The household questionnaire gathered data on household characteristics such as household composition, socioeconomic status, and food insecurity; and on caregiver characteristics such as education, childcare knowledge and experience, women's empowerment and involvement in decisionmaking, and mental and physical health. It also gathered data on six types of childcare practices: (1) early infant feeding practices (i.e., around birth); (2) current child-feeding practices of index child including meal frequency, dietary diversity, intake of animal sources foods; (3) child feeding practices for younger sibling, (4) preventive and curative health-care seeking practices (e.g., antenatal care, childhood immunization, treatment for diarrhea); (5) hygiene practices (e.g., child, maternal, and house cleanliness); and (6) discipline practices (see list of baseline and final survey modules in Annex 2.1).

Anthropometric measurements (height and weight) were taken on the caregivers, the target child (12-41 months of age), and his/her younger sibling if any. Weight was measured using a UNICEF Seca scale (SECA Ltd., Birmingham, U.K.) accurate to 0.1 kg, and height was measured using wooden length boards accurate to 1 mm (Irwin Shorr Productions, Olney, Maryland, U.S.A.). Weight and length were measured by fieldworkers who were standardized using recommended protocols.

Child health and appetite were assessed using a visual analogue scale, where caregivers are asked to rate the health (or appetite) of her child as compared to other children his or her age and indicate where it fell on a 10-cm-long line drawn on the questionnaire. Caregivers were also asked to recall whether the child had symptoms of illnesses (diarrhea, cough/cold, pneumonia, or fever) in the two weeks prior to the survey. Finally, spot-check observations were used to assess the cleanliness of the child, caregiver, and the household.

2.4.1.1.4 Selection of survey households

The survey households were selected after conducting a complete census of the general population residing in the communities included in the program evaluation. A pool of households eligible for inclusion in the survey was generated for each cluster by identifying those households that had a child in the target age range, where the child's mother also resided in the same house. Eighty-five households were selected at random (without replacement) from each cluster, and the survey administered to the first 75 of these households. When possible, the field team conducted more household interviews than the minimum 75 required to provide a larger sample size within the existing logistical constraints of the survey.

The census gathered information on household composition, headship, and the date of birth of all children under 5 years of age. For all children under 5 and for women of reproductive age who were identified as being either pregnant or lactating, data were gathered on whether they had ever been program beneficiaries.

2.4.1.2 Qualitative study

An exploratory qualitative study was carried out before implementing the baseline survey to assist in the design of the quantitative survey questionnaires. This study was carried out in communities outside of the study area and provided the basis for designing the questionnaires

used in the survey. More information on this study is provided in Chapter 4 and in Menon et al. 2002a.

2.4.1.3 Operations research studies

Operations research (OR) methodologies, including both qualitative and quantitative approaches, were used to gather information on the effectiveness of delivery of the interventions in the communities included in the evaluation. Although the first round of OR was intended to inform program implementation, it is described here along with the second round (which was more pertinent to the impact evaluation), since both rounds of OR were linked and shared the same broad objectives. These were to (1) assess program implementation and quality; (2) improve quality of operations and fidelity to program approaches and plans; and (3) gain sufficient knowledge about implementation issues and quality in the two program approaches to allow appropriate interpretation of impact evaluation results.

Two rounds of OR were undertaken, with the first occurring in July-September 2003, shortly after full implementation of the BCC strategy, and the second approximately one year later, in June-August, 2004. The first round (2003) focused on the first broad objective and aimed to assess the fidelity of implementation of the program relative to plans, assess the quality of delivery of the various services, and explore perceptions of stakeholders (i.e., participants/beneficiaries⁷ and field implementers) regarding the operations and quality of services provided by the program.

This first round of OR (OR1) was followed by a process of consultation with program implementers, including management and field staff. The consultative process was directed toward the second broad objective: through this process, program operations were improved and better aligned with program design.

After allowing approximately 5 months for staff to implement corrective actions/program improvements, a second round of OR was undertaken, with the objectives of assessing implementation of the corrective actions/program improvements identified during the consultative process. Notably, this round also served to document differences between the two program approaches in program implementation, staff work context-related factors (job motivation, supervision, etc.), and finally, beneficiary mothers' knowledge and experiences with trial and adoption of specific infant and young child-feeding and care practices promoted in the BCC, in order to gain understanding of constraints to adoption of these practices in the two program areas. By doing so, the OR process enriched the impact evaluation through identifying any relevant differences between program areas in implementation, staff-level factors, or participants'/beneficiaries' experiences.

2.4.1.4 Cost-effectiveness study

Two cost studies were conducted to assess the cost and cost structures of the two program approaches, one in 2003-2004 (Maluccio and Loechl 2004) and the other in 2005 (Maluccio and Loechl 2006). In evaluating costs, the cost studies considered direct program costs, off budget

⁷ "Participants" include those who access any of the available range of services; "beneficiaries" are those who live in households receiving food assistance.

program costs, as well as private (beneficiary household) costs. Knowledge of the structure of costs helped identify which components of program costs were truly different between the program approaches, thus, which were important to consider in the estimates of cost-effectiveness.

The difference in costs between the program approaches was applied to the difference in program outcomes at the final survey to calculate the relative cost-effectiveness of the two targeting approaches. Details of the cost-effectiveness calculations, the various considerations that went into the calculations, as well as the implications of the results, are discussed in Chapter 11.

2.5 Conclusions

This chapter presented an overview of the evaluation process, focusing on the consideration of program theory in the design of the evaluation activities. In subsequent chapters, we discuss how the program theory was translated into program design activities as well as the specific evaluation activities, such as the operations research and the impact evaluation.

3. WV-HAITI'S MCHN PROGRAM

3.1 Introduction

This chapter provides an overview of the various services provided by the WV-Haiti MCHN program in the Central Plateau of Haiti (see map below). It describes the services and their main delivery points and briefly describes the organizational and staffing structure of the program.

3.2 Services Provided

As with many other USAID Title-II funded MCHN programs, the program implemented by WV-Haiti in the Central Plateau of Haiti provides a range of MCHN services. These services are targeted to pregnant and lactating women as well as to children between 0 to 5 years of age. Many of the services offered are open to all members of the community in these groups, while some other services are provided in a targeted fashion. Food assistance, for instance, is targeted, and is provided on a conditional basis, i.e., program beneficiaries are required to use other preventive services in order to receive monthly food rations. Box 3.1 presents a summary of the program service delivery points (these are also described in more detail in section 3.3) and Table 3.1 provides an overview of the general and targeted services provided by WV program.

Box 3.1 Five service delivery venues for the WV-Haiti food-assisted MCHN program

Rally Posts (RPs): Health education, growth monitoring and promotion (GMP), immunizations, and other preventive health-care services are provided to all participants, and program beneficiaries are identified.

Mothers' Clubs (MCs): Small peer groups of beneficiary mothers (or mothers of beneficiary children) gather for facilitated discussions of health and nutrition topics laid out in the program's behavior change and communication (BCC) strategy and curriculum.

Food Distribution Points (FDPs): Venue for distribution of monthly food rations to beneficiaries.

Pre- and postnatal consultations: Preventive health care and education are offered.

Home visits: WV health staff provide additional education and referrals to mothers of newborn infants, severely malnourished children, and children whose growth has faltered.

The research was undertaken in the Central Plateau region, where WV operates in all 12 communes and serves a population of approximately 600,000. The evaluation covered three communes: Hinche, Thomonde, and Lascahobas.

Table 3.1 Services provided by the WV MCHN program in Central Plateau

Group	Services available to all	Targeted services to program beneficiaries	Conditional services^a
(1) Pregnant women	<ul style="list-style-type: none"> - Prenatal consultations and check-up - Tetanus toxoid immunizations - Iron-folate supplements - Mothers' Clubs 	<ul style="list-style-type: none"> - Home visits (in late pregnancy) 	<ul style="list-style-type: none"> - Food assistance
(2) Lactating mothers with children 0-6 months of age	<ul style="list-style-type: none"> - Postnatal consultations and check-up - Postpartum vitamin A supplements (via home visits) - Mothers' Clubs 	<ul style="list-style-type: none"> - Home visits (targeted to mothers of newborn children) 	<ul style="list-style-type: none"> - Food assistance
(3) Children 6-59 months of age	<ul style="list-style-type: none"> - Immunizations - Vitamin A capsules - ORS - Deworming - Growth monitoring and promotion - Group education (at Rally Posts) 	<ul style="list-style-type: none"> - Mothers' Clubs (targeted differently in the preventive and recuperative approaches, see chapter 4) - Home visits (targeted to severely malnourished children) 	<ul style="list-style-type: none"> - Food assistance
(4) Women 15-49 years old	<ul style="list-style-type: none"> - Family planning 		

^a Food assistance is conditional upon beneficiaries attending monthly Rally Posts (children) or pre-natal (pregnant women)/postnatal (lactating women) clinics and monthly Mothers' Clubs.

Map of Haiti



Source: <https://www.cia.gov/library/publications/the-world-factbook/geos/ha/html>.

3.3 Program Service Delivery Points

As highlighted in Box 3.1, the WV MCHN program offers services at five major points of contact between program staff and participants. These are (1) *Rally Posts* (RPs), where health education, growth monitoring and promotion, and preventive health care are provided and beneficiaries are identified; (2) *Mothers' Clubs* (MCs), where smaller groups of participants gather to discuss health and nutrition topics in the context of the program's behavior change and communication (BCC) strategy; (3) *Food Distribution Points* (FDPs), where beneficiaries receive their monthly food rations; (4) *Pre- and Postnatal Consultations*, where pregnant and lactating women receive preventive health care and education; and (5) *Home Visits*, where beneficiary households with a newborn infant, a severely malnourished child, or a child with growth faltering are visited by the WV health staff. A brief description of the services offered and operational plan at each of these service delivery points is provided below.

3.3.1 Rally Post services

Rally Posts (RPs) are open to all pregnant women, mothers with children less than 5 years of age, and women 15 to 49 years old in the communities attended. Services provided include health and nutrition education, growth monitoring and promotion of children under 5 years of age, immunization, vitamin A supplementation, deworming, free distribution of ORS, and information about the family planning component.⁸

The RPs are the main entry point into the program and are used to identify beneficiaries and to refer them to the appropriate program services. New beneficiaries are identified at the RPs every month, and new eligible children are admitted on a monthly basis. For pregnant and lactating women, however, new entrants are invited into the program only every four months.

Monthly weighing of beneficiary children and monthly attendance at the RP by the child's caregiver are mandatory for receiving food assistance. Either the mother or another caregiver can take the child to the RP.

RP meetings are held on a monthly basis in each community and are managed primarily by the health agent responsible for that community. The health agent is usually assisted by at least two other health agents and two *colvol*s. The sequence of activities described in the implementation plan is as follows (also see Figure 3.1 below):


- A) The education session is the first activity to be carried out at the RP. The sessions run for about 10 minutes, depending on the topic. WV sets up a calendar of monthly topics, which include the following topics: immunization, pre- and postnatal care, preparation for child delivery, diet for pregnant and lactating mothers, weaning techniques, description of kwashiorkor and marasmus, hygiene and environment, diarrhea and preparation of oral rehydration salts, acute respiratory infections, family planning, and HIV prevention. It is expected that appropriate visual materials will be used to communicate more effectively. The health agent or *colvol* is also expected to

⁸ WV-HAITI offers hormone pills and three monthly injections. Women can receive these services administered by WV nurses at mobile clinics, in health centers during pre- and postnatal consultations, or at Area Development Program clinics.

conduct several sessions on the same topic throughout the day to allow all mothers to participate and to restrict the size of the group to 10-15 persons.

- B) After the education session, the health agent registers attendance of each participant. The health agent determines whether the participant is due to receive immunization, vitamin A supplementation, and/or deworming tablets. She also updates the health cards with the information on immunization and vitamin A supplementation, and for food aid beneficiaries, she signs attendance on the ration card. This same health agent also does the distribution of vitamin A (every six months for children less than 5 years of age), anti-helminths tablets (every six months to children 2-5 years of age), and oral rehydration salts (ORS) sachets (three sachets per month per household).
- C) The next activity at the RP is growth monitoring and promotion. Each child is weighed and the weight is recorded on the growth chart printed in the health card kept by the caretaker. It is expected that at least the caregivers of malnourished children, especially of moderately and severely malnourished children (M2 or M3 for weight-for-age according to the Gomez classification), will receive brief counseling about feeding practices and prevention of childhood illnesses.
- D) After growth monitoring and promotion, children are directed to receive their immunizations. Children receive vaccinations based on their age and previous immunization history (previously verified by the health agent in Step 2 above).

Figure 3.1 Flow of activities at the Rally Post



3.3.2 Mothers' Club services

Originally, the Mothers Clubs (MCs) were designed as a venue for small peer group discussions between program health staff and beneficiary mothers on general health and nutrition topics. Monthly attendance at the MCs is a requirement for receiving food assistance. Unlike the RPs and the food distribution points, mothers are required to attend the MCs themselves and cannot send another family member to use this service. This ensures that mothers are the direct participants in the education activities offered at the MCs.

Results from our initial formative research (Menon et al. 2002b) suggested that the MCs would be an ideal setting for the more comprehensive behavior change and communication (BCC) strategy that the research team in partnership with WV staff was developing. Reasons for

this are that the MCs are located close to the mothers' homes and include only a small group of participants, resulting in minimal distraction (especially compared to the RPs). Thus, the WV program decided to use the MCs as the primary venue for BCC activities.

The MCs are implemented as monthly small peer group discussion sessions, facilitated by health workers trained in technical aspects of nutrition and child health as well as in adult education techniques. Chapter 4 describes the re-organization of MCs following the formative research and related development activities, and provides more detail on how the MCs' organization differs between preventive and recuperative approaches.

3.3.3 Food Distribution Point (FDP) services

The distribution of food aid commodities to the MCHN beneficiaries of the WV program occurs on a monthly basis at special distribution points. Beneficiaries from several communities (*localités*) are scheduled to receive their food rations at a central distribution point on the same day. The distribution is done by community and starts with beneficiaries from the communities that are far away from the distribution point. Within each community food rations are provided first to lactating and pregnant women and then to caregivers of child beneficiaries.

There are a total of 10 central FDPs covering 50 RPs in the area included in the research (including preventive and recuperative program communities). Unlike at the MCs, a beneficiary can designate another family member to collect the food ration by handing over the beneficiary card to this person.

The beneficiary households receive both direct (individual) and indirect (household) rations. The amounts and commodities vary with respect to the beneficiary category (see Table 3.2). Even if a household has two direct beneficiaries participating in the program, only one indirect ration is provided. The caloric content of the rations is also indicated in the table below.

Table 3.2 Composition of direct and indirect food rations, per beneficiary category

Type of commodity	Children 6-23 months of age (preventive approach) Undernourished children 6-59 months of age (recuperative approach)		Pregnant and lactating women (both approaches)	
	Direct child ration (kg)	Indirect household ration (if child is beneficiary) (kg)	Direct women ration (kg)	Indirect household ration (if mother is beneficiary) (kg)
WSB	8			
SFB		10	5	5
Lentils		2.5	2	2
Vegetable oil	2		1.5	1.5
Caloric value (kcal/day)	1,534	1,414	1,233	1,233
Total caloric value	2,949		2,467	

The sequence of activities at the FDPs is as follows:

- A) Eligibility of the beneficiary is verified by food monitors and health agents mainly based on the information on the beneficiary card. This card contains information about the beneficiaries and indicates attendance by the beneficiary at the other MCHN activities (i.e., Mothers' Clubs, Rally Posts, and pre- and postnatal consultations) that are required in order to receive the food rations.
- B) Once their eligibility is verified, the beneficiaries proceed to collect their rations. A team of trained beneficiaries assists the WV staff during the distribution and is responsible for opening food sacks, measuring out appropriate amounts of each of the commodities, and handing over the food to beneficiaries.
- C) Finally, once the beneficiary (or designee) has received his/her rations, the food monitor verifies the rations received and the beneficiary card. In some cases, the food monitors re-weigh the rations to verify that the right amounts have been given to the beneficiary. Once this final check is complete, the food monitor signs the beneficiary card to indicate that the correct ration has been delivered to that beneficiary.

3.3.4 Pre-and postnatal consultations

Prenatal consultations are provided to all pregnant women in the WV program areas who decide to use the WV services. Typical prenatal services are provided and include physical examinations, provision of iron-folate supplements, tetanus toxoid immunizations, etc. Attendance at prenatal consultations is mandatory in order to receive food rations.

Postnatal consultations are also provided to all women who have recently given birth. The key services at the postnatal consultations include physical examinations for the mother and the newborn infant, provision of postpartum vitamin A (if the mother did not already receive this via a home visit soon after delivery), and other health services. As with the prenatal services, attendance at postnatal consultations is mandatory in order to receive food rations.

The pre- and postnatal consultations are provided by WV nurses, usually at the WV clinics or dispensaries or at mobile clinics in the more remote areas of Central Plateau.

3.3.5 Home visits

WV health staff make home visits in the areas covered by the program services to check on women in late pregnancy, mothers of newborn infants, or to follow up on severely malnourished children. The services provided at home visits usually include only a visit by the health agent or nurse, except for mothers of newborn infants who also receive a dose of postpartum vitamin A in the first home visit after delivery.

3.4 Program Organization/Staffing

The program services described in the preceding sections are provided by two teams of staff: all health services are provided by a team of health staff, while all food distribution

activities are conducted by a team of Commodities staff. The organizational structure for both of these staff teams are provided in Annexes 3.1 and 3.2. The roles of the main WV-Haiti program implementers and the staff organizational structure are outlined below.

WV-Haiti health promoters and assistant health promoters (previously called health agents and *colvols*, respectively). The health promoters (HPs) and assistant health promoters (AHPs) are the direct implementers of the MCHN program in the field and thus are the frontline staff in contact with the program participants. They are in charge of the interventions being delivered at the RPs and MCs and of assisting the food monitors at the FDPs. They attest to the attendance by beneficiaries at the different program activities (which determines their eligibility to receive food rations). HPs are WV employees and receive a monthly salary. AHPs, previously called *colvols*, used to be community volunteers who received only a small incentive from WV. However, with the growth of the program over time, it became apparent that *colvols* were in fact more than volunteers in terms of their time commitment to the program activities, and their level of responsibilities. In 2003, they were promoted to AHPs and received an increase in salary. They also started to participate in all training for health staff.

WV-Haiti food monitors. Food monitors are responsible for the distribution of the food rations at the FDPs and for ensuring that only eligible beneficiaries receive the food. They are also responsible for verifying the beneficiary lists provided by the health team. This is done mainly by consulting the beneficiary card.

WV-Haiti health (MCHN) and commodity supervisors at the local level. MCHN supervisors are nurses responsible for the supervision of the health agents and *colvols*, and commodities field supervisors are responsible for the supervision of the food monitors. The MCHN supervisors generate monthly lists of beneficiaries eligible to receive the food rations, based on lists of attendance prepared by the health agents for each service delivery point. The Commodity Section uses these lists to program the food amounts needed per distribution point.

WV-Haiti management staff at the national and regional levels in MCHN and Commodity. The Assistant of the Regional Health Coordinator in Hinche supervises the nurses. This assistant is, in turn, supervised by the Regional Health Coordinator. The National Health Coordinator for WV is based in Port-au-Prince and oversees the activities in all the program areas of WV in Haiti. The organizational structure for the health component of the program is presented in Annex 3.1.

The commodity supervisors work under the supervision of the Assistant of the Regional Commodity Officer. This Assistant is, in turn, supervised by the Regional Commodity Officer in Hinche. The organizational structure for the food component of the program is outlined in Annex 3.2.

In the evaluation areas, the health team consisted of 4 nurse supervisors, 20 health promoters (HPs), and around 20 assistant health promoters (AHPs). The Regional Coordinator supervises the nurse supervisors as well as other nurses who provide services in other communes of the Central Plateau region. Each HP-AHP pair provides services to about 300 children. The remoteness and dispersed nature of households in this region of Haiti necessitate lower health worker-client ratios than in other more dense regions.

The commodities team provides services for the entire Central Plateau DAP area, with no distinction between staff who serve the preventive and recuperative areas.

3.5 Conclusions

This chapter described the basic service provision structure of the WV MCHN program in the Central Plateau. The next chapter describes how the program design steps outlined in Chapter 2 were used to design the preventive program approach, strengthen the recuperative approach, and design the BCC strategy to be used in the MCs for both approaches. The subsequent chapter provides a detailed view of how the two rounds of operations research were used to assess and improve the services delivered at the venues described in this chapter. Together, these chapters provide a solid example of how applied program research can be used in a collaborative fashion with implementation staff to design and implement high quality programs.

4. GETTING IT RIGHT FROM THE START: DESIGNING THE PREVENTIVE AND RECUPERATIVE APPROACHES⁹

4.1 Introduction

This section describes the process used to develop the preventive program and to strengthen the recuperative program. The two main sets of interventions that needed to be adapted in order to design a truly preventive approach were the food distribution (targeting and duration of intervention) and the education sessions at the Mothers' Clubs. It was agreed at the onset of the study that one of the most critical components of a good preventive or recuperative food-assisted MCHN program was an effective behavior change and communication (BCC) strategy. The research and development process involved in helping WV-Haiti design and plan the implementation of a new BCC strategy for its program is described in this chapter.

The design of the food assistance component for both program approaches is described in the next section, followed by the research and development process used to design the BCC strategy. Implementation protocols were already in place for the preventive health-care components of the program at the Rally Posts (i.e., immunization, vitamin A supplementation, and deworming) and for pre- and postnatal consultations, and these were not modified (see Chapter 3). These activities are therefore not discussed in this chapter, as is the case for the home visit protocols. Thus, the chapter focuses on the screening of beneficiaries for food assistance (at the RPs), the food distribution protocols (implemented at FDPs), and the BCC strategy mostly implemented at the MCs. The chapter concludes with a description of the similarities and differences between the preventive and the recuperative program approaches being compared in this study.

4.2 The Food Assistance Component

The protocol for food assistance for the recuperative program remained as originally designed by WV-Haiti. For this approach, *malnourished* children (identified during growth monitoring)¹⁰ are enrolled in the program between 6-59 months of age for 9 months. According to WV-Haiti management, the rationale for providing food supplements to malnourished children for 9 months is based on programmatic experience, which suggests that 9 months is sufficient for most children to recover from malnutrition. Note, however, that to our knowledge, this assumption has not been verified by evaluation research. Furthermore, while anecdotal evidence suggests that the duration of participation for children in Title II programs is between 6 and 12 months, there are no systematic reviews available of the average duration of enrollment of children in recuperative programs implemented by other Cooperating Sponsors.

Design of the food assistance component for the preventive program was based on empirical research showing the benefits of providing nutrition interventions early in life in order to *prevent* malnutrition. Discussions were held with program managers at WV to review the

⁹ Cornelia Loechl led the writing for this chapter.

¹⁰ Malnourished children are defined as M2 and M3 according to the Gomez classification. In this classification, normal (N) corresponds to $\geq 90\%$ of the median of the weight-for-age CDC/NCHS/WHO standards; mild malnutrition (Grade M1) to 75% to $< 90\%$; moderate malnutrition (Grade M2) to 60% to $< 75\%$; and severe malnutrition (Grade M3) to $\leq 60\%$ (Cogill 2003).

scientific rationale for a preventive approach and discuss how the benefits of food supplements could be maximized, while remaining within the resource capacities of the program. It was decided that children would be enrolled in the preventive program between 6-18 months of age. This age range was selected because research suggests that the period between 6 and 24 months is the age of maximum response to nutrition interventions (see Chapter 2, section 2.4.1.1.). Beneficiaries would continue to receive food supplements up to the age of 24 months, thus ensuring that even those who enter the program as late as at 18 months of age would receive 6 months of supplementation.

Food assistance for pregnant and lactating women also remained as originally designed by WV, i.e., pregnant and lactating women are eligible to receive food assistance for 6 months during pregnancy and for the first 6 months of lactation. Food assistance to pregnant and lactating women was to be exactly the same in preventive and recuperative program communities.

Information regarding ration size and composition - for direct and indirect rations - is provided in Chapter 3.

4.3 The BCC Component

The rationale for implementing a BCC strategy in the context of a food-assisted MCHN program is that, in addition to providing food for children, it is also important to ensure that mothers are well informed and able to adopt optimal child-feeding and caregiving practices. The key aspects of care and feeding that are particularly important to *prevent* malnutrition during the vulnerable period of 0-23 months of age are breastfeeding, appropriate complementary feeding practices, and other preventive and curative health-related practices like good hygiene, timely immunization, appropriate home health care, and care-seeking during illness. Mothers also need to be educated about how to address childhood malnutrition and how to detect signs of severe malnutrition or childhood illness. These topics were the focus of the BCC strategy developed for WV-Haiti's preventive and recuperative programs.

The strategy was designed in two phases: a research phase and a development phase (see Figure 4.1).

In the following section, details of each step of both phases are presented. The data from the research phase were used to assess the adequacy of current infant and young child-feeding practices and to identify constraints and facilitating factors to adopting optimal feeding practices. Following this, programmatic actions to improve nonoptimal practices and sustain optimal practices were developed consultatively with WV-Haiti staff. This was achieved through the different steps of the program development phase.


4.3.1 Research phase

4.3.1.1 Review of existing communication materials

The first review of program communication materials commonly used by PVOs in Haiti was initiated in November-December 2001 and has been reported previously (Menon et al. 2001).

The purpose was to gather information on the existing nutrition and health education models currently used in Haiti. Two guides used by different PVOs in Haiti were reviewed. It was found that both guides covered breastfeeding practices and practices related to the prevention and treatment of childhood illnesses quite extensively. Messages related to complementary feeding practices, however, were minimal and psychosocial care was generally absent from both education packages. Messages related to complementary feeding focused mainly on nutrient density and dietary diversity and did not address feeding frequency or portion size.

Figure 4.1 Steps of the design process of the BCC component¹¹



4.3.1.2 Preliminary qualitative study

As a first step in the formative research process, a rapid qualitative study was conducted in January 2002 in the Central Plateau to gather information on general patterns of infant and child-feeding practices. The data were used for two purposes: (1) to design the formative research required for the development of the BCC strategy (Menon et al. 2002a) and (2) to guide the development of the baseline survey for the evaluation (see Chapter 2). The results suggested that the current infant and child-feeding practice departed substantially from international feeding recommendations, especially with regard to the recommendations to exclusively breastfeed infants up to 6 months of age, and to complement breast milk with frequent feeding of energy- and micronutrient-dense complementary foods after 6 months of age.

¹¹ All BCC communication materials and information about the development of the program are available at the following FANTA website: www.fantaproject.org.

4.3.1.3 Formative research study

A more extensive formative research study was undertaken between May and August 2002. The objectives of the study were (1) to gather in-depth information on current infant feeding practices and their determinants; (2) to identify constraints and factors that may facilitate adoption of optimal feeding practices; (3) to conduct recipe trials to develop improved complementary foods based on the use of donated, fortified food commodities and other nutrient-rich foods locally available; and (4) to understand current WV program activities in the Central Plateau of Haiti and identify potential program delivery points that could be used for the behavior change and communication (BCC strategy) (Menon et al. 2002b).

Several data collection techniques were used, including individual and group interviews with mothers of young infants, grandmothers, fathers, and WV program staff. Participatory group recipe trials with groups of local women were conducted to develop recipes for enriched complementary foods and to discuss their feasibility, acceptability, and affordability under real-life conditions in the program areas. Finally, observations of WV's program activities were conducted in the Central Plateau area as well as on the island of La Gonâve to understand the implementation of different program components and to explore the feasibility of enhancing ongoing educational activities. Key results of the formative research are summarized below.

Infant and young child-feeding practices: The formative research provided information that allowed us to characterize typical infant and young child-feeding practices in rural Haiti and to understand the rationale for these practices. Specific factors likely to either facilitate or constrain adoption of optimal practices were also identified for each specific dimension of child-feeding practices studied.

Development of enriched complementary foods: The recipe trials confirmed that traditional complementary foods are low in micronutrient-density, although they are generally of adequate energy density. The process also demonstrated that it was feasible for the recipe trial participants to develop a number of improved recipes using traditional preparation methods, fortified donated commodities, and adding locally available nutrient-dense foods such as fish, eggs, beans, and vitamin A-rich vegetables and fruit.

Exploring the potential of different program delivery points for the behavior change and communication (BCC) strategy: Observations at the **Rally Posts** (where growth monitoring, immunization, and health education activities are held) indicated that while the Rally Posts may be a promising entry point for the BCC program, some aspects of program implementation, such as the timing of the education sessions, the communication techniques of staff, and the allocation of time for communication and counseling, could be strengthened to improve their potential for effective communication with participants. The **food distribution points** (where food commodities are distributed) were identified as the least promising delivery point for the BCC intervention because of their crowded, busy, and distracting environment. The **Mothers' Clubs** (group meetings held in the communities and used primarily for discussions on health education topics) were seen to be the best forum for group communication and discussions, and thus a promising "main" venue for the BCC program. However, it would be important to strengthen the content and teaching and communication approaches used to foster more effective learning and behavior change communication.

4.3.2 Development phase

4.3.2.1 Identification of priority programmatic actions for the BCC component


Priority programmatic actions for the BCC component were identified through discussions with WV staff at all levels. These discussions were held through a series of workshops involving decisionmakers and program staff within WV-Haiti, as well as staff from the U.S. Agency for International Development (USAID) Mission in Haiti and other private voluntary organizations (PVOs) working in the area of child nutrition in Haiti. The workshop discussions focused on prioritizing practices to be promoted through the BCC component, as well as on reviewing the design and the technical and operational aspects of the BCC strategy.

The selection of priority actions for the BCC component was facilitated greatly by the use of a decision tool that summarized and organized the formative research results in the form of a matrix (presented in Annex 4.1). The matrix compares current practices in the program areas to best practices, as summarized in the Guiding Principles (PAHO/WHO 2003). The matrix also summarizes results of the formative research regarding facilitating factors and constraints that could influence the ability of program participants to adopt recommended practices. An example of the decision tool is provided in Box 4.1.

Box 4.1 Example from decision matrix to organize formative research results on exclusive breastfeeding				
Goal	Practices to promote	Practices in Haiti	Facilitators	Constraints
Exclusive breastfeeding (EBF) for 6 months	<ul style="list-style-type: none">- Avoidance of pre- and post-lacteal feeds- Using expressed breast milk when mothers leave home	<ul style="list-style-type: none">- Pre-lacteals and post-lacteal liquids and gruels widely used- Breast milk expression not widely practiced	<ul style="list-style-type: none">- Concept of EBF already introduced- Positive role models exist- No cultural barriers to use expressed breast milk	<ul style="list-style-type: none">- Gruels given when mothers leave home- Lack of training and support on breast milk expression, use and conservation of expressed breast milk

As a next step to the results matrix, we developed a “program-planning matrix” that examined the programmatic actions that would be necessary to address each specific constraint or facilitating factor (see Annex 4.2). The consideration of feasible programmatic actions (presented in the second column) was based primarily on the existing WV program infrastructure and capacity (human, financial, technical). However, future needs and other supporting programs (particularly to support the BCC component) were also considered and these are presented in the third column of Annex 4.2. The program planning discussions held with WV-Haiti focused on identifying modifiable behaviors, constraints, and facilitating factors that could be addressed within the current programmatic options available to them. An example of how constraints to exclusive breastfeeding were addressed is presented in Box 4.2.

Box 4.2 Example from program-planning matrix to address constraints to exclusive breastfeeding

Constraints to exclusive breastfeeding	Program options within current structure and delivery system
<i>Gruels given because mothers need to leave home for work or other activities</i> <i>Use of expressed breast milk is rare; milk expression unknown in some areas</i>	 BCC program: ensuring adequate training in the use and appropriate storage of expressed breast milk

4.3.2.2 Development of the BCC strategy

Following the formative research process and the discussion of the results with WV-Haiti, the BCC strategy was developed. This was done using a “BCC strategy planning matrix,” which outlines the various aspects that need to be addressed in order to ensure that the behavioral change objectives defined through the program planning discussions are achieved. The matrix is presented in Annex 4.3. It identifies, for each age-specific set of practices to promote, (1) *who* needs to be targeted in order to ensure that the desired feeding practice is achieved, (2) *when* the communication related to a specific practice has to reach the identified audience in order to maximize its effectiveness, (3) *where* the communication has to take place in order to reach the desired audience at the right time, (4) *how* specific practices should be promoted at the different program venues and for different program audiences, and (5) *what* is needed to ensure that the communication strategies identified for each type of practice, program venue, and participant are implemented appropriately. An example of the matrix regarding expression of breast milk is presented in Box 4.3.

Box 4.3 Example of BCC strategy planning matrix on expression of breast milk

Practice to promote: Use expressed breast milk as needed (avoid other liquids and foods)

Who will messages be targeted to?	Lactating mothers/fathers/grandmothers
When will messages be delivered?	First 1-2 months of lactation
Where will communication be delivered?	Postnatal consultations and Mothers’ Clubs
How will communication be delivered?	<ul style="list-style-type: none"> - Individual counseling at postnatal consultations - Group discussions and problem solving related to expression of breast milk at Mothers’ Clubs (including demonstration and practice)
What is needed to help with communication?	<ul style="list-style-type: none"> - Training of health staff in communication methods and content of practices to encourage - Provision of resource materials for

4.3.2.3 Development of BCC materials and training plans

Following the identification of the BCC strategies to be used at the different program venues, program communication materials were developed for use in the BCC component. Since the Mothers’ Clubs were identified as the most promising main venue for the BCC, the

material development process focused on materials to be used at the Mothers' Clubs. Further, WV was already in the process of developing other simple materials for use at the Rally Posts.

The materials developed for use in the Mothers' Clubs focus mainly on infant and young child-feeding practices. These practices were considered the most important to address in a program whose goal was to *prevent* malnutrition among children 0-24 months old. Other WV materials were available that covered general aspects of health care and care during illness for infants and children. Details about the communication materials have been reported previously (Loechl et al. 2003b).

The development of the BCC materials consisted of five steps:

- A) *A second review of program communication materials used in Haiti:* In conjunction with the formative research process, a review of two additional sets of BCC materials related to infant feeding in Haiti was conducted to identify newly developed local materials that could potentially be adapted for use in the WV program. The two modules on breastfeeding and young child feeding developed by Freedom from Hunger (FFH) in 2001 and used in conjunction with the FFH Credit for Education program in Haiti were identified as the most appropriate for adaptation and permission was obtained to use them. They addressed breastfeeding practices as well as complementary feeding practices, and included messages related to feeding frequency, responsive feeding, portion size, psychosocial care, and good hygiene practices during food handling. Each module consisted of several learning sessions. These learning sessions were highly detailed and comprehensive. Each session included a set of explicit instructions to the fieldworker, accompanied by activities for them to carry out with the group of participants in order to achieve the objectives of the learning session. The materials were intended for use with a communication strategy that incorporated the principles of adult learning as well as of trials of improved practices.
- B) *Pretesting and adaptation of newly developed messages:* A first step in the adaptation of the FFH materials was the pretesting of new messages developed based on the formative research. The pretest process consisted of four individual and two focus group discussions in the areas where the BCC component was to be implemented. For each item, the interviews gathered information on participant comprehension, the believability of the message, the perceived importance and benefit of the actions implied in the message, and whether the participants would consider changing their practice after hearing the message. After modification of the messages according to the results of the pretest, they were incorporated into the communication materials. An example of the adaptation of the message on expression of breast milk is presented in Box 4.4.

Box 4.4 Example of message adaptation for expression of breast milk

New message tested	Modified message included in communication materials
If the mother has to leave home, she can express breast milk so that another person can give this to the child with a little spoon when she is away.	If the mother has to leave home, she can express breast milk <i>in a cup</i> so that another person can give this to the child with a little spoon when she is away. [To avoid that they express it in a bottle]

C) *Adaptation of BCC sessions based on the formative research and WV program context:* Based on the results of the pretest exercise and the current infant feeding guidelines, the content of FFH modules on breastfeeding and young child feeding was adapted in collaboration with a local firm in Haiti. The adaptation also took into account findings from the formative research. The materials were also adapted to the programmatic context of WV as this differed considerably from the context of the Credit for Education program that FFH had used them in. In the preventive program the order of the topics was changed to be age-specific and to take into consideration the notion that critical pieces of information should reach mothers at what is likely to be the most appropriate learning moment for each set of practice. In addition, the length of the learning sessions was extended, while still maintaining the same structure as with the FFH learning sessions. An example of a learning session that includes instructions on the expression of breast milk is presented in Box 4.5.

Box 4.5 Example of learning session on expression of breast milk

Title: Always promote breastfeeding

Overall objective: To analyze constraints to exclusive breastfeeding and identify solutions

Activity	Objective	Methods used to achieve objective
1	To share experiences with exclusive breastfeeding and find solutions to problems encountered	<ul style="list-style-type: none"> ➤ Paired discussion among participants about breastfeeding experience ➤ Question/answer session on recommendation of exclusive breastfeeding for first 6 months ➤ Review of proper child position during breastfeeding, feeding frequency, and care for breasts and nipples by participants
2	To learn how to express breast milk and how to conserve it	<ul style="list-style-type: none"> ➤ Demonstration of expression of breast milk by experienced mother ➤ Discussions of how to conserve expressed breast milk and how to give it to the child
3	To drink water while breastfeeding to avoid fatigue	<ul style="list-style-type: none"> ➤ Story about drinking water while breastfeeding and further explanations by health promoter

D) *Testing and adaptation of visual aids for BCC:* The instructions for conducting a learning session are accompanied by visual materials. For several of the learning sessions, a large-format, laminated chart on child growth, development, and feeding is

used to facilitate discussion of infant and child-feeding recommendations in relation to the physical development of a child. In addition, a series of images is used to support verbal presentations of the health staff. The images present scenes to illustrate stories and specific feeding recommendations.

These visual materials developed by FFH were adapted in collaboration with a local firm to ensure that the technical information was up-to-date and relevant, and also to ensure that the materials would be culturally relevant and accepted. For example, visual instructions showing manual breast milk extraction techniques were adapted for use in the Mothers' Clubs.

E) *Adaptation of BCC training guides:* The FFH training materials included manuals and resource materials for training of trainers as well as for training of field staff. For the WV staff, the trainers' guide and toolkit for the modules on breastfeeding and young child feeding were adapted to reflect the changes in the content of the learning sessions. In addition, the schedule of learning sessions was created specifically to address the needs of the WV program. The manual on adult learning principles and practices, which is used along with the training materials on breastfeeding and young child feeding, needed only slight adaptations in terminology.

4.3.2.4 *Training of WV-Haiti staff*

The formative research suggested that although the WV health promoters and assistant health promoters were highly motivated to transfer skills and knowledge related to child health to the participants in the Mothers' Clubs, they were constrained by a lack of training in appropriate methods of adult education. Therefore, WV program staff was trained in the use of these communication methods that are grounded in principles of adult learning. Using this approach to teaching and learning, program staff learned how to create a learning environment where people feel safe and respected, how to facilitate group discussions, offer open-ended questions, create dialogue, animate role plays, and build on the ideas of the participants. In addition, the staff was trained in the technical issues related to breastfeeding and young child feeding.

The training was done in two steps. First, the supervisory-level staff was trained in a "training of trainers" session, followed by the training of field staff in WV. Details of the training are described elsewhere (Loechl et al. 2003b).

Training of trainers was done in two stages. In the first stage, all MCHN staff above the level of health promoters and assistant health promoters (i.e., the MCHN National Coordinator, Regional Coordinators, and field supervisory staff) was trained in the use of adult learning principles for communication. In the second step, the same staff was trained over a period of six days on the use of the adapted communication materials on breastfeeding and young child feeding.

Training of field staff (i.e., health promoters and assistant health promoters) was conducted through a six-day workshop. The training was conducted by a group of five WV supervisors who had previously been trained in the Training for Trainers workshops. The health promoters and assistant health promoters were trained in the use of the breastfeeding and young

child learning sessions, and the use of the technical content of the sessions was linked to the principles and practices of adult education.

4.3.2.5 *Development of the implementation plan for BCC intervention*

The implementation plan for the BCC activities at different program points was developed and finalized through a round of discussions held with WV program staff. A first outline of the implementation plan was developed together with the national MCHN coordinator for WV. This plan was further complemented through several meetings with the regional MCHN coordinator and the field supervisory staff in Hinche.

4.4 The Implementation Plan for the Preventive and Recuperative Program Approaches

The WV MCHN program (both preventive and recuperative) offers services at five major points of contact between program staff and participants. These are described in Chapter 3. Following the development of the preventive approach and of the BCC strategy, changes were made to only three service delivery points: (1) the Rally Posts (RPs); (2) the Mothers' Clubs (MCs); and (3) the Food Distribution Points (FDPs). Table 4.1 presents a summary of the activities conducted at these three service delivery points and compares the intervention packages offered in the preventive and recuperative approaches at these delivery points. Shaded areas highlight where the two program approaches differ. Services offered at the pre- and postnatal consultations and at home visits are not included in the table because they are described in Chapter 3 and are exactly the same for both program approaches.

4.4.1 *At Rally Posts (RPs)*

The services to be provided at the RPs are identical for the recuperative and preventive programs, with the only difference being the criteria for selection of food assistance beneficiaries. In the recuperative program group, children are selected based on their nutritional status, whereas in the preventive program group, the selection of beneficiaries is based on age.

In the *recuperative* program, 6-59-month-old children are eligible for food assistance if they are malnourished (M2 and M3¹²) based on the weighing results from growth monitoring activities. There is no defined upper age limit for admittance, although there is an upper age limit for eligibility for program services. For instance, a child can be admitted at 58 months, but s/he would have to exit the program one month later when s/he reaches the age of 59 months.

In the *preventive* program, children are eligible for food assistance based on their age: *all* children between 6 and 18 months of age are eligible to enter and remain in the program until they reach 24 months of age. Thus, in this approach, the upper age limit for admitting children into the preventive program is 18 months, to ensure that all children in the program receive food assistance and other services for at least six months (up to 24 months of age). In the preventive

¹² Malnourished children are defined as M2 and M3 according to the Gomez classification. In this classification, normal (N) corresponds to $\geq 90\%$ of the median of the weight-for-age CDC/NCHS/WHO standards; mild malnutrition (Grade M1) to 75% to $< 90\%$; moderate malnutrition (Grade M2) to 60% to $< 75\%$; and severe malnutrition (Grade M3) to $\leq 60\%$ (Cogill 2003).

Table 4.1 Package of interventions provided in the recuperative and preventive approaches

Recuperative Approach	Preventive Approach
At the Rally Post:	At the Rally Post:
<i>Preventive health and nutrition activities:</i> <ul style="list-style-type: none"> - Vitamin A supplementation (6-59 mo old children and women in first mo. postpartum) - Immunization - Anti-helminth (12-59 mo children) - Family planning - Iron supplementation (pregnant and lactating women) 	<i>Preventive health and nutrition activities:</i> <ul style="list-style-type: none"> - Vitamin A supplementation (6-59 mo old children and women in first mo. postpartum) - Immunization - Anti-helminth (12-59 mo children) - Family planning - Iron supplementation (pregnant and lactating women) - Identification of children 6-23 mo – admission in food distribution program
<i>Growth monitoring:</i> <ul style="list-style-type: none"> - Weighing - Screening of malnourished (M2,M3; 6-59 mo) IF MALNOURISHED (and 6-59 mo) - Brief individual counseling - Admission in food distribution program 	<i>Growth monitoring:</i> <ul style="list-style-type: none"> - Weighing - Screening of malnourished (M3) (24-59 mo) IF MALNOURISHED (and 24-59 mo) - Brief individual counseling - Admission in food distribution program
<i>Record keeping:</i> health promoters maintain records of nutrition/health status of all children measured	<i>Record keeping:</i> same as recuperative
At the Mothers' Clubs	At the Mothers' Clubs
<i>Behavior change communication sessions with small groups of participants, organized:</i> <ul style="list-style-type: none"> - For <i>pregnant and breastfeeding (BF)</i> women: specific to stage of pregnancy & lactation - For <i>mothers of malnourished children:</i> fixed set of sessions Maximum continuous attendance of mothers at the MCs: 21 months	<i>Behavior change communication sessions with small groups of participants organized:</i> <ul style="list-style-type: none"> - For <i>pregnant and lactating</i> women: same as recuperative - For <i>mothers of 6-23 months</i> old children: age-specific set of sessions Maximum continuous attendance of mothers at the MCs: 30 months
<i>Topic of BCC sessions</i> <ul style="list-style-type: none"> - <i>Pregnant women:</i> Diet, dangerous signs, preparation of child delivery, BF - <i>Lactating women:</i> BF, complementary feeding (CF), child development - <i>Malnourished child (6-59 mo):</i> causes of malnutrition and recuperation of malnourished children, CF, child development, BF, immunization, diarrhea and hygiene in food preparation, handling and storage; other topics: HIV/AIDS, family planning 	<i>Topics of BCC sessions:</i> <ul style="list-style-type: none"> - <i>Pregnant women:</i> Same as recuperative - <i>Lactating women:</i> Same as recuperative - <i>Malnourished child (24-59 mo):</i> Same as malnourished child from recuperative group - <i>Child 6-23 months:</i> CF, child development, hygiene in food preparation, handling and storage, causes of malnutrition; other topics: diarrhea, immunization, hygiene, use of <i>moringa olifeira</i>, HIV/AIDS, family planning, home gardening, vitamin A, parasites, water treatment, and ARI infections
	(continued)

Recuperative Approach	Preventive Approach
At the Food Distribution Post	At the Food Distribution Post
<i>Verification of eligibility (malnutrition)</i>	<i>Verification of eligibility (age)</i>
<i>Food distribution:</i> MALNOURISHED CHILDREN (M2, M3; 6-59 MO): for up to 9 months duration Monthly ration: 1 individual (direct) and 1 family (indirect)	<i>Food distribution</i> MALNOURISHED CHILDREN (M3; 24-59 MO): for up to 9 months duration ALL CHILDREN 6-23 MONTHS: up to the age of 23 months (maximum 18 months in program) Monthly ration: 1 individual (direct) and 1 family (indirect)

Note: Shaded areas correspond to aspects that were different between the two program approaches.

approach, severely malnourished children between 24 and 59 months of age are also eligible to participate in the preventive program. As in the recuperative approach, these children (classified as M3 according to the Gomez classification) are identified through the regular growth monitoring activities done at the RPs. The services provided for the severely malnourished children in preventive program communities are also similar to those received by children in the recuperative approach. They include (1) distribution of food rations for 9 months, (2) two meetings for the mothers where issues related to malnutrition and recuperation are discussed, and (3) home visits by health staff during the first weeks after identification.

In both the preventive and the recuperative programs, pregnant and lactating mothers (until their infant reaches 6 months of age) are also eligible for food assistance.

For mothers of children 6-23 months old in the preventive approach and mothers of malnourished children in the recuperative approach, monthly attendance at the RPs and at MCs is mandatory to be eligible to receive the monthly food rations offered by the program. Pregnant and lactating women are also required to participate in MCs and pre- and postnatal consultations to be eligible for the monthly food rations.

4.4.2 At the Mothers' Clubs (MCs)

The formative research identified the MCs as the ideal setting for effective BCC activities; the reasons being that MCs are usually located close to the mothers' homes and include only a small group of mothers, resulting in minimal distractions (especially compared to the RPs). Thus, WV decided to use the MCs as the primary venue for its BCC strategy.

Organization of MCs and timing of sessions: Results of the formative research emphasized the need to reorganize the MCs to include peer groups of mothers of a particular physiological state and/or child age. Based on the BCC implementation plan, separate MCs were organized for pregnant mothers, lactating mothers, mothers of children 6-23 month old (in preventive areas), and mothers of malnourished children (in recuperative areas). For the pregnant and lactating MCs from both the preventive and the recuperative groups, and for the MCs with mothers of 6-23-month-old children in the preventive group, the schedule of sessions for the MCs was planned to be specific to the physiological status of mothers (pregnant/lactating)

and to the age of the child (6-23 months), and to address practices that are immediately relevant for the child's health, development, and growth at a particular age.

Duration of participation: The duration of participation by mothers at the MCs (and other BCC activities) also differs between the two program groups; mothers in the preventive program may benefit from the BCC activities for longer than mothers in the recuperative program. For example, a mother in the preventive group should start attending the MCs during pregnancy for a period of 6 months, continue to attend the same club throughout her first 6 months of lactation and subsequently, for another 18 months, when the child is between 6 and 23 months of age. Thus, mothers in the preventive approach may attend the MCs for up to 30 monthly sessions, without interruption.

In the recuperative approach, this continuity of participation is not the norm. In order to be eligible for the MCs, mothers from this group must be either pregnant or lactating, and/or have a malnourished child 6-59 months of age. Thus, the length of participation of mothers in the MCs depends on their physiological status and on their child's nutritional status. Mothers are eligible to attend the MCs for 6 months during pregnancy, 6 months during lactation (the first 6 months postpartum), and up to 9 additional months if they have a malnourished child at any point in time between 6 and 59 months of age.¹³ Thus, the maximum possible non-interrupted time mothers in this program group can attend the MCs is 21 months, and this will happen only if they start attending during pregnancy, continue through lactation, and have a malnourished child when they reach 6 months postpartum (the malnourished child can be the 6-month-old one or an older child, as long as s/he is less than 60 months of age).

Content of MC learning sessions: The modules on breastfeeding and young child feeding were adapted from materials developed by FFH (see previous section). The modules are organized into learning sessions, which are designed to be completed in about one hour. Each of these sessions covers specific key practices using a variety of communication methods. All the sessions consist of a set of instructions to the health promoters/assistant health promoters, accompanied by activities for them to carry out with the group of participants. Other communication materials are used for topics other than infant and child feeding, such as an album of images with key messages related to other aspects of maternal and child health.

The MC learning sessions covered a range of topics and skills related to infant and young child feeding (Box 4.6). The material covered in the learning sessions closely followed current international "Guiding Principles" for infant and young child feeding (PAHO/WHO 2003; WHO 2005). They were also adapted to address key local constraints to following the Guiding Principles.

Families who participate in WV programs are encouraged to care for their children by accessing RP preventive health-care services such as immunizations, by seeking help from the health agent or other medical professional when their child is sick, and by using the ORS that is distributed at RPs. These caregiving and care-seeking practices are promoted within the larger MCHN program, and thus in both preventive and recuperative areas.

¹³ Children are eligible to re-enter the recuperative program if they are still undernourished one year after having exited the program.

Box 4.6 Mothers' Club learning sessions

Prenatal learning sessions (last trimester)

1. The importance of good breastfeeding practices
2. Good breastfeeding practices

Postpartum learning sessions (0-6 mo)

3. Always promote breastfeeding
4. How to breastfeed better
5. Lactational amenorrhea method
6. Start giving other rich foods to complement breast milk when children are 6 months old
7. Learning to eat: how to breastfeed and feed children less than 12 months old
8. Preparing nutritious foods for children

Sessions for mothers of older children

(6-23 months in preventive areas; malnourished children 6-59 months in recuperative areas)

9. Helping children eat well in health and in sickness
10. Variety of food combinations appropriate for children 6-12 months
11. Protecting your food – protecting your children
12. Feeding children beyond 12 months of age
13. What we can do to combat malnutrition

Timing and schedule of MC sessions: The sessions were designed to be timely and to teach women about key practices before the practices should be initiated. For example, two sessions on breastfeeding were meant to be covered in prenatal MCs, during the last trimester of pregnancy. Similarly, information on appropriate complementary feeding was meant to be introduced in MCs when the child was 4-6 mo of age. At around 6 mo, an important session allowed women to practice preparation of recommended gruels/recipes (Session 8 in Box); this session was covered twice in preventive areas.

The first 8 of the 13 learning sessions were used identically in MCs in both program areas. The last five sessions also covered similar material, but in recuperative program areas, mothers joining MCs had children of varying ages and topics were covered in a different order, and to meet the needs related to feeding somewhat older children. The full schedule of learning sessions at the MCs is presented in Annex 4.4 (A through C). For the MCs with mothers of malnourished children in the recuperative group, the schedule consists of a fixed set of nine learning sessions, which do not take the age of the child into consideration because the purpose is to discuss the recuperation of malnourished children, irrespective of their age (Annex 4.4 C).

Facilitation of learning sessions: The learning sessions were facilitated by health staff trained in adult education principles and techniques, as well as in the content of the learning sessions. Throughout the learning sessions, a variety of interactive techniques were used, including demonstrations and role-playing, and discussion and practice were encouraged. Recommended practices were discussed and local practices compared to recommendations.

Constraints to following recommendations were identified and discussed. Chapter 5 discusses some of the key elements of the facilitation when the operations at the MCs are described.

4.4.3 Food Distribution Points (FDP)

Activities at the FDPs are identical for the two program groups and were intended to be implemented as described in Chapter 3. The only difference is the duration of participation, which is a maximum of 9 months for children in the recuperative communities and 18 months for children in the preventive communities.

4.5 Conclusions

This chapter presented an overview of the process used to develop the preventive and recuperative program approaches, with a focus on the BCC strategy. In the next chapter we examine the quality and fidelity of implementation of these approaches, which was assessed through two rounds of operations research studies. Notably, we discuss how the operations research results were used consultatively with WV-Haiti to improve program implementation. The next chapter thus follows the same thread initiated in this one, and illustrates how action-oriented and program-linked research can be used to improve the design and implementation of programs.

5. DELIVERING IT WELL: THE OPERATIONS RESEARCH PROCESS¹⁴

5.1 Introduction

This chapter discusses the operations research (OR) process used in the present study, which was outlined in Chapter 2. In this evaluation, the OR process was used first to assess the quality of implementation; the information generated by the research was then used to strengthen program processes and implementation. OR was also used to assess differences in program implementation between the two program approaches, and by doing so, to identify critical implementation issues (if any) that might lead to a differential impact of the two program approaches on child outcomes.

In presenting the methods and outcomes of the OR process, this chapter demonstrates how program-oriented research can be used to benefit program implementation and impact evaluations. It also highlights the successful consultation process used in this study, which ensured that research findings were effectively communicated to program implementers and was used for action. The OR process used in this study thus illustrates how the translation of knowledge into effective action is enhanced by joint collaboration and good communication between researchers and program implementers.

5.1.1 Integration of operations research into implementation and evaluation processes

OR typically focuses on program processes and implementation issues, and is often referred to as “process evaluation.” In this chapter, we use the term “operations research” as synonymous to process evaluation since we focus primarily on the process of program *delivery*.


While the usefulness of OR for program improvement is well-established in the family planning and reproductive health literature, there are relatively few documented examples from nutrition programs in the published literature. There are even fewer documented examples of the integration of OR with impact evaluation. Yet, in addition to facilitating program improvement, OR has the potential to enhance impact evaluations in several ways. Information about processes can clarify the pathways through which impact occurs (see, e.g., Robert et al. 2006) and strengthen plausibility when evaluators attribute impact to an intervention. When programs fail to meet objectives, OR results describing implementation problems can provide insights into programmatic factors that could have constrained success.

OR can also play a role in studies such as this evaluation in Haiti, where program approaches are being compared. In this case, comparing the efficiency of implementation of the two approaches is important to help interpret the impact evaluation results. If, for instance, one program proves to have a greater impact on growth than the other program, it is important to be able to rule out the possibility that the greatest impact may in fact be due to better implementation, rather than to the truly greater effectiveness of one approach compared to the other. Thus, when comparing different program approaches, OR should be an integral component of impact evaluation in order to provide a “fair test” of concept.

¹⁴ Mary Arimond led the writing for this chapter.

In this project, OR was conceived from the outset as an integral part of the process, with multiple purposes for its use. Figure 5.1 provides a schematic overview of the different steps involved in overall program development and evaluation, and situates two rounds of OR in this scheme.

Figure 5.1 An “engaged” model of evaluation and program development



Detailed results from both rounds of OR have been previously reported (Loechl et al. 2004; Menon et al. 2005); this chapter provides a summary of objectives, methods, and main results from this process.

5.2 Objectives

The broad objectives of the OR were to (1) assess program implementation and quality of service delivery; (2) use the information to strengthen the quality of operations and ensure fidelity to program approaches and implementation plans; and (3) gain sufficient knowledge about implementation issues and quality of service delivery in the two program approaches to allow appropriate interpretation of impact evaluation results. Two rounds of OR were undertaken, with the first occurring in July-September 2003, shortly after full implementation of the BCC strategy, and the second approximately one year later, in June-August 2004. The first round (2003) focused on the first broad objective and more specifically aimed to:

- A) Assess the fidelity of implementation of the program relative to plans at all service delivery points;
- B) Assess the quality of delivery of the various services; and
- C) Explore perceptions of stakeholders (i.e., participants/beneficiaries¹⁵ and field implementers) regarding the operations and quality of services provided by the program.

This first round of OR (OR1) was followed by a process of consultation with program implementers, including management and field staff. The consultative process (described in Sections 5.3.3 and 5.4.2) was directed toward the second broad objective: through this process, program operations were improved and better aligned with program design.

After allowing approximately 5 months for staff to implement corrective actions/program improvements, a second round of OR was undertaken, with these specific objectives:

- A) To assess implementation of the corrective actions/program improvements identified during the consultative process;
- B) To document differences between the two program approaches in
 - 1) Program implementation at the Rally Posts and Mothers' Clubs;
 - 2) Staff -level factors that could lead to differential implementation between the two approaches (e.g., job satisfaction; motivation; perceptions of supervision, etc.);
 - 3) Beneficiary mothers' knowledge and experiences with trial and adoption of specific infant and young child-feeding and care practices promoted in the BCC,

¹⁵ "Participants" include those who access any of the available range of services; "beneficiaries" are those who live in households receiving food assistance.

in order to gain understanding of constraints to adoption of these practices in the two program areas.

The second round of OR (OR2) allowed continued documentation of program operations and quality of implementation, and allowed us to document improvements in operations. This round of OR also allowed us to meet the third broad objective of OR in this study, which was to enrich the impact evaluation through identifying any relevant differences between program areas in implementation, staff-level factors, or participants'/beneficiaries' experiences.

In the following sections, we define the program context and the service delivery points that were studied in the OR, as well as the stakeholder groups whose views were sought. This is followed by a section describing the research methods used in both rounds of OR, the main results from each round of OR, and a description of the consultative processes that were used to discuss the results with program staff.

5.3 Methods

5.3.1 Program context and stakeholders

A first step in any OR process is to map out the program context and relevant stakeholders so as to ensure that the OR process is comprehensive in its reach. The program context for the OR includes the processes involved in the delivery of services, the norms established by the program for implementation and service delivery, and the main venues for the delivery of services in the WV program. In the OR, we focused on three of the five service delivery points used by the program (see Chapter 3 for description): the Rally Posts (RPs), the Mothers' Clubs (MCs), and the Food Distribution Points (FDPs). The additional two program delivery points - prenatal and postnatal consultations, and home visits - were not included in the OR due to financial and logistical constraints, but also because the services provided at those venues were identical for the two program groups.

The stakeholder groups from whom we sought views in the OR were participants in program activities, beneficiaries of the food assistance component of the program, and the WV administrative and field staff at different levels. WV staff included the health promoters, assistant health promoters (community program volunteers) and food monitors, who are the direct program implementers, their supervisors (the MCHN supervisors and the commodity supervisors), the regional commodity officer, and the regional and national health coordinators.

5.3.2 Data collection methods

A variety of qualitative and quantitative methods were used in the OR process. They include structured observations at service delivery points, structured exit interviews with participants and beneficiaries at selected service delivery points, semi-structured interviews with participants and beneficiaries in their homes, and with health staff at program offices, and focus group discussions with various groups of WV staff. A summary of the research methods used in OR1 and OR2 is presented in Table 5.1.

Table 5.1 Data collection methods used in the OR process

	OR1	OR2
Research Method	Where (with whom)	Where (with whom)
Structured observations	<ul style="list-style-type: none"> - RPs (n = 19) - MCs (n = 20) - FDPs (n = 10) 	<ul style="list-style-type: none"> - RPs (n = 20) Growth monitoring and promotion at RPs: 5 children/RP (n = 100) - MCs (n = 20)
Structured exit interviews	<ul style="list-style-type: none"> - RP (participants/ beneficiaries): 2-3/RP (n = 59)^a - MC (beneficiaries): 2-3/MC (n = 41) - FDP (beneficiaries): 4-5/FD (n = 45) 	<ul style="list-style-type: none"> - RP (beneficiaries only): 2-10/RP (n = 128)^b
Semi-structured interviews	<ul style="list-style-type: none"> - Homes (beneficiaries) (n = 30) 	
Semi-structured interviews	<ul style="list-style-type: none"> - WV offices or other venue with: - Health promoters (n = 20) - Regional health coordinator - National health coordinator - Regional commodity officer 	
Structured interviews		<ul style="list-style-type: none"> - WV offices or other venue with: - Health Promoters (HP) (n = 19) and Assistant Health Promoters (AHP) (n = 19)
Focus group discussions	Various locations with: <ul style="list-style-type: none"> - HP (2 groups) - AHP (2 groups) - Food monitors (1 group) - MCHN supervisors (1 group) - Commodity supervisors (1 group) 	Various locations with: <ul style="list-style-type: none"> - Beneficiary mothers (7 and 5 groups of mothers from preventive and recuperative areas, respectively)

^a In addition to the 59 exit interviews with a random sample of respondents, exit interviews with “tracked” respondents were also conducted (n = 38). These were participants who were identified as they arrived at the RP and were followed through the different activities they went through during their visit at the RP.

^b Mothers were randomly selected, but there were several criteria. Mothers were selected if they had a beneficiary child, and if they had attended at least 5 MCs. Because the interview included an assessment of maternal knowledge, and because we wanted a “fair” comparison between areas, we chose to include only women with this level of exposure. Else, a random selection would have been likely to include women with higher exposure in the preventive program areas, and could have included many respondents with no exposure in the recuperative areas.

The service delivery points observed during OR1 consisted of a subset of all RPs and MCs. As previously described, the overall evaluation included 20 clusters, with one health agent responsible for program services in each cluster. In OR1 we aimed to observe one RP and one MC in each agent’s area of coverage; we achieved 19, rather than 20 RPs and 20 MCs. Note that at that time, each health agent was responsible for between 2 and 5 RPs and up to 12 MCs. Therefore our sample represents a small subset of these. FDPs cover larger geographic areas, so it was possible to observe all FDPs (n = 10) in the evaluation area. For the exit interviews with mothers at RPs, MCs, and FDPs, selection was done randomly; the target and achieved sample sizes were 2-3 per RP for general exit interviews (achieved = 59); 2 per RP for tracked respondents (achieved = 38); 2-3 per MC (achieved = 41); and 4-5 per FDP (achieved = 45).

OR2 gathered data at RPs and MCs as in OR1, but not at the FDPs. The same approach was used in both rounds to select 1 RP and 1 MC per health agent, but the two rounds did not necessarily include the same RPs and MCs. This was due to timing and logistical constraints,

which required the selection of program distribution points that held sessions during the period of the fieldwork and could be visited by the research team.

A similar mix of methods as in OR1 was used, but with some differences in target groups and venues. Differences between the two rounds of OR were generally dictated by differences in the specific objectives of each round; e.g., in OR2 it was necessary to employ more open-ended methods (group discussions) with beneficiary mothers in order to understand their experiences and the constraints they encountered with trial and adoption of recommended practices. Conversely, staff interview methods in OR2 were more structured; structured interview instruments were developed based on themes emerging from staff focus groups in OR1.

More detailed descriptions of methods are available in Loechl et al. 2004 (OR1) and Menon et al. 2005 (OR2).

5.3.3 Consultative workshop

Findings from OR1 were presented by the IFPRI-CU team in Haiti in February 2004. A general presentation of the results to PVOs and USAID Mission staff was followed by a 1.5 day consultation of the IFPRI-CU team and the WV-Haiti management staff. Participants from WV-Haiti included field-level supervisors as well as senior management. The main objectives of the consultation were

- 1) To review and discuss the findings of the OR;
- 2) To discuss the constraints to implementation that were identified in our assessment and to identify potential solutions to address these constraints and to strengthen program operations and quality of service delivery; and
- 3) To prioritize the constraints to be addressed and the potential solutions to be implemented, and to develop an “action plan” for implementing the selected actions.

A matrix was used to guide the discussions and to facilitate prioritization of the constraints to be addressed by the program in the short term (see the example in Box 5.1, and the full matrix in Annex 5.1). At the outset of the workshop, the matrix listed the operational constraints identified at the different service delivery points. During the workshop, the group filled in a column listing potential solutions and one summarizing discussions of the feasibility of adopting the proposed solution(s) in the short term. The group also assessed the potential impact of addressing each problem or constraint, both on program operations and also on the overall impact of the program on its targeted beneficiaries. Summary judgments on potential impact were also recorded on the matrix, in the final column.

Toward the end of the workshop, discussion centered on prioritizing actions to be taken, using three criteria: (1) the possibility of identifying a solution (corrective measure) that was within the scope of current program activities; (2) the feasibility of implementing these corrective measures, given the program’s current financial and human resources; and

Box 5.1 Example of prioritization matrix used at the consultative workshop

Constraint(s) identified	Potential corrective action(s)	Feasibility of implementing corrective action(s)	Potential impact of improving this aspect on program effectiveness and impact
Children are weighed, weights are recorded, nutritional status is assessed. But information is not used to communicate with mother and give her tailored individual advice regarding her child's growth and progress.	Retrain personnel to ensure that each mother is: - Informed of child's weight, nutritional status, progress (gained/lost weight) - Given brief message: a) <i>encouragement</i> (if child growing well); b) <i>recommendation</i> to participate in other program activities (e.g., Mothers' Clubs) (if child not growing well); c) <i>scheduled</i> for home visit (if severely malnourished).	Feasible; will require that supervisors motivate staff and monitor that these actions are implemented. Not feasible to require more in-depth individual counseling due to overcrowding of Rally Posts and scarcity of staff time.	Mothers reported in qualitative research that they like to know their child's weight and progress. This information can help raise awareness and motivate mothers to engage more in behavior change and communication (BCC) activities and adopt recommended practices, which in turn can increase impact on child nutrition.

(3) judgments concerning the potential impact of implementing these corrective measures on the overall program's effectiveness and impact. It was considered most important to identify simple and low-cost solutions, which, if implemented, could have a major impact on operations and on effectiveness. These criteria for identifying and prioritizing actions were carefully gauged in the discussions held in Haiti.

5.4 Results

5.4.1 First round of operations research (ORI)

Detailed results have been reported previously (Loechl et al. 2004); key results are presented here in order to illustrate how OR was used for the purposes identified above.

5.4.1.1 Key results from the Rally Posts (RPs)

Structured observations at the RPs gathered information on operations, including the sequencing and flow of activities, the time allocated to different activities, the quality of implementation of the different activities (registration, education sessions, growth monitoring and promotion (GMP) activities, the distribution of vitamins, deworming tablets and oral rehydration salts, and immunization). A qualitative assessment of the general ambiance and quality of the venue for the RP was also done. Exit interviews with beneficiaries assessed access, services used and the perceived importance of the different services offered by the program. Semi-structured interviews and focus groups with health staff gathered information about their perceptions related to the functioning and quality of services at the RPs and their general impressions relative to their current responsibilities, the constraints on their performance, the supervision structure of the program, the coordination between the health and food

distribution (commodity) components of the program, and to gather their suggestions relative to potential program improvements.

Program operations and service delivery at the Rally Posts

OR1 results suggested that RPs were generally operating according to the program's implementation plan, and in general were providing participants with nearly the full range of planned services and activities. Areas where the quality of service delivery appeared to call for improvement were the group education sessions, and the communication between staff and mothers during the growth monitoring and promotion (GMP) activities. More specifically, the following constraints to operation and quality of service delivery were observed:

- ❖ RPs were crowded, with a high participant-to-staff ratios, and bottlenecks (particularly at registration) and long waiting times for mothers;
- ❖ The crowded and noisy conditions also impacted negatively on the quality of group education sessions;
- ❖ Education sessions were generally held only once, near the beginning of the RP, and late arriving mothers could therefore not attend these sessions;
- ❖ Education session topics generally did not follow the plan for the month;
- ❖ Communication between staff and mothers during GMP was limited and messages given to mothers about their child's growth were often vague and inaccurate;
- ❖ Measurement errors when weighing children resulted in the misclassification of a substantial proportion of children into inaccurate categories (Gomez classification); and
- ❖ There were problems with availability of supplies (e.g., vitamin supplement capsules, ORS, deworming medicine).

Staff perceptions of Rally Post services and activities

Staff perceived the RPs to be functioning well in general, and both health promoters and supervisors viewed the RPs as making important contributions to the families they served. Further, they noted that the RPs were of importance because they served the whole community, not just those identified as being eligible for food assistance.

Staff also mentioned many of the same problems and constraints identified by the RP observations, including crowding, high participant-to-staff ratios, and lack of a wide range of supplies and equipment, including chairs and tables as well as scales, cold chain, and medical supplies. In addition, they cited late arrival by participants as a problem as the participants then would miss the (single) education session. Health promoters requested more training for assistant health promoters, so that they could better assist them with their multiple tasks. Staff also identified the lack of transport for themselves as a deeply felt problem, and one which could result in RPs starting late and longer waits for participants.

5.4.1.2 Key results from the Mothers' Clubs (MCs)

The OR1 observations of MCs focused on assessing the quality of teaching and facilitation at the MCs, and also on assessing whether the MCs had been constituted based on child age subgroups, as described in Chapters 3 and 4. Exit interviews with beneficiaries assessed issues related to access to the MCs and their perceptions regarding the importance of the topics discussed at the MCs. Interviews with health staff gathered information on their perceptions and opinions regarding the importance and quality of the services offered at the MCs.

Program operations and service delivery at the Mothers' Clubs

Results pertaining to the quality of delivery of the learning session were very positive, as health staff demonstrated both good mastery of technical content and good use of new facilitation skills. However, the MC observations indicated that the reorganization of MCs in the preventive group was still a work in progress. More specifically, analysis of operations at the MCs revealed the following:

- ❖ Quality of education was high, both in terms of technical content and teaching and communication skills.
- ❖ Organization of MCs based on child age in preventive program communities did not follow the plan; this resulted in many mothers not receiving appropriate learning sessions for their child's age.

Staff and beneficiary perceptions of the Mothers' Clubs

All health staff felt that the MCs were, in general, functioning well, and they expressed a variety of positive impressions in the semi-structured interviews and focus groups. These included that the MCs provided a good forum for an exchange of ideas, and that exposure to MCs did lead to behavior change on the part of the mothers, and that mothers enjoyed the MCs. Staff also felt that the MCs had improved since the development of the new BCC strategy and subsequent staff training.

Some of the problems mentioned with the MCs were logistical in nature, and included late arrivals and absenteeism, and frustrations that women did not always bring ration cards with them. Lack of transport for staff, poor venues, and lack of seating for mothers were also commonly cited; less common were inadequate staff time to prepare for the MCs, and timing/scheduling of MCs. In addition, a number of staff mentioned the challenges in eliciting participation, especially in newly formed MCs.

Beneficiaries were asked for their views of the MCs during the semi-structured interviews in their own homes. Views expressed were generally very positive, with mothers reporting that they felt the MCs were important for their children's health. Most felt that the MCs were the easiest of the MCHN services to access, as the clubs were organized close to women's homes, and the time commitment was not excessive.

5.4.1.3 Key results from the Food Distribution Points (FDPs)

Information gathered about the functioning of the FDPs covered the following domains: overall organization of the FDPs including timing, crowding, and waiting times; quantity of food relative to allocated rations; quality of food; and staff and beneficiary perceptions of FDP operations.

Program operations and service delivery at Food Distribution Points

There are many challenges inherent in commodity food distributions; these include tracking and accounting for food at all points in time, and physically transporting large quantities of commodity foods on bad roads and in all seasons. These inherent challenges require that food be distributed from a centralized point. The result of this, for beneficiaries, can be long travel times to reach the distribution point and overcrowded distribution venues. These realities were reflected in OR1 results for the FDPs:

- ❖ The quality of distributed commodities was generally good, with almost all sampled bags/containers free of infestation or visible changes in color.
- ❖ FDPs were crowded, with high attendance (range 117-375 beneficiaries), high beneficiary-to-staff ratios, and long waiting times (average of 4 hours).
- ❖ FDPs often started late, primarily due to staff transport problems.
- ❖ On average, the amounts of food distributed conformed to the plan for the FDP (kgs and number of beneficiaries), but the averages masked frequent measurement errors in allocating household food rations.

Staff and beneficiary perceptions of Food Distribution Points

Staff interviews and focus groups revealed that health staff and commodities staff shared some common understandings, but that views diverged on several other issues. Both health and commodities staff viewed the delays in starting food distribution to be a major problem, and both attributed delays to transport problems, including bad roads, rains, and equipment breakdowns, as well as availability of transport.

In contrast to this shared perception of transport problems, numerous health staff viewed crowding at the FDPs as a problem, while this was not raised as a point of concern in interviews/focus groups with commodities staff. Similarly, several health staff raised concerns about hygiene practices during food handling at the FDPs, but these concerns were not raised in interviews with commodities staff.

Views also diverged on problems related to confirmation of eligibility for individuals. Individuals could be denied rations for a number of reasons related to record keeping (beneficiary lists, entry and exit dates on a master list and on ration cards, etc.) or in cases where ration cards were forgotten. Health staff tended to view commodities staff as being overly restrictive or even punitive, whereas commodities staff felt constrained to follow strict rules regarding eligibility. Commodities supervisors expressed more flexibility regarding rules, but

this flexibility did not appear to have been communicated to frontline staff. All agreed that better communication as beneficiary lists were prepared could help.

In general, there appeared to be a need for more dialogue and collaboration between these two arms of the program.

As would be expected, beneficiaries expressed great appreciation for the food that they received, and many specifically expressed appreciation for the wheat-soy blend (WSB), because they understood that it contained vitamins that were beneficial for their children. The main problem for beneficiaries was the long-time commitment involved in attending the FDP, which they related both to the distance traveled, and the long waits at FDP. Despite the variability we observed in ration sizes, only two respondents expressed this as a problem.

5.4.2 Outcomes of consultative workshop

The consultative workshop described previously led to a variety of agreements on the corrective actions to be implemented by the program at the different delivery points. A summary of these is provided below and further details are presented in the matrix in Annex 5.1.

5.4.2.1 Solutions for the Rally Posts

Organization

- ❖ Few solutions were identified to solve overcrowding in the short term because resources for hiring more staff were unavailable. Splitting RPs to decrease participant/staff ratios could not be done except in very few cases.
- ❖ Improve training of assistant health promoters (AHPs): The management staff indicated that this measure had already been implemented and that AHPs were now being included in all field staff training. The AHPs had also recently been “promoted” and their salary was raised from 30% of the health promoters’ salary to 50%.¹⁶
- ❖ Revise sequence of activities at the RP; use of tickets to ensure that mothers are being attended in the order in which they arrive.

Group education

- ❖ Increase the coverage of the education sessions by offering more than one education session at the RPs.
- ❖ Keep education sessions short; use sessions to deliver simple messages and reinforce topics taught at the MCs.

¹⁶ These changes were accompanied by changes in job titles, to recognize and formalize the difference. However, for consistency, we refer to the two levels of staff as health promoters and assistant health promoters, respectively, throughout this report.

- ❖ Enforce use of 12-month calendar of education topics to ensure that participants are exposed to the complete set of topics planned by the program (supervisors to take responsibility).

Supplies

- ❖ Program management to pursue efforts to maintain adequate supplies of vitamins, ORS, deworming tablets, and vaccines (through coordination with main providers (Ministry of Health, UNICEF, etc.)).
- ❖ No action recommended for field staff on this issue.

Growth monitoring and promotion (GMP)

- ❖ Retrain and increase supervision of health staff conducting GMP to reduce measurement errors (and consequent misclassification of children).
- ❖ Simplify GMP process, but ensure that at least the following is provided: all mothers should be informed about the child's weight, nutritional status, progress since last weighing; and provided with brief individual advice. The advice should consist of praising and encouraging the mother if the child is growing well; if the child is not growing well, the mother should be advised to follow up with other program activities and adopt recommended practices discussed at the MCs (e.g., exclusive breastfeeding, use of enriched porridges, etc.).

5.4.2.2 Solutions for the Mothers' Clubs

Organization of MCs by child age subgroups

- ❖ Pursue additional efforts to ensure the organization of MCs follows implementation plan, i.e., separate MCs for mothers of children aged 6-11 months and those of children aged 12-23 months.
- ❖ Supervisors to revise rationale for the new organization of MCs with local staff and provide necessary support for implementation.

Quality of teaching and facilitation skills

- ❖ Ensure effective supervision and feedback to maintain a high quality of education, achieve a greater consistency of performance, and maintain staff enthusiasm and motivation, and interest of beneficiaries/participants.

5.4.2.3 Solutions for the Food Distribution Points (FDPs)

Logistical problems

- ❖ Continue ongoing efforts to address constraints identified in OR1.

- ❖ Pursue efforts to improve transport and logistical problems as possible (it was well recognized that several of the transport and related logistical problems could not be fully overcome in the Central Plateau, where road conditions are bad and availability of fuel is a major constraint).

Errors in amounts of food received

- ❖ Strengthen the existing supervision system to reduce errors in the measurement of food rations (e.g., supervisors could systematically verify rations provided to a subsample of beneficiaries).

Communications difficulties between commodity and health staff

- ❖ Strengthen the supervision to ensure good communication among all staff and with the beneficiaries (treat cases of miscommunication individually, as needed).
- ❖ Organize joint preparation (between health and commodity staff) of beneficiary lists, in advance of each FDP.

5.4.2.4 Follow-up steps from the consultative workshop

The February 2004 consultative workshop was a key step in the process of program improvement. All staff present at the meeting participated very actively in the dialogue, and controversies were not avoided, but were skillfully managed by senior staff. The process used to identify and prioritize feasible solutions was effective, and this same process also appeared to ensure among staff a sense of ownership of solutions and responsibility for taking action.

Notably, discussions also revealed that management had already initiated actions to resolve several of the problems identified in OR1, through such activities as joint planning between health and commodities teams. It was noted that the action plan described above would rely heavily on a strong and effective supervision system. At the time of the consultation, the management staff also indicated that they had already been taking steps to strengthen the supervision of the program at all levels, and the health team presented their newly developed supervision plan for the region.

5.4.3 Second round of operations research (OR2)

The second round of operations research (OR2) was carried out in mid-2004, approximately 5 months after the consultative workshop. As noted earlier in this chapter, this round was undertaken to (1) assess implementation of the corrective actions and program improvements identified during the consultative process and (2) to document differences between the program approaches in program implementation at the RPs and MCs, in staff-level factors that could lead to differential implementation, and in beneficiary mothers' knowledge and experiences with trial and adoption of specific infant and young child-feeding and care practices promoted in the BCC. The methods used in OR2 are summarized in Table 5.1 (page 55).

For the second round of OR, the decision was made not to return to FDPs for more observations. A number of the problems identified at the FDPs were deemed not amenable to

change, while others were to be addressed through improvements in communications. Direct observation at FDPs in 2004 was not considered likely to capture changes in communication.

This section presents the results of findings related to program implementation and operation, including improvements since OR1 and differences between program groups at OR2. In considering the differences at OR2, we also present differences between the groups in staff knowledge, motivation, and perceptions of supervision.

5.4.3.1 Key results from the Rally Posts

Organization: Flow of activities, staffing, and crowding

OR2 observations confirmed that the new sequence of activities proposed during the February 2004 consultation had been successfully implemented in most RPs. Staff nearly universally (97%) preferred the new sequence and all agreed that RPs were more “smooth and calm.” However, registration was still identified by observers as the most important bottleneck in almost all RPs. Regarding the crowding and long waits identified in OR1, management agreed to continue to consider splitting RPs when possible, and to consider training more assistant health promoters. Findings from the 10 RPs observed both in 2003 (OR1) and 2004 (OR2)¹⁷ show that while the average number of staff remained the same (4 staff per RP), the variability and number of participant-to-all staff ratios decreased (maximum dropped from 59:1 to 33:1). In sum, it appeared that while overall staffing remained the same, staffing levels had become more consistent for those RPs observed in both years.

Comparisons between program groups in 2004 for the 20 RPs observed (10 in preventive and 10 in recuperative program areas) show that average participation and participation-to-health promoter ratio were higher in preventive compared to recuperative RPs. The age distribution among children attending the RPs was also different between program groups. Among the randomly selected children for whom GMP was observed, 94% were between 6-23 months in the preventive RPs as compared to 50% in the recuperative RPs. Only 2% of the children observed in the preventive RPs were 24 months or older as compared to 46% in recuperative areas. The difference could have had implications for coverage of preventive health services (supplementation, deworming, ORS, and vaccinations) and is assessed in Chapter 7.

Group education sessions

A number of improvements compared to OR1 were observed at OR2 in the group education sessions at RPs. These included an increase in the number of sessions per RP (from average of 1 to 2), a smaller number of women attending each session (average dropped from 30 to 14), a longer duration of the education sessions (from 10 minutes in OR1 to 22 minutes in OR2), and a greater use of visual materials (from 16% in OR1 to 84% in OR2). With very few exceptions, the topics of the education sessions during OR2 followed the WV calendar, which was rare in OR1. No differences between program groups were observed in 2004.

¹⁷ As noted in the methods section of this chapter, due to timing and logistics, it was not possible to observe all the same RPs in both rounds. However, half (10) of the RPs were observed both in 2003 and in 2004; comparisons focusing on the number of participants and the staff/participant ratio focus on these 10 RPs because RPs vary widely in these aspects and thus, comparisons between 2003 and 2004 would be misleading if all 20 RPs observed at each round were included in the comparisons.

Growth monitoring and promotion

Quality of measurement. OR1 observations had revealed errors both in measurement and in plotting of weights. During OR2, observations focused on one aspect of technique that had been covered in the retraining, which was to wait until the needle on the scale had stopped moving before reading the weight. Marked improvements on this aspect were observed between the two rounds, with the proportion of readings of the scale with the needle “swinging a lot” reduced from 21% in OR1 to 1% in 2004. Improvements were seen in both program areas, but in 2004, a somewhat larger proportion of measurements were correctly taken in RPs from the preventive (68%) compared to the recuperative (50%) area.

Quality of communication with mothers. Communication with mothers during GMP was identified as particularly problematic in OR1. Retraining was planned to ensure that mothers were informed and counseled appropriately, as described above in Section 5.4.2. Information on the quality of communication was gathered both through direct observation of GMP activities (n = 100) and through exit interviews with mothers (n = 128). Tables 5.2, 5.3, and 5.4 provide detailed results comparing OR1 and OR2 and comparing preventive to recuperative RPs. Key results suggest that

- ❖ Overall, mothers were much more likely to be given information and brief counseling in 2004 than in 2003;
- ❖ Mothers in preventive area RPs were somewhat more likely to be informed of their child’s weight, nutritional status, and progress (gain/loss) and were more likely to receive brief individual counseling than mothers in the recuperative area;
- ❖ Among those given information on their child’s progress, information was more likely to be accurate in preventive (98%) than in recuperative areas (76%);
- ❖ Two messages selected during the consultation—promoting use of the special gruels, and reinforcing importance of attendance at MCs—were rarely used in OR2 (10% and 2% of observations, respectively), and the most common counseling messages, while not inappropriate, remained vague (“feed more food”).

Availability of supplies

Because of the perceived difficulty of resolving the problem of availability of supplies, and the fact that this problem was largely outside of WV’s control, the participants in the consultative workshop concluded that expectations should be reasonable. It was considered feasible and necessary for WV-Haiti management to be persistent in attempts to maintain supplies, and for management to maintain good communication with collaborating agencies. However, these management efforts were not judged to be likely to completely resolve this issue. OR2 results confirmed this reality, and no differences between the two program groups in these aspects were identified.

Table 5.2 Types of information communicated to mothers during growth monitoring at Rally Posts, by year, by program area, and by research method

Program area	Overall	Overall	Preventive	Recuperative	Preventive	Recuperative
Year	2003	2004	2004			
Method	Observation	Observation	Interview	Observation	Observation	Interview
Number of children	n = 38	N = 100	n = 128	n = 50	n = 50	n = 85
	Percent of children					
Child was weighed	100	100	100	100	100	100
Weight was plotted / recorded on health card	76	100	n/a ^a	100	100	n/a ^a
Nutritional status was assessed	84	100	n/a ^a	100	100	n/a ^a
Mother was told weight	42	89	96	96	82	98
Mother was told nutritional status	24	81	47	96	66	50
Mother was told if child gained or lost weight	n/a ^b	80	62	96	64	64
Mother was given individual message	37	71	55	84	58	55
Mother was told nutritional status if child's weight-for-age is:						
Normal	22	85	46	95	65	47
M1 / "orange" ^c	13	79	50	100	71	63
M2 / "yellow" ^d	40	50	67	n/a	50	n/a
Mother told if child gained or lost weight:						
Normal	n/a ^b	82	62	95	57	64
M1 / "orange"	n/a ^b	82	67	100	76	75
M2 / "yellow"	n/a ^b	50	100	n/a	50	n/a
Mothers given individual message if:						
Normal	22	71	49	81	52	52
M1 / "orange"	33	71	73	100	62	88
M2 / "yellow"	45	67	100	n/a	67	n/a

^a Women were not asked during exit interviews if weight was plotted or if nutritional status was assessed.

^b This information was not recorded by observers in 2003.

^c OR-2004 occurred during a period of transition between two types of health card; one indicated classifications using the Gomez system (M1, etc.) and the other showed different color zones for children falling below cutoffs using the Z-score system.

^d Only six of the children for whom growth monitoring was observed were classified as M2 or "yellow"; all six were in the recuperative program area.

Table 5.3 Types of advice given to mothers at Rally Posts in 2004, by child's progress^{a, b}

	Overall (n = 100)	Gained weight (n = 52)	Lost weight (n = 40)	Same weight (n = 8)
Percent of children				
Mother was praised	27	50	3	0
Reinforced importance of MCs	2	0	5	0
Told to feed enriched gruels	10	6	18	0
Told to feed more food	37	27	48	50
Given other advice	16	12	20	38
No message after weighing	29	29	30	25

^a Data are from direct observation of growth monitoring and promotion.

^b Some women were given more than one type of advice, so percents sum to more than 100.

Table 5.4 Advice given at Rally Posts in 2004, by child's progress and program area^{a, b}

	Gained weight		Lost weight	
	Preventive (n = 24)	Recuperative (n = 28)	Preventive (n = 21)	Recuperative (n = 19)
Percent of children				
Mother was praised	58	43	0	5
Reinforced importance of MCs	0	0	10	0
Told to feed enriched gruels	4	7	29	5
Told to feed more food	38	18	67	26
No message after weighing	13	43	19	42

^a Data are from direct observation of growth monitoring and promotion.

^b Some women were given more than one type of advice, so percents sum to more than 100.

5.4.3.2 Key results from the Mothers' Clubs: Improving organization and maintaining quality of communication

Improving organization of MCs

Clear improvements were observed in 2004 in organizing the MCs according to the two program approaches: in both preventive and recuperative areas, staff had succeeded in grouping women appropriately, according to their physiological status (pregnant, lactating in both program approaches), their child's malnutrition (recuperative approach) and/or their child's age group (preventive approach). In both program areas, the content of observed sessions was also found to be appropriate for maternal status and/or child age or status. MCs were also of appropriate size in 2004 (range, 7-17 women), whereas in 2003, four of the 20 MCs observed were considered too large (more than 20 women).

Maintaining high-quality communication

OR1 had documented a very high quality of communication and facilitation of MC sessions, so maintaining that level of quality was perceived as a potential challenge. Five domains of quality were observed in both years: (1) technical content; (2) session management and organization; (3) teaching and facilitation skills; (4) attitudes displayed (with observer to explicitly note how attitude was demonstrated); and (5) atmosphere at the learning session.

Within each of these domains, the observation tool listed a number of specific and observable practices; these are detailed in Annex 5.2. The results presented in Table 5.5 show that the generally high level of quality noted in 2003 was maintained in 2004. The range of scores observed, however, suggests that there was still notable variability in health workers' performance in OR2. No major differences were observed between program areas.

Table 5.5 Quality of information and facilitation at Mothers' Clubs, by year and by program area^a

	Overall	2004		
	2003 (n = 20)	2004 (n = 20)	Preventive (n = 10)	Recuperative (n = 10)
	Percent or mean (range)			
1. Technical content				
- Percent of key session points presented correctly	83% (64-100%)	84% (42-100%)	82% (42-100%)	86% (64-100%)
- Percent of key points where incorrect information was presented	14% (7-33%)	10% (0-58%)	14% (0-58%)	2% (0-21%)
- Percent of key points not presented during session	9% (0-40%)	6% (0-36%)	4% (0-18%)	8% (0-36%)
2. Session management and organization (score 0 to 6)	5.1 (3-6)	4.8 (3-6)	5.0 (4-6)	4.5 (3-6)
3. Teaching and facilitation skills (score 0-9) ^b	6.5 (2-9)	6.7 (2-9)	7.2 (5-9)	6.1 (2-9)
4. Attitudes displayed (score 0-4)	3.0 (0-4)	2.5 (0-4)	2.7 (1-4)	2.2 (0-4)
5. Atmosphere of learning session (score 0-6)	3.2 (1-5)	3.4 (1-6)	3.4 (1-6)	3.4 (1-6)

^a For each summary score presented, the HP (or assistant) received a score of one for each positive checklist behavior s/he demonstrated. These were then summed to create the score for each domain.

^b There were 11 possible "behaviors" to observe in the domain of teaching and facilitation skills. However, two behaviors relating to use of visual aids were not applicable for several of the observed sessions, and so those two behaviors were not included in the summary score.

5.4.3.3 Summary of improvements and differences in implementation between program areas

Table 5.6 summarizes changes implemented since the February 2004 workshop and identifies the few areas where implementation differed between the two program areas. In the table, the third column indicates "no differences" between the groups if any observed differences are small and considered of no practical significance. In the fourth column, the potential influence of the observed differences is described.

The overall picture presented is one of impressive success in implementing the program improvements that had been identified and prioritized in the workshop. For most changes identified and prioritized at the consultative workshop, significant progress was seen in both preventive and recuperative RPs, with no large differences between the two program groups.

Among the few differences observed between the two program areas, the most notable are related to participation at RPs. As noted in Table 5.6, higher participation in preventive areas could negatively impact the quality of services through crowding and shortage of time for health staff to focus on each participant. We found, however, that quality of measurement in growth

Table 5.6 Summary of program changes in WV-Haiti MCHN program, and differences between preventive and recuperative program areas in 2004

Program element	Change between 2003 and 2004	Differences between preventive and recuperative program areas in 2004	Potential influence of differences on impact
At the Rally Posts (RPs)			
Organization—sequence of activities and bottlenecks	Participants now provided with a number on arrival, per plan; sequence of activities is much more consistent and follows plan, as compared to 2003.	No differences.	N/A
Organization—staffing and participation	No change in size of RPs for those observed in both 2003 and 2004. No change in average number of staff, but range in number of staff at RPs reduced from 1-7 to 4-5; participant-to-staff ratio slightly reduced, and maximum ratio decreased from 59:1 to 33:1.	No difference between program groups in number of staff ; the number of participants and the participant-to-staff ratio were higher in preventive RPs.	No differences in staff costs; higher participant-to-staff ratios could increase waiting time or decrease quality of service in preventive areas.
Age distribution of children	No change between 2003 and 2004.	Greater proportion of younger children (< 2 years) at preventive RPs; many fewer older children (2-5 years) at recuperative RPs.	Could have implications for coverage (e.g., lower immunization coverage for children < 2 years of age in recuperative areas; lower coverage for ORS, Vitamin A and deworming for older children in preventive areas).
Group education sessions	Many positive changes successfully implemented: More education sessions; Longer sessions; Smaller groups of women; Follow monthly plan; More use of visuals.	No differences.	N/A

(continued)

Program element	Change between 2003 and 2004	Differences between preventive and recuperative program areas in 2004	Potential influence of differences on impact
Growth monitoring and promotion			
- Quality of measurement	Much improvement in targeted technique (reading scale when needle is still); other problems persist (incorrect hanging of scales, slight inaccuracy of scales).	More improvement in weighing technique observed in preventive RPs; scales also slightly more accurate relative to standard weight.	Effectiveness of recuperative program could be reduced if children are poorly measured and consequently misclassified as malnourished or well-nourished.
- Quality of communication	Major increases in the proportion of mothers informed of child's weight, nutritional status, and child's progress. Mothers also more likely to receive brief advice. Frequency of delivery of specific recommended messages still low; messages still tend to be vague, although not necessarily incorrect.	Increases were larger in preventive RPs; recommended messages given more frequently; information on progress more likely to be accurate.	Impact of GMP could be higher better in preventive areas if advice triggers action
At the Mothers' Clubs (MCs)			
Organization	Marked improvement in organizing MCs according to plan, and in providing education session appropriate to the group.	No differences.	N/A
Quality of education	On average, high quality of education at MCs maintained since 2003 in five domains: technical content; session management; teaching skills; attitudes, and atmosphere. Quality still varies a lot between health staff.	No differences.	N/A

monitoring was slightly better in preventive compared to recuperative areas, and so was the quality of individual counseling of mothers. Finally, the age distribution of children attending the RPs was quite different, with a greater proportion of younger children, and fewer older children attending preventive area RPs. This could have implications for coverage of various services (vaccination, vitamin A supplementation, ORS distribution, and deworming), a topic that will be discussed in Chapter 8.

In addition to assessing changes and differences in implementation between program areas, we also explored two additional sets of factors that could influence the effectiveness of the program—and had the potential to differ between program areas. These are staff-level factors such as knowledge, motivation, and workload (reported in the next section) and the mothers' knowledge and experiences with trial and adoption of recommended practices (reported in Chapter 8).

5.4.3.4 Staff skills and motivation

Information from staff focus groups (OR1) was used to guide development of structured interview tools, used during OR2. Health promoters and assistant health promoters from 19 of the 20 study clusters were interviewed. Interview questions covered the following domains: job satisfaction; motivation; perceptions of supervision; technical knowledge, and time allocation and workload. Detailed results are reported in Menon et al. 2005; a brief summary is provided here.

Job satisfaction. Results were nearly identical in the two program areas, all staff reported that, overall, they were at least “somewhat satisfied” with their jobs (29%). A similar number (32%) reported being “very satisfied” and the remaining 40% were “satisfied.”

Motivation. Staff were asked 24 questions reflecting various qualities of the work environment and their own perceptions and beliefs about their work, each of which could bear on motivation. For example, staff were asked if they agreed/disagreed that MCs could change behaviors; they themselves received adequate training; program management valued their work; they were satisfied with their salary, etc. (see Annex 5.3 for the full list of questions). The 24 questions were summarized into scales reflecting the following dimensions:

- ❖ Feeling of being valued by management and participants (1 scale);
- ❖ Confidence in themselves and enjoyment of work (1 scale);
- ❖ Adequacy of training, salary, and support from supervisors/management (3 scales);
- ❖ Discouragement and plans to leave (2 scales).

Results for these scales showed nearly identical scores for staff in the two program areas. Staff had many positive perceptions and generally felt valued and well supported, adequately trained, and confident, and they reported enjoying their work and viewing it as important. There was more variability in responses related to salary.

Perceptions of supervision. Staff were also asked a series of 29 questions regarding their perceptions of the quality of supervision that they received. For example, staff were asked how frequently their supervisor consulted them before changing their activities, and how often the supervisor took concerns of frontline staff up to senior management. The full list of questions is provided in Annex 5.4. The responses were summarized into two scales. The first reflected staff perceptions of the adequacy of supervision in terms of tasks; i.e., did the supervisor visit, provide appropriate feedback, ensure adequate supplies, and help organize activities. The second captured whether staff felt supported, valued, and respected by their supervisor.

Results were once again nearly identical between the two program areas. Scores for each of the two scales ranged from 15-75, and mean scores were 58 (supervisor performs expected tasks) and 55 (supportive supervision). The mean scores are toward the higher end of the range in each case but ranges were quite wide, reflecting diversity in levels of satisfaction with supervision.

Technical knowledge. Staff were asked 44 factual questions, covering “core” material in the MC sessions (26 questions) and additional questions covering background material related to nutrition. Results were strongly positive and consistent, and did not differ between program areas. When all 44 questions were summed into a score, the average score was 88% correct (range 79-94%). On the “core” material, scores were even better, with an average score of 96% correct (range 90-100).

Time allocation/workload. Information on time use was gathered from staff and management. Staff were asked to report the number of service delivery points they were responsible for organizing and/or staffing each month (RPs, MCs, and FDPs). In addition, senior management outlined expectations for the number of days each health agent or *colvol* would spend in activities such as reporting, training and meetings with supervisors, community meetings, etc.

When all activities are summed, 15-16 workdays are required to cover service delivery points and all meetings, training, reporting responsibilities, etc. In addition, health promoters must find time for home visits with new mothers and in households with malnourished children.

Despite the higher participation and larger number of MCs in preventive program areas, health promoters in both areas reported very similar workloads. Assistant health promoters in preventive program areas reported working approximately 10 hours per month more than their counterparts in recuperative areas; the largest difference was in time spent on MCs.

In summary, we found no substantial differences in staff-related factors between the two program areas, and certainly no differences that were likely to have significant implications for program implementation. We found that the work context for WV health staff was very positive, with the presence of motivating factors like good training, good relationships among staff, and a feeling of being valued by management and beneficiaries. Staff also perceived supervisory practices to be appropriate and supportive, and staff time allocation to program tasks appeared to be reasonable. We also found few de-motivating factors. Finally, staff appeared to possess both very good technical knowledge and the communication skills needed to ensure the effectiveness of the BCC arm of the program.

5.5 Conclusions

The two rounds of OR described in this chapter allowed us to meet the three objectives outlined earlier, i.e., to assess program implementation and service delivery, to strengthen program implementation, and to gain sufficient knowledge about implementation and quality of service delivery in the two program approaches to allow appropriate interpretation of impact evaluation results. In the process, the operations research studies also provided several opportunities for active engagement and discussion among the external IFPRI-Cornell research team and the internal WV-Haiti implementation team.

The OR process also provided an in-depth understanding of several key steps along the program theory pathway laid out in Chapter 2. More important for the impact evaluation, it showed that there were few, if any, differences between program groups in program delivery. Program services were being delivered largely as planned in both program areas and the quality of delivery of the BCC strategy was high. In addition, the careful comparisons between program groups at different service delivery points as well as staff knowledge, motivation, and perceptions of supervision showed very few differences in these aspects between program areas.

In sum, these results suggest that differences in program impact on child outcomes are highly unlikely to be explained by differences in program delivery. They lead us to infer that any differences found in our primary impact measure—i.e., nutritional status—can reasonably be attributed to the design of the program approaches, and not to differences in implementation.

6. PROGRAM PARTICIPATION AND HOUSEHOLD AND CAREGIVER CHARACTERISTICS

6.1 Introduction

This chapter describes the program participation patterns in the preventive and recuperative program communities. It examines participation among pregnant and lactating women as well as among the children. The chapter also compares the characteristics of households and caregivers who participated in the program with those who never participated in the program.

We examine participation patterns as well as factors associated with participation to enable better interpretation of the program impact results presented in Chapters 7 through 10 and the cost-effectiveness results in Chapter 11. In examining the factors associated with participation, we focus on those factors that are *not* expected to be impacted by any of the program inputs, such as age, maternal education and house quality, and access to services. We do not, on the other hand, examine the association between participation and household assets or food security because the program can be expected to improve both of these outcomes through the provision of monthly food assistance.

6.2 Data and Analysis

6.2.1 Data and variables

The data for this chapter are derived from the household survey described in Chapter 2 (section 2.4.1.1.). We compare maternal and household characteristics between program participants and nonparticipants. The following variables were used to assess household participation in the different components of the program.

Any exposure to the program: Data were gathered on exposure to the program for the index child (12-41 months old), the younger sibling (if the index child had one; 0-11 months old), and any other sibling in the same age range as the index child (12-41 months). For each child, the respondent mother was asked if she had received food assistance either when pregnant or breastfeeding that child. She was also asked if the child had ever been enrolled in the program after s/he was 6 months old, either in the preventive program or the recuperative program. Finally, the respondent was asked if she was currently pregnant or lactating and receiving food assistance. If the response was affirmative for exposure to program benefits for any of the children or for the respondent herself, the household was considered as having ever participated in the program.

Current household participation: The respondents were also asked whether she or any of her children was currently enrolled in the program. If the response was affirmative to any of the questions about current participation, the household was considered as currently participating in the program.

Index child participation: The mothers of the index children were asked if the child had ever been enrolled in the program after the child was 6 months old. They were also asked if they

had been enrolled in the program when pregnant with the index child or breastfeeding the index child. Thus, for each index child, we assessed whether the child had been exposed to the program *in utero*, via the mothers' participation when breastfeeding, or whether the child him/herself was a direct program beneficiary.

Timing and level of exposure: We examined the timing and level of exposure of the child to the program by asking whether the mother had participated when pregnant and breastfeeding and whether the child had participated after s/he was 6 months old. It would be expected that child and maternal exposure during pregnancy and lactation would be the same between the two program groups, but that exposure would be different once the child reached 6 months of age. This is because in preventive program communities, all children 6 months or older were eligible to receive program benefits, whereas in recuperative program communities, only children with malnutrition were eligible.

Participation in Mothers' Clubs: Respondents were asked whether they had attended MCs or not when they were pregnant with or breastfeeding either the index child, a younger sibling, or another child between 12 and 42 months of age. However, since respondents were likely to have attended several MC sessions when their child was enrolled in the food assistance program, we did not directly ask about maternal attendance to MCs, but rather we asked about whether they had ever missed a session, and if so, how frequently she had missed sessions.

Use of Rally Posts: Since the Rally Posts are the entry point into the program, it could be argued that differences in participation between groups could be driven by differences in participation at the RPs. We gathered data on whether children had ever been taken to the RP and whether they had been to the RP in the last month.

6.2.2 Analysis

We examined program participation in each program group, using random effects regression approaches to evaluate whether the participation variables differed significantly between program approaches. We also examined use of Rally Posts in the last month by child age, to evaluate whether the two program approaches led to differential use of the RP services for children in different age groups.

For household and caregiver characteristics, we examined differences by program approach and exposure, using random effects regression approaches to control for the study design.

6.3 Results

6.3.1 Program participation

6.3.1.1 Household/caregiver-level participation/exposure

Table 6.1 provides information on differences between program communities in program participation rates by *any member of the household*,¹⁸ either at the time of the survey or at any

¹⁸ Household participation includes participation by any beneficiary child, or the child's mother when either pregnant or lactating. Given the possibility of multiple children participating in any household,

point in time since the program started, three years before. Results show that current participation is almost 2.5 times higher in preventive compared to recuperative communities. There is, however, no difference between program communities in the proportion of households ever exposed to the program benefits. As expected by design, the average duration of participation at the household level is significantly higher in the preventive group compared to the recuperative group.

Table 6.1 Program participation at the *household* level (includes participation by all children in the household, as well as pregnant/lactating women)

	Preventive	Recuperative
	(%)	
Households currently participating	43.4	17.4*
For households currently participating: total duration of participation (mean (SD))	15.5 (7.0)	12.0 (6.2)*
Households that ever participated (including current participation)	83.2	82.7
For households who ever participated: total duration of participation (mean (SD))	15.8 (5.1)	10.6 (6.2)*

Note: * $p < 0.05$.

6.3.1.2 Child-level participation/exposure

Table 6.2 shows the use of the program services for pregnant and lactating women and index children. In both program communities, enrollment by mothers when they were pregnant with the index child was around 57% and enrollment when they were breastfeeding the index child was 63%. The number of times mothers received food assistance during pregnancy (average of 4 months) and the first 6 months of lactation (average of 5 months) was also similar between program communities.

For children 6 months or older, enrollment patterns are markedly different between program communities, as expected by design (i.e., different targeting mechanisms). Children in the preventive communities, on average, were enrolled at 8 months of age (they are eligible from 6 months on), while children in the recuperative program were enrolled later, on average at 14 months. This was also to be expected, given the timing of growth faltering in this population and the actual time it takes for a child to reach the cutoff points for WAZ that classify them into M2 or M3 malnutrition levels.

Consistent with program design, children in the recuperative program received food for a shorter duration than children in the preventive program (7.5 versus 12). About 73% of the children in the preventive program communities had ever been enrolled in the program, while 28% of children in the recuperative program had ever been enrolled. Current child enrollment in the program was 38% in the preventive communities versus 14% in the recuperative communities.

Timing of exposure: Table 6.3 presents the timing of exposure to the program among index children (and their mothers) in the two program communities. The proportion of children never exposed to the program either *in utero*, while being breastfed, or after they were older than 6 months old is very similar in the program groups. However, among children who have been exposed to the program, the timing of exposure is quite different among the two program groups:

a larger proportion of children in the recuperative group has been exposed only in pregnancy or in early infancy (via their mother's participation), while in the preventive group, slightly over 50% of children have been exposed to the program both through their mother's participation (*in utero* and in the first six months of breastfeeding) and after they were 6 months or older. These patterns of exposure are generally as expected, based on the design of the two approaches.

Table 6.2 Enrollment of index children in the food assistance program, by program group

	Program communities	
	Preventive Percent or Mean (SD)	Recuperative Percent or Mean (SD)
Received food assistance when pregnant (q305)	57.2	58.2
Number of times received food when pregnant	4.0 (2.0)	4.0 (2.0)
Participated in Mothers' Club for pregnant women? (q305)	62.8	62.9
Number of times attended MC when pregnant	5.0 (2.0)	5.0 (2.0)
Received food assistance when breastfeeding	66.3	62.2
Number of times received food when breastfeeding	5.0 (1.0)	5.0 (1.0)
Participated in Mothers' Club for breastfeeding women	69.8	64.8
Number of times participated in MC for breastfeeding women	5.0 (1.0)	5.0 (1.0)
Child <u>ever</u> received food assistance (survey)	73.1	28.2*
Child <u>currently</u> receiving food assistance (survey)	37.6	14.1*
Age when child was first enrolled	7.7 (2.1) Median – 7; Range (6-23)	13.6 (3.9)* Median – 12 Range (7-37)
Age when child last received program benefits	22.8 (2.9) Median – 23 Range (7-36) ¹	21.5 (6.7) Median – 23 Range (7-40)
Total number of times received WSB	11.7 (4.3) Median – 12; Range (1-24)	7.5 (4.3)* Median-7 Range (1-24)

Notes: * $p < 0.05$; ¹ The range in the preventive group goes to 36 months because malnourished children older than 24 months also receive food assistance.

Table 6.3 Timing of exposure for index children, by program group

Timing of exposure	Preventive N = 748 Percent	Recuperative N = 750 Percent	Overall N = 1,500 Percent
Never exposed in pregnancy, lactation, or child level	22.1	24.9	23.5
Only in pregnancy	0.3	3.6	1.9
Only during lactation	0.9	5.7	3.3
Only at child level	10.4	8.8	9.6
Only pregnancy and lactation	3.6	37.6	20.7
Pregnancy, lactation, and child level	52.4	16.5	34.4
Pregnancy and child	0.9	0.5	0.7
Lactation and child	9.4	2.4	5.9

Participation by child age: Table 6.4 presents data on program participation (ever or current) by child age and program. Note that this table includes both index children and their younger siblings. Patterns of participation by child age at the final survey are quite different between program communities, both for any exposure to the program as well as for current participation. In both program groups, there are no children directly exposed to the program in the 0-6 month age group (as expected). In the preventive program group, current and ever exposure rise to over 50% in the 6-11 month age group, while only 5% of children in that age group are exposed to the program in the recuperative group. Among older children—within the age group eligible for the preventive program, i.e., children in the 12-17 and 18-23 month age groups—participation is around 75%, both for ever and current participation. In the same age groups in the recuperative communities, participation is around 18% in the 12-17 month age group and somewhat higher in the 18-23 month age group. For children over 24 months of age, current exposure is very low as expected (since only severely malnourished children over 24 months old are enrolled in the preventive program), while past/any exposure is between 65% and 77% among all age groups above 24 months. The prevalence of severe underweight (WAZ<-3 Z-scores) is 2.2% among children older than 24 months in the preventive program, and this is consistent with the participation rates among children in this age range, especially those over 30 months of age.

Table 6.4 Receipt of food assistance among index children, by child age and program participation

Age group	Preventive			Recuperative			All children		
	n	Ever	Current	n	Ever	Current	n	Ever	Current
Mean participation (index children + young siblings)	906	63.8	54.1	936	22.9	50.7	1,842	43.1	53.1
Mean participation (12-41 mo)	748	73.1	37.6	752	28.2	14.1	1,500	50.6	25.8
- 0-6 mo	99	0.0	0.0	125	0.0	0.0	224	0.0	0.0
- 6-11 mo	60	53.3	53.3	60	5.0	5.0	120	29.2	29.2
- 12-17 mo	197	76.1	75.1	187	19.3	18.2	384	48.4	47.4
- 18-23 mo	153	77.1	73.9	119	27.7	21.1	272	55.5	50.7
- 24-29 mo	180	74.4	7.7	167	29.9	11.9	347	53.0	9.8
- 30-35 mo	115	67.8	2.6	140	32.1	11.4	255	48.2	7.4
- 36 mo and older	102	64.7	1.9	138	34.8	7.9	240	47.5	5.4

In the recuperative program, rates of past/any exposure among children over 24 months of age range between 28% and 35%. Current exposure is highest among children in the 12-17 and 18-23 month age groups, and gradually decreases beyond this age. These patterns mirror the age distribution of malnutrition—and therefore eligibility for the program among children in recuperative communities; i.e., the prevalence of underweight is highest in children 12-23 months of age. Notably, the participation rates are almost halved among children over 24 months of age, compared to those between 12 and 23 months of age, even though the prevalence of underweight is not markedly lower among older children (see Chapter 7). This could suggest that fewer older children are taken to the RPs even in the recuperative communities (since eligibility to receive food assistance is contingent upon children being taken to the RP for screening).

Exposure to MCs: Table 6.5 presents data on exposure to MCs among the respondents. As expected, based on the patterns of participation in the food assistance program, MC exposure

is not different between the program groups in pregnancy and lactation. The results also show that almost 50% of respondents reported missing at least one MC session. The proportion of respondents who reported missing an MC session “only rarely” was higher in the preventive compared to the recuperative group, while those who reported missing sessions “often” was higher in the recuperative compared to the preventive group.

Use of Rally Post services: The use of RPs was not different between program groups when examined in terms of whether an index child had ever been taken to an RP since 2002 (Table 6.5). However, a slightly larger proportion of children in the preventive communities had been taken to the RP *in the past one month* compared to children in the recuperative communities, and the total number of times the child had been taken to an RP in the last year was slightly higher in the preventive communities than in the recuperative communities.

Table 6.5 Participation in Mothers’ Clubs and use of Rally Post services

	Preventive Percent	Recuperative Percent
Mothers’ Club participation		
Participated in Mothers’ Club for pregnant women?	62.8	62.9
Number of times attended MC when pregnant	5.0 (2.0)	5.0 (2.0)
Participated in Mothers’ Club for breastfeeding women	69.8	64.8
Number of times participated in MC for breastfeeding women	5.0 (1.0)	5.0 (1.0)
MC participation by mother when child was enrolled in food assistance program ^a		
- Never missed an MC session	51.3	48.7
- Frequency of having missed MC sessions		
- rarely	40.3	27.4
- sometimes	51.1	58.3
- often	8.6	14.3
Rally Post utilization		
- Percent of children ever taken to RP (between 2002 and 2005)	96.7	97.1
- Percent of children taken to RP in month preceding survey	52.6	49.7
- Number of times taken to RP in one year preceding survey	7.5 (3.2)	7.1 (3.2) ^b

^a All beneficiary children’s mothers are obligated to attend MCs. We did not gather data on the number of MCs attended when children were enrolled in the program, but asked mothers about how often they had missed.

^b $p < 0.05$.

Use of RP services by child age, however, differed between the two program communities, particularly for older children (Table 6.6). Use of RP services was very high for children who were less than 12 months old, and not different between program communities. On the other hand, RP participation rates were higher for children 12-23 months old in the preventive communities compared with the recuperative communities. For children older than 23 months, the proportion of children taken to RPs was higher in the recuperative communities than in the preventive communities. These significant differences could reflect household and caregiver understanding of the targeting mechanisms of the two program approaches, such that in the preventive program communities, caregivers were more diligent about bringing children less than 24 months of age, while in the recuperative communities, they were more likely to bring older children (who might be identified as being malnourished). The lack of difference between program communities in RP use for children younger than 12 months is reassuring, however, since it implies that regardless of the targeting mechanism for food assistance, younger children

are still taken to the RPs in the recuperative communities for immunization and other preventive services.

Table 6.6 RP participation, by child age and program group

Age group	Child was taken to RP in the past month		
	Overall	Preventive	Recuperative
< 12 mo	86.0	88.5	83.5
12-18 mo	72.1	77.9*	65.7*
18-24 mo	64.7	73.5	53.4
24-30 mo	43.2	38.6	47.9
30-36 mo	33.5	28.2	37.8
36-42 mo	31.1	20.6	38.9

* $p < 0.05$ (difference by age for index children 12-41 months of age, within program group).

Program uptake and targeting: We define *uptake* as the proportion of eligible children in each program area who are currently enrolled in the program, and *targeting* as the percentage of children enrolled in each of the programs who are eligible to receive food assistance under that program approach. Since the program services were provided in all the communities covered by the survey, uptake is largely driven by household decisions to bring the child to the RP for enrollment in the food assistance arm of the program. For estimates of uptake as well as targeting, we use data from our final survey data. In preventive areas, uptake was high (75%) and targeting was very effective, with up to 93% of current participants (survey) being in the targeted age range (6-24 months) (Table 6.7). In the recuperative areas, however, only 29% of currently underweight children were enrolled in the program, and only 57% of those enrolled were underweight at the time of the final survey.

Table 6.7 Program uptake and targeting, by program group

	Program communities	
	Preventive Percent	Recuperative Percent
Program uptake (survey) - percent of targeted group who <u>currently participate</u>	74.6	29.1
Targeting: percent of current participants who meet eligibility criteria (survey)	93.2	56.6

It should be noted that estimates of uptake and targeting in the recuperative group cannot be assessed accurately with a cross-sectional survey. This is because the eligibility criterion is underweight at the time of enrollment, not at the time of the survey. In trying to estimate uptake and targeting based on measurements taken at the time of the survey, we introduce errors due to the fact that (1) some children may have recovered since they were enrolled in the program and are no longer underweight (thereby appearing as non-eligible); (2) some children may have become underweight only recently and have not been screened into the program yet (thereby appearing as not covered); (3) some children might have been enrolled in the past but have now relapsed (or never recovered) (also appearing as not currently covered¹⁹).

¹⁹ Note that children in the recuperative group can re-enter the program after one year of exiting from the program, or if they are severely malnourished at the end of the 9-month period. Children in the preventive group re-enter if they are malnourished (WAZ < -2) between 24-59 months of age. The proportion of children previously enrolled who later re-entered the program was 8.5 percent (18 children) in the recuperative approach and 1.5 percent in the preventive approach (8 children).

In summary:

- ❖ Enrollment and duration of participation in the food assistance program during pregnancy and the first six months of lactation was not different between program communities.
- ❖ Enrollment and duration of participation after the child reached 6 months of age was significantly different between the program communities, as expected by design.
- ❖ Timing of exposure to the program benefits was different between the program communities, with more children having been exposed only through their mother's participation (i.e., during pregnancy and lactation) in the recuperative communities, versus a greater proportion of children having been exposed both through their mother's participation and as direct beneficiaries (after 6 months of age) in preventive communities. This was also expected by design.
- ❖ Child age is closely associated with participation rates (particularly current participation). In the preventive program communities, participation is high between 6 and 24 months of age as expected, while in the recuperative communities, participation is highest when children are between 12 and 23 months of age, when undernutrition is at its peak.
- ❖ MC exposure is not different between program groups for participation during pregnancy and lactation. The frequency of missing MC sessions is slightly lower in the preventive group than the recuperative group.
- ❖ Overall RP participation is reasonable (half of the children were taken in the month preceding the survey, and more than 95 percent had ever been taken to the RP) and similar between program groups. Age differences in RP attendance between program groups were quite marked, however, with fewer children above 24 months of age taken to RPs in the preventive communities.
- ❖ There were significant differences in program uptake and efficiency of targeting in the two groups. Uptake was much higher in the preventive than in the recuperative approach, and more than 90 percent of children currently enrolled in the preventive group met the targeting criteria. Although it is difficult to interpret both uptake and targeting in the recuperative group, the results point to differences that could be driven by community and household perceptions about the program.

6.3.2 Household and caregiver characteristics

In this section, we briefly examine some household and caregiver characteristics, focusing on those that are not expected to be influenced directly by the program. In addition to comparing household and caregiver characteristics by program group, we also compare household and caregiver characteristics of program participants and nonparticipants in both program groups. We focus on comparisons of characteristics of household or caregivers who had *ever* participated in the WV MCHN program to those that had never participated in the

program. Since current participation is largely driven by child eligibility (which differs between the two approaches), we consider comparisons based on any exposure to be more relevant to understanding the drivers of participation.

6.3.2.1 Household characteristics

Table 6.8 provides information on household characteristics by program group, and comparing participants and nonparticipants within each program group.

Table 6.8 Household characteristics, by program and by program participation (ever participated)

	Variable	Preventive (n = 748)		Recuperative (n = 752)		Preventive (n = 126)		Recuperative (n = 622)		Preventive (n = 130)		Recuperative (n = 622)	
		Nonparticipant		Participant		Nonparticipant		Participant		Nonparticipant		Participant	
		Mean		SD		Mean		SD		Mean		SD	
		Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
No. of rooms in house	q27a	2.2	0.7	2.2	0.8	2.2	0.6	2.2	0.8	2.1	0.7	2.2	0.8
Household size	hhsz	6.6	2.3	6.6	2.3	6.1	2.2	6.6 ^a	2.3	6.1	2.6	6.7 ^a	2.2
		%		%		%		%		%		%	
Own house	q24a	91.8		90.0		93.7		91.5		86.9		90.7	
-Own house		3.7		5.3		2.4		4.0		7.7		4.8	
Rents		4.4		4.7		4.0		4.5		5.4		4.5	
Free housing													
Own land on which house is	q24b	78.6		75.9		81.7		78.0		73.8		76.4	
Floor material	q27b	94.9		95.1		95.2		94.9		94.6		95.2	
Earth/sand/rock		5.1		4.9		4.8		5.1		5.4		4.8	
Concrete													
Wall material	q27c	25.9		27.8		19.8		27.2		20.0		29.4	
Clissade/earth		3.2		1.1		4.0		3.1		0.0		1.3	
Wood/plank		2.8		4.8		0.0		3.4		6.2		4.5	
Stone blocks/stones		66.0		65.2		73.8		64.5		72.3		63.7	
Palissade													
Roof material	q27d	50.4		50.8		53.2		49.8		50.8		50.8	
Thatched roof		49.6		49.2		46.8		50.2		49.2		49.2	
Aluminum													
Drinking water source	q21	28.2		31.3		22.2		29.4		31.5		31.2	
Public tap water		3.1		1.9		2.4		3.2		3.8		1.4	
Public open well		0.1		0.8		0.0		0.2		0.8		0.8	
In compound covered/protected well		6.8		4.8		8.7		6.4		5.4		4.7	
Protected spring		58.2		56.6		65.1		56.8		56.2		56.8	
Unprotected spring		2.3		3.5		0.8		2.6		2.3		3.7	
River													

^a Difference between participants and nonparticipants (within program group) is significant at $p < 0.05$ (random effects regression analysis).

Overall, there were no significant differences between the program groups in terms of general household characteristics such as house construction, number of rooms in the house, home ownership, and sources of drinking water. There were also no significant differences in these characteristics among participant and nonparticipant households in the two program groups. However, household size was significantly different between those who had ever participated in the program and those who had never participated. In both program groups,

household size was larger among households that had ever participated in the program, compared to households that had never received program benefits.²⁰

6.3.2.2 Caregiver characteristics

Table 6.9 provides information on caregiver characteristics by program group, and comparing participants and nonparticipants within each program group. There were no significant differences between the program groups on any of the maternal characteristics examined. The sets of program communities were similar in terms of major determinants of child nutrition, such as caregiver education, education of the caregiver's partner, marital status, and employment status. The program communities were also similar in terms of characteristics such as maternal BMI, communications between the caregiver and her partner, ownership of assets, control over household purchases, financial and material support, involvement in decisionmaking, and availability of household help.

Maternal work-related characteristics were slightly different between the program communities, but differences were of small magnitude and nonstatistically significant.

When comparing caregivers from households who had *ever* received program benefits with those that had never received benefits (Table 6.9), differences were seen in communications between the respondent and her spouse/partner, women's ownership of assets, and marital status. In the recuperative group, communication between respondent and spouse/partner appeared to be better among participants than among nonparticipants. In both program groups, women who had ever participated in the program scored slightly higher on the asset ownership scale than those who had never been exposed to the program, and a slightly larger proportion of participants were married (or had a partner) compared to women never exposed to the program.

Although Table 6.9 shows some trends toward less time spent on working away from home among participants in both program communities, these differences are not statistically significant. However, they could reflect trends that indicate that women who work away from home for long hours are unable to participate in the program. Conversely, it could also signify a program impact, i.e., that program participation reduced the amount of time that mothers had to spend away from the home on work because the program provided a food transfer to the household.

In summary:

- ❖ There were no significant differences in household or caregiver characteristics among program groups at the time of the final survey, although caregivers in the preventive communities were slightly more likely to be working away from home.
- ❖ There were few differences in household characteristics among participant and nonparticipant households. However, households exposed to the program were likely to be larger in size than households never exposed to the program.

²⁰ Note that all households in both the baseline and the final survey had at least one child between 12 and 41 month of age, as this was a selection criterion for inclusion in the survey.

Table 6.9 Caregiver/respondent characteristics, by program group and by program participation (ever participated)

	Variable	Preventive (n = 748)		Recuperative (n = 752)		Preventive (n = 622)				Recuperative (n = 622)			
		Non-participant		Participant		Non-participant		Participant		Non-participant		Participant	
		Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Education of the respondent mother (years)	<i>edu_resp</i>	2.0	2.7	1.9	2.7	1.8	2.5	2.0	2.7	2.0	2.7	1.9	2.7
Education of the partner/husband	<i>edu_part</i>	3.2	3.5	3.2	3.4	3.1	3.4	3.2	3.5	3.5	3.6	3.1	3.4
Maternal body mass index	<i>bmi</i>	21.8	3.3	21.7	3.5	21.9	3.6	21.7	3.3	21.4	3.3	21.8	3.5
Couple communications scale	<i>q802comm</i>	6.2	1.8	6.2	1.8	6.0	1.8	6.2	1.8	5.7	1.8	6.3 ^a	1.8
Women's ownership of assets (scale)	<i>q803poss</i>	2.1	1.0	2.1	1.1	1.9	1.1	2.2 ^a	1.0	2.0	1.1	2.1 ^a	1.1
Control over purchasing (scale)	<i>q805purc</i>	3.2	2.4	3.3	2.5	3.4	2.4	3.2	2.5	3.6	2.6	3.3	2.5
Financial/material support (scale)	<i>q810supp</i>	1.6	1.1	1.6	1.2	1.7	1.2	1.6	1.1	1.5	1.1	1.6	1.2
Social support (financial/material/ emotional) (scale)	<i>q810supn</i>	2.2	1.5	2.1	1.5	2.3	1.5	2.2	1.5	2.0	1.4	2.1	1.5
Involvement of respondent in HH decision-making	<i>q817invtasks</i>	8.6	3.0	8.8	3.0	8.4	3.2	8.7	3.0	9.1	2.9	8.7	3.1
Number of tasks mother gets help with (scale)	<i>ntasks</i>	5.4	3.0	5.4	3.0	5.2	3.2	5.4	2.9	5.2	3.1	5.4	3.0
Number of childcare tasks mother gets help with (scale)	<i>ch_help</i>	2.1	1.2	2.1	1.3	2.0	1.3	2.2	1.2	2.0	1.3	2.2	1.2
		%		%		%		%		%		%	
Currently pregnant	<i>q34</i>	14.6		16.6		18.3		13.8		18.5		16.2	
Married or partnered	<i>q201</i>												
Yes married		27.3		30.2		22.2		28.3 ^a		24.6		31.4 ^a	
yes, placee (civil union)		62.0		56.3		60.3		62.4		55.4		56.4	
yes, partnered, but partner doesn't live with her		2.0		2.5		3.2		1.8		3.1		2.4	
yes, live with a man		1.6		2.4		3.2		1.3		3.1		2.3	
no, not in union		7.1		8.6		11.1		6.3		13.8		7.6	
Ever been to school	<i>q207</i>	52.8		52.0		52.4		52.9		55.4		51.3	
Schooling level(only those who went to school)	<i>q207c</i>												
primary		90.4		89.6		93.7		89.7		88.5		89.9	
secondary		8.0		7.7		5.6		8.5		9.2		7.4	
higher		0.0		0.1		0.0		0.0		0.0		0.2	
Currently employed	<i>q208</i>	76.6		74.7		77.8		76.4		73.1		75.1	
Respondent's occupation	<i>q209</i>												
owns own land or family land		41.7		42.3		40.5		42.0		39.2		42.9	
owns land of other persons		2.5		1.9		2.4		2.6		0.0		2.3	
business		0.8		0.8		0.8		0.8		0.0		1.0	
market/trade		29.3		26.2		29.4		29.3		27.7		25.9	
office/institution		1.1		1.2		1.6		1.0		0.8		1.3	
manual labor		3.6		3.5		5.6		3.2		5.4		3.1	
unpaid work		0.1		0.1		0.0		0.2		0.8		0.0	
fisherman		1.3		2.3		0.0		1.6		3.1		2.1	
retired		0.0		0.0		0.0		0.0		0.0		0.0	
other		0.0		0.3		0.0		0.0		0.8		0.2	
Unemployed		19.5		21.5		19.8		19.5		22.3		21.4	
Location of work	<i>q209d</i>												
home		16.7		18.9		22.2		15.6		20.0		18.6	
away from home		34.2		34.8		35.7		33.9		32.3		35.4	
both		29.5		24.7		22.2		31.0		25.4		24.6	
Unemployed		19.5		21.5		19.8		19.5		22.3		21.4	
Duration away from home, when working	<i>q209f</i>												
> 1 day		2.7		2.4		3.2		2.6		1.5		2.6	
whole day		22.1		23.9		23.8		21.7		26.2		23.5	
1/2 day		31.0		27.5		24.6		32.3		26.9		27.7	
< 4 hours		8.0		5.7		6.3		8.4		3.1		6.3	
Unemployed/work at home		36.2		40.4		42.1		35.0		42.3		40.0	
Childcare arrangements when works outside	<i>q213</i>												
always bring with her		4.9		6.4		4.0		5.1		5.4		6.6	
always leave with someone else		49.3		45.9		46.0		50.0		46.9		45.7	
both		8.7		6.9		7.9		8.8		6.9		6.9	
Unemployed/work at home		37.0		40.8		42.1		36.0		40.8		40.8	

^a Difference between participants and nonparticipants (within program group) is significant at p<0.05 (random effects regression analysis).

- ❖ Among caregivers ever exposed to the program, differences were seen in communication between the respondent and her spouse/partner, ownership of assets by the respondent, and marital status of the respondents.

6.4 Summary of Results

This chapter examined differences in program participation between the preventive and recuperative program communities, and looked at caregiver and household characteristics associated with having ever participated in the program. Our findings indicate that program participation differences are driven largely by differences in targeting mechanisms for children, since participation during pregnancy and the first six months of lactation are not different between program communities. This leads to substantial differences in the timing and duration of exposure to the program between the program groups. Our assessment of the use of Rally Posts in the two program communities suggests that differences in program enrollment by child age could be driven by the caregivers' understanding of program targeting, since fewer older children are brought to preventive RPs and fewer younger children to recuperative RPs.

We found no significant differences in household or caregiver characteristics among program groups at the time of the final survey, although caregivers in the preventive communities were slightly more likely to be working away from home. There were also few differences in household characteristics between households who had ever participated and those who had never participated in the program. Among caregivers ever exposed to the program, however, differences were seen in communications between the respondent and her spouse/partner, and the respondents' ownership of assets and marital status.

In conclusion, our assessment is that differences in program participation between preventive and recuperative communities appear to be largely driven by the design and targeting mechanism of the two program approaches, rather than differences in household or caregiver characteristics. However, a more extensive analysis of the determinants of participation to develop typologies of participants and nonparticipants based on the data on women's work patterns and other characteristics could be useful to provide a deeper and more nuanced understanding of the drivers of participation.

7. NUTRITIONAL IMPACT OF THE PREVENTIVE APPROACH COMPARED TO THE RECUPERATIVE APPROACH

7.1 Introduction

This chapter presents the main results of the effectiveness study comparing the impact of the preventive and recuperative approaches on attained growth and on the prevalence of undernutrition (wasting, stunting, and underweight). It first presents the results of the probability cluster randomized design and follows with additional results that document the plausibility of the results.

7.2 Objective of the Impact Evaluation

The main objective of the impact evaluation was to compare the effectiveness of the preventive and the recuperative approaches in reducing community-level childhood undernutrition. Our main hypothesis was that, compared to the recuperative approach, the preventive approach would significantly increase the mean WAZ, HAZ, and WHZ and reduce the prevalence of stunting, underweight, and wasting among children 12-41 months of age who lived in the program communities and had the potential to have been exposed to the program in the previous three years (see below).

7.3 Intervention and Comparison Groups

The intervention packages compared are described in detail in Chapter 4. As a reminder, the preventive approach differs from the recuperative approach in three main aspects (see Table 7.1 for summary): (1) the targeting mechanism (undernutrition for recuperative and age for preventive); (2) the focus and timing of the BCC intervention (care and feeding of the undernourished child for recuperative; optimal care, and feeding practices for 6-23-month-old child for preventive); and (3) the timing and duration of eligibility (9 months from the time the child is undernourished for recuperative; 18 months from the time the child is 6 months of age). Both approaches include a similar package of interventions for pregnant women and for lactating women until their child reaches 6 months of age.

Table 7.1 Key differences between recuperative and preventive approach intervention packages

Intervention component	Recuperative approach	Preventive approach
Targeting mechanism	Children 0-59 months of age with weight-for-age Z-scores (WAZ) < -2	All children 6-23 months of age ^a Children 24-59 months severely malnourished (M3) ^b
Focus of BCC intervention at Mothers' Clubs	Care and feeding of undernourished child; note that other learning sessions on child-feeding and care practices may also be used.	Timely delivery of age-specific, relevant, and action-oriented messages on optimal breastfeeding, caregiving, and complementary feeding practices of children 6-23 months of age.
Timing of eligibility	When undernourished	When 6-23 months of age
Duration of eligibility to receive food and BCC	9 months (originally determined by WV)	18 months (entire period when the child is between 6 and 23 months of age)

^a Note that children 24-59 months of age with WAZ < -3 in the preventive group are eligible for program benefits for up to 9 months, similar to those in the recuperative group.

^b Severely malnourished children (M3 according to the Gomez classification) are those whose weight-for-age is between 60 and 75% of the median of the weight-for-age CDC/NCHS/WHO standards (Cogill 2003).

7.4 Methods

7.4.1 Evaluation design

The evaluation used a community-level randomized pre-post design, whereby 10 paired clusters of communities were randomly assigned to either the preventive or the recuperative program group. The baseline survey was conducted between May and September 2002 and the post-evaluation survey was conducted exactly three years later, between May and September 2005, to minimize seasonal variations. All components of the intervention packages, except the newly developed BCC strategy, were implemented immediately following the baseline survey, i.e., in August-September 2002. The full BCC package, however, was implemented several months later (in May 2003).

The main outcomes of the evaluation were mean height-for-age Z-scores (HAZ), weight-for-age Z-scores (WAZ) and weight-for-height Z-scores (WHZ), and the prevalence of childhood stunting, underweight, and wasting.

Details about the evaluation design, including sample size estimation, matching and selection of community clusters, the age groups selected for the impact assessment, and the survey design, are described in Chapter 2, Section 2.4.1.1.

7.4.2 Data analysis

Data used in this impact analysis were obtained from the surveys described in Chapter 2 and in Annex 2.1.

The outcomes of interest of the impact evaluation were mean HAZ, WAZ, and WHZ, and the prevalence of stunting, underweight, and wasting. We used the new WHO reference standards to derive these indicators based on children anthropometric measurements (WHO 2006). Differences between program communities in the mean HAZ, WAZ, and WHZ were tested using a pair-wise comparison at the cluster level (and a paired t-test for statistical significance). Additional analyses used random effects regression modeling with child-level data adjusting for the clustering at the pair level (Murray 1998) to test differences between the groups in child anthropometry, adjusting for child age and gender, and maternal height and schooling. Note that the approaches did not control for additional caregiver or household characteristics because many of these determinants of child anthropometry may have been impacted by the program, e.g., household socioeconomic status, food security, and caregiver knowledge and feeding practices. Differences in the prevalence of stunting, underweight, and wasting at the cluster level were tested using random effects logit approaches (xtlogit in Stata 9) that adjusted for the clustering at the zone level and controlled for child age, gender, maternal height, and schooling.

Our power calculations were not conducted to detect differences between program communities in *changes over time* and thus, the statistical significance of these differences is not

reported. Where relevant, differences between baseline and end line within each intervention group (preventive and recuperative) are reported.²¹

7.5 Results

The results section is organized as follows. The first subsection presents the results of comparisons between the preventive and recuperative group at baseline to assess the success of the randomization process. The main impact results based on the community-level probability design are presented next, and focus on difference between the two groups at final survey for the three main anthropometric outcomes: mean HAZ, WAZ, and WHZ, and the prevalence of stunting, underweight, and wasting. The next subsection examines the plausibility of the results, describing changes from baseline to final survey and addressing issues such as dose-response and age-specific response to the intervention. The final subsection compares the prevalence of morbidity symptoms between the preventive and recuperative groups.

7.5.1 Baseline characteristics

Results of the paired cluster-level comparison of mean HAZ, WAZ, and WHZ presented in Table 7.2 show that there were no differences between the program groups in nutritional status at baseline. The same is true for differences in the prevalence of stunting, underweight, and wasting tested using random effects regression approaches to adjust for clustering at the zone level. Similarly no differences were found between the two groups in individual-level comparisons of child age, gender distribution, or reported child-feeding practices (Table 7.2).

Comparisons between the two program groups to test differences in community, household, and caregiver resources are also presented in Table 7.2. The findings suggest that the randomization process was effective because very few differences between the groups were found in spite of the large number of variables compared. For instance, at the community level we found that the pairs of clusters were largely comparable in terms of key geographic characteristics and access to various services such as closest town, school, market, and health services (not shown). Caregiver and household characteristics of the two groups were also very comparable (see Table 7.2 for subset of variables compared). For instance, no differences between the groups were found in caregiver age, education level, occupation, or in gender of household head, household size, or access to basic water services.

In sum, the recuperative and preventive program communities were very similar at baseline, suggesting that randomization was successful.

²¹ Even though our sample size was estimated for comparisons between program groups at the endpoint, they are adequate for examining the significance of change *within* each group between baseline and final surveys. However, our sample size is not adequate to detect the statistical significance of the difference of differences.

Table 7.2 Comparison of child, maternal, and household characteristics between program groups at baseline^a

Indicator	Recuperative	Preventive
	[n = 10 clusters] Mean (SE)	[n = 10 clusters] Mean (SE)
Nutritional status indicators – cluster level^b		
Height-for-age Z-score (HAZ)	-1.65 (0.10)	-1.69 (0.04)
Weight-for-age Z-score (WAZ)	-1.02 (0.06)	-0.97 (0.08)
Weight-for-height Z-score (WHZ)	-0.18 (0.03)	-0.18 (0.05)
Other child characteristics (individual level)^{c,d}		
	N = 792	N = 7881
Stunting prevalence (%)	37.4	36.7
Underweight prevalence (%)	17.8	17.6
Wasting prevalence (%)	4.3	5.2
Age (mean, SD)	29.4 (7.6)	29.3 (7.9)
Gender (% female)	48.0	51.4
Breastfed within 1 hour (%)	19.3	16.2
Fed meals at least minimum recommended number of times (3 times/day) at 12-23 months (%)	58.7	57.6
Mean number of food groups consumed by child (mean, SD)	5.1 (1.5)	5.0 (1.5)
Consumed meat, fish, or eggs in previous 24 hours (%)	87.3	89.2
Caregiver characteristics^d (n = 1,514)		
	N = 765	N = 759
Age (mean, SD)	30.8 (7.0)	30.8 (8.0)
Maternal height at baseline (mean, SD)	157.9 (11.6)	157.6 (15.4)
Years of schooling (mean, SD)	1.4 (2.3)	1.6 (2.5)
Never attended school (%)	53.2	50.7
Occupation		
- % Unemployed	16.1	16.5
- % Farming	43.1	42.5
- % Trade/market	32.7	32.0
Household characteristics^d (n = 1,514)		
	N = 765	N = 755
% male head	90.8	90.1
Occupation of head		
- % Unemployed	2.0	1.4
- % Farming	85.5	86.8
Household size (mean, SD)	6.8 (2.3)	6.7 (2.3)
% who own house	94.1	91.1
% who have electricity	2.1	1.9
% who have sanitation facility	57.3	56.0
% who have tap water in the house (%)	1.6	0.9

^a None of the differences between the groups were statistically significant.

^b Differences in mean were tested using cluster-level pair-wise comparisons and paired t-test.

^c Differences in the prevalence of stunting, underweight, and wasting were tested using a random effects logit model controlling for cluster effect.

^d Differences in other child, caregiver, and household characteristics were tested at the individual level, with t-tests (for means) and chi-square tests (for proportions).

7.5.2 Intervention impact: Results from probability design analysis

Table 7.3 shows the differences in mean anthropometric outcomes between the two program communities at the end of the intervention period. Both unadjusted, cluster-level means, and means adjusted by random effects regression modeling for cluster effect and for child age and gender are presented. The preventive group had higher mean HAZ, WAZ, and WHZ than the recuperative group. All differences were statistically significant, except the unadjusted difference in mean HAZ. Results of random effects regression approaches controlling for child age and gender, however, show that all three mean anthropometric indicators are significantly higher for the preventive compared to the recuperative group. Additional models controlling for maternal height and schooling in addition to child characteristics showed similar results (not shown).

Table 7.3 Mean anthropometric outcomes at final survey, by program group

Child anthropometric outcome	Recuperative			Preventive			Difference (preventive – recuperative)
	n	Mean	SE	n	Mean	SE	
HAZ							
Unadjusted (n =10 clusters/group) ^a	10	-1.68a	0.05	10	-1.53	0.06	0.15
Adjusted for child age and gender ^b	746	-1.67*	0.05	735	-1.53	0.05	0.14
WAZ							
Unadjusted (n clusters) (ICC: 0.021)	10	-1.21*	0.04	10	-0.97	0.06	0.24
Adjusted for child age and gender	746	-1.20*	0.05	735	-0.96	0.05	0.24
WHZ							
Unadjusted (n =10 clusters)	10	-0.46*	0.04	10	-0.23	0.06	0.23
Adjusted for child age and gender	746	-0.46*	0.05	735	-0.22	0.05	0.24

Note: * Statistically significant ($p < 0.05$).

^a Statistical significance of differences in unadjusted means was tested using a paired t-test of cluster level means.

^b Mean random effects regression models were used to analyze child-level data and adjust for the clustering at the pair level and for child age and gender.

Differences between program communities in the prevalence of stunting, underweight, and wasting at final survey are shown in Figure 7.1. At the end of the study, the prevalence of stunting was 4 percentage points lower among children from the preventive (33.9%) compared to the recuperative group (38.2%), while for underweight the difference was 6 percentage points (14.8% for preventive vs. 20.8% for recuperative) and for wasting, the differences in favor of the preventive group was 4 percentage points (3.7% vs. 7.4% for preventive and recuperative, respectively).


7.5.3 Results of plausibility analysis

Two sets of additional analyses were conducted to assess the plausibility of the results. The first one is an analysis of differences between preventive and recuperative communities among the age group that was estimated to have the greatest potential to respond to the intervention; this group is composed of children 24-35 months of age at the end of the study—i.e., children who had been exposed to the preventive program benefits from the time when they were 6-23 months of age. Anthropometric measures at final survey were compared between these fully exposed children and those who had been exposed only partially, i.e., who were either 6-23 months at final survey, or were 36-41 months. The second analysis examines the difference between groups in the nutritional status of children who were in their first year of age at the time of the final survey. These children have been minimally exposed to the interventions themselves (since direct benefits for children start at 6 months of age), but their mothers were exposed while pregnant and lactating. Since the two programs offer exactly the same services to pregnant and lactating women, we did not expect any major differences in the growth of children in the first 6 months of age, but possibly thereafter.

Before assessing these aspects of plausibility, we also examine changes from baseline in the prevalence of stunting, underweight, and wasting (see Figure 7.1). Baseline results are combined for both program groups because they did not differ at baseline. Using adjusted prevalences for baseline and final surveys in both groups, the results indicate that stunting decreased by 3.5 percentage points since baseline in the preventive communities, while it went

up very slightly in the recuperative communities. The prevalence of underweight went up by 2.8 percentage points in the recuperative communities, while it went down by 2.8 percentage points in the preventive communities. Wasting went up by 4.0 percentage points in the recuperative areas while it decreased slightly in the preventive areas.


Figure 7.1 Prevalence of undernutrition among children 12-41 months at baseline (groups combined) and end line, by program group



Notes: Groups were not different at baseline. Random effects logit models adjusted for child age and sex were used to assess statistical significance of differences between program communities. P values are for differences between preventive and recuperative at final survey. Numbers in figure are adjusted prevalences derived from the random effects logit models.


When looking at the magnitude of differences between program communities for the age range of children exposed to the intervention between 6 and 23 months of age, larger effect sizes are indeed observed for most indicators (Figures 7.2 and 7.3). Among the children 24-35 months at final survey, differences in mean anthropometric indicators between preventive and recuperative communities were larger than for the other age groups (differences in favor of the preventive group were +0.21 Z-scores for HAZ, +0.34 for WAZ, and +0.27 for WHZ). Differences in favor of the preventive group in the *prevalence* of underweight were also of larger magnitude in this age group compared to the sample as a whole (7.3 percentage points). For stunting, however, large differences were seen both in the 12-23 and the 24-35 month age groups (6.5 and 6.3 percentage points), but not in the older age group (36-41 months). In fact, in the older age group, stunting was lower in the recuperative areas compared to the preventive (but

Figure 7.2 Mean anthropometric outcomes, by child age and program group at final survey



Notes: Random effects linear regression models adjusted for child age and sex were used to assess statistical significance of differences between program communities for each age group. P values are for differences between preventive and recuperative at final survey. Numbers in figure are adjusted means derived from the random effects models.

Figure 7.3 Prevalence of stunting, underweight, and wasting, by child age and program group at final survey




Notes: Random effects logit models adjusted for child age and sex were used to assess statistical significance of differences between program communities for each age group. P values are for differences between preventive and recuperative at final survey. Numbers in figure are adjusted prevalences derived from the random effects logit models

nonsignificant, $p>0.5$). For wasting, larger differences were seen among the 12-23 month age group (4.7 percentage points) but not for the other age groups, where differences were similar to the overall differences in wasting between the groups. Since the periods of 12-23 and 24-35 months of age are those of greater nutritional vulnerability, these results are largely as expected.

The level of utilization of the different program services in preventive and recuperative communities is discussed in Chapter 6. One point worth highlighting in the context of the impact of the program on nutritional status is the fact there were no differences between program groups in the mothers' participation during pregnancy and lactation (see Table 6.2, Chapter 6). Differences in participation between the groups start when the child is 6 months of age, as expected, because by design the two programs have different eligibility criteria for children 6 months and older (age for the preventive group; and malnutrition for the recuperative group).

Figure 7.4 shows the mean weight-for-age Z-scores of index children (12-41 months) and of their younger siblings (0-11 months). Although the children in the preventive group track above the children in the recuperative group at all ages from 0 until 41 months of age, the differences are stronger after about one year of age. This pattern was expected since mothers in the preventive and recuperative groups received exactly the same package of intervention until the child reached 6 months, and participation in both groups was similar. On average, children

Figure 7.4 Weight-for-age Z-scores of index children (12-41 mo) and their younger siblings (0-11), by age and program group at final survey




Note: Numbers in figure are adjusted means derived from random effects regression models comparing the program communities and adjusting for child age and sex.

in the preventive approach entered the program at approximately 7 months of age, and thus, it is highly plausible that differences in children's growth between the two approaches start becoming significant only after this age. A striking finding of the WAZ curves shown in Figure 7.4 is the steep drop in mean WAZ observed during the first year, in spite of the fact that more than half of the mothers in both program areas participated in the program during pregnancy and the first 6 months of lactation. Similar drops were also observed for HAZ (not shown).

7.5.4 Differences in morbidity between the two program groups


Figures 7.5-7.8 present the differences between the preventive and recuperative groups in the prevalence of symptoms of infections. Overall, there were no consistent differences between the groups, except for fever, where the recuperative group had a consistently higher prevalence than the preventive group (the differences were statistically significantly different only among the 24-29 month old children, however).

Figure 7.5 Percentage of children who had fever in the past two weeks, by age and program group (final survey)




Note: * $p < 0.05$ using separate random effects logit models for each age group, adjusting for child sex and comparing the program communities.

Figure 7.6 Percentage of children with cold/cough in the past two weeks, by age and program group (final survey)




Note: Statistical significance was tested using separate random effects logit models for each age group, adjusting for child sex and comparing the program communities.

Figure 7.7 Percentage of children with diarrhea in the past two weeks, by age and program group (final survey)



Note: Statistical significance was tested using separate random effects logit models for each age group, adjusting for child sex and comparing the program communities.

Figure 7.8 Percentage of children with fast breathing in the past two weeks, by age and program group (final survey)



Note: Statistical significance was tested using separate random effects logit models for each age group, adjusting for child sex and comparing the program communities.

7.6 Summary of Results

7.6.1 Probability results

- ❖ There were no differences between the two program communities at baseline in mean anthropometric indicators or in the prevalence of stunting, underweight, or wasting.
- ❖ At final survey, children in preventive communities had higher mean HAZ, WAZ, and WHZ, and lower prevalence of stunting, underweight, and wasting than children in recuperative communities. All differences were statistically significant.
- ❖ The differences in mean HAZ, WAZ, and WHZ (adjusted for child age and gender and clustering effect) in favor of the preventive group were +0.14, +0.24, and 0.24, respectively. For the prevalence of stunting, underweight, and wasting, differences in favor of the preventive group were 4, 6 and 4 percentage points, respectively. The magnitude of these effects is modest, but comparable to effects seen in other studies

that compared baseline and post-intervention or intervention and control group at the end of an intervention (Caulfield, Huffman, and Piwoz 1999; Swindale et al. 2004).

7.6.2 Plausibility results

- ❖ Compared to baseline, children's nutritional status appears to have deteriorated among the recuperative group, especially with regards to the prevalence of underweight and wasting; by contrast, wasting remained the constant over time in the preventive group, and both underweight and stunting decreased over time. These results suggest that the preventive approach may have helped mitigate the deleterious effects on childhood malnutrition of the economic and political crisis that occurred in Haiti during the study period.
- ❖ Children who were exposed to the preventive approach for the entire period between 6 and 23 months (i.e., who were 24 to 35 months at final survey) benefited more from the intervention than children who were only partially exposed. This was particularly true for mean HAZ, WAZ, and WHZ, and underweight. For stunting, children 12-23 months old and 24-35 months old showed larger benefit, while for wasting, children 12-23 months old showed the largest benefit.
- ❖ Analysis of younger siblings of index children (i.e., infants 0-11 months of age at the final survey) shows no difference between the preventive and recuperative groups - a result that was expected, given that the two programs offered exactly the same preventive services to mothers during pregnancy and the first 6 months of lactation. Neither program approach, however, succeeded in preventing the steep decline in mean anthropometric indicators during the first year of life.
- ❖ No differences in morbidity symptoms were found between the preventive and recuperative groups, with the exception of fever, which was less prevalent among children 24-29 months of age in the preventive group.

7.7 Conclusions

The study confirms that the preventive program approach, which targets food assistance and BCC to all children 6-24 months, is more effective at reducing undernutrition than the recuperative approach, which targets malnourished children younger than five years of age. The evidence of the greater effectiveness of the preventive over the recuperative program approach is strong from a probability perspective. It is also highly plausible, given the larger impact found among children fully exposed to the program, compared to those who were exposed only partially during the period of greatest nutritional vulnerability (i.e., when they were 6-23 months of age).

The following three chapters explore some of the intermediary mechanisms that may be responsible for the greater effectiveness of the preventive compared to the recuperative approach. These include changes in maternal knowledge and practices related to child feeding and care (Chapter 8), improved household food security (Chapter 9), and other potential household-level socioeconomic benefits (Chapter 10).

8. IMPACT OF THE PROGRAM ON KNOWLEDGE AND PRACTICES²²

8.1 Introduction

This chapter describes the impact of the program on the anticipated outcomes of the behavior change communications activities, i.e., caregiver knowledge and practices. In examining the impact of the program on these outcomes, we attempt to track different stages leading to sustained behavior change, moving from knowledge to trial and adoption of recommended child-feeding and care practices, to childcare practices reported by mothers in the final survey. Throughout, we maintain a focus on the feeding and care practices promoted in the Mothers' Club (MC) learning sessions described in Chapter 4, which constituted the main venue for delivery of the BCC.

8.2 Data and Variable Creation

8.2.1 Maternal knowledge

We assessed caregiver knowledge related to infant and child-feeding practices using the knowledge questions from the baseline survey instrument as well as a more extensive knowledge test that was developed for the operations research (OR) study conducted in 2004. The questions from the baseline survey focused on knowledge about the ideal duration of breastfeeding, the appropriate introduction of complementary foods, and appropriate feeding frequency for children in different age groups. The knowledge test from the OR-2004 questionnaire was specifically designed to assess the respondent's knowledge about topics taught at the MCs and Rally Posts (RPs). Thus, in addition to questions regarding infant and young child feeding, this knowledge test also included more general health-related questions such as prevention of HIV infection, danger signs in pregnancy, prevention of worms, hand-washing practice, etc. Using both of these sets of questions, we constructed a variety of scales to capture respondent knowledge about child feeding and care. These are described below.

8.2.1.1 Infant and Young Child-feeding Knowledge Test (same as in baseline questionnaire)

The survey instrument included a module on appropriate timing of introduction of complementary foods and one on appropriate feeding frequency for infants and children of different ages. Both are described briefly below.

Knowledge of timing of introduction of complementary foods. The questionnaire assessed respondents knowledge related to the introduction of liquids and foods from six different food groups (water/liquids, semisolids, staple foods, vegetables, eggs, and meats) to infants, using types of liquids and foods commonly fed to infants and young children in this area (Menon et al. 2002b). Respondents were asked when (in months after the birth of an infant) they thought it was appropriate to introduce each liquid or food. The data on the age of introduction of

²² Mary Arimond led the writing and analysis for sections pertaining to infant feeding practices in this chapter. Purnima Menon and Mduduzi Mbuya conducted analysis and writing pertaining to maternal knowledge, as well as awareness, trial, and adoption of recommended practices.

individual liquids and foods were then used to create a summary scale of the overall knowledge of a respondent about appropriate introduction of foods, using the current PAHO/WHO Guiding Principles on complementary feeding to define appropriateness (PAHO/WHO 2003). For each food group, the scoring distinguished between appropriate introduction (i.e., at between 6 and 8 months of age) and introduction that was either too early (before 6 months of age) or too late (9 months or older). Introduction of any of the foods in the food group at the appropriate time was assigned a score of 1, and introduction of any of the foods in the inappropriate window was assigned a score of 0. The scores for a total of six food groups were then added up to create a summary knowledge scale that ranged from a possible minimum of 0 to a maximum of +6. Details regarding the scoring are provided in Annex 8.1.

Knowledge of appropriate feeding frequency. Six questions in the feeding knowledge questionnaire asked respondents about their knowledge of the appropriate frequency of feeding meals and snacks to infants and young children in three different age groups (6-8 months, 9-11 months, and 12-23 months). These data were combined to create an overall scale that assessed the knowledge of appropriate feeding frequency. The scoring of the variables was based on current age-specific recommendations on the frequency of feeding complementary foods to breastfed children between 6 and 23 months of age (PAHO/WHO 2003). The detailed scoring is presented in Annex 8.1. Based on this scoring, respondents whose answers for each age group matched or exceeded the currently recommended meal and snack frequency for that age group received a score of 1, while those whose responses indicated a lower frequency than the recommendations received a score of 0. The scale thus created went from a possible minimum of 0 to a maximum of +6.

Overall feeding knowledge (weighted scale). The two scales described above—knowledge regarding introduction of new foods and knowledge of age-appropriate feeding frequency—were combined with information on knowledge about the optimal duration of breastfeeding to create a scale to assess overall feeding knowledge. The two 6-point scales were each divided by 3, to contribute up to 2 points each to the overall feeding knowledge scale; the overall scale ranged from 0 to +6.

The respondents' knowledge about the optimal duration of breastfeeding was assessed by asking then how many months they believed infants and young children should be breastfed. Responses to this question were also coded based on the current recommendations for feeding infants and young children, i.e., that children should be breastfed up to at least 24 months of age. Responses that were further away from 24 months received lower points than responses closer to 24 months, e.g., a response between 0 and 5 months received -2 points, while a response of 24 months and beyond received the maximum of +2 points.

The points on breastfeeding knowledge were summed with the points on the two other weighted scales to create an overall feeding knowledge scale that ranged from a possible minimum of 0 to a possible maximum of +6. As with the previous feeding knowledge scales, the detailed scoring for this overall scale is presented in Annex 8.1.

8.2.1.2 General Nutrition and Health Knowledge Test - based on topics covered at MCs and RPs

Box 8.1 presents the questions asked on the knowledge test related to the health and nutrition topics covered at the MCs and RPs. Responses were evaluated relative to the discussion of those topics at the MCs (or, where relevant, the RPs). Each variable was scored to contribute one point to an overall knowledge score. For most open-ended questions, we assigned one point on the question/topic if even one correct response was provided by the mother. For a few questions, certain responses were considered much more appropriate or important than

Box 8.1 Knowledge test questions on topics covered at MCs and RPs

a. Breastfeeding knowledge (maximum score: 8)

- Until what age do you think you should breast feed your child?
- How long after birth should a baby start breastfeeding?
- What should a mother do with “first milk” or colostrum?
- What should a mother do if she thinks her baby is not getting enough BM?
- Should infants < 6 be given water in addition to BM if the weather is hot?
- What should the baby < 6 months be fed if mother needs to be away?
- Can a mother who is not well fed produce enough breast milk?
- What are the things you can do to increase milk production?

b. Complementary feeding knowledge (maximum score: 9)

- What are the special foods mothers could make to complement breast milk?
- Will feeding a child enriched gruel in the evening cause indigestion?
- How many times per day should a child 6-8 months old eat (meals and snacks)?
- How many times per day should a child 9-11 months old eat (meals and snacks)?
- How many times per day should a child ≥ 12 months old eat (meals and snacks)?
- At meal times, how much food should a child 6-9 months old be offered?
- At meal times, how much food should a child ≥ 12 months old be offered?
- Should a one-year old child eat only the same foods as the rest of the family?
- What are some of the things you can do to encourage young children to eat food?

c. Knowledge about child illness (maximum score: 4)

- Do children need an extra meal per day after they have been sick?
- For how long do children need an extra meal after they have been sick?
- What should you do when your child has diarrhea?
 - o Oral rehydration salts (ORS)
 - o Food and diet-related answers

d. Knowledge about general health issues (maximum score: 5)

- What can you do to protect a child against polio?
- What can you do to protect a child against HIV/AIDS?
- When should you wash your hands?
- How can you protect a child from getting worms?
- How can you make drinking water safe?

e. Knowledge about malnutrition (maximum score: 2)

- Why do you think children get malnourished?
- How can we help malnourished children recuperate?

others; in such cases, the most appropriate response(s) was assigned a full score (1 point), and the less appropriate response(s) was assigned a partial score (0.5 point). And finally, for the question that asked about the use of the lactation amenorrhea method (LAM), only those respondents who mentioned *all* three necessary criteria for LAM were assigned a point. The coding of correct responses to the questions is shown in Annex 8.2.

The scores for a total of 28 questions were then added up to create a summary knowledge scale that ranged from a possible minimum of 0 to a maximum of 28. The total score on the knowledge scale was also transformed into a variable that provided information on the proportion of the 28 questions the respondent had answered correctly. We also created more focused knowledge scales to capture respondent knowledge about specific topics, i.e., on breastfeeding, complementary feeding, feeding sick children, general health, and malnutrition (using the questions shown in Box 8.1, a through e).

8.2.2 Trial and adoption of key recommended practices

The BCC strategy was developed comprehensively to improve overall infant and young child-feeding practices. At the same time, it promoted certain specific feeding practices and recipes that were developed based on the formative research and recipe trials described in Chapter 4 (Menon et al. 2002b). Based on the theory that behavior change occurs in stages (Prochaska and DiClemente 1983), we used qualitative and quantitative approaches to understand the stages through which mothers exposed to new practices would have to progress for the program to have an impact, i.e., being aware of the new practice, then trying it, and finally, adopting the new practice. Thus, we first examined awareness, trial, and adoption of these key new practices using qualitative research methods in the operations research study described in Chapter 5 and in more detail in Menon et al. (2005). We then designed a survey module for examining awareness, trial, and adoption based on the results and insights from this qualitative research for inclusion in the final impact survey.

Focusing on seven key recommended practices and new recipes (Box 8.2), we asked respondents who had been exposed to the MCs if they had ever heard about the practice at the MCs, whether they had ever tried it at home and if so, how often they had tried it. We also asked about reasons for never trying a practice or for not adopting it, defining “adoption” as doing the practice more than just 1 or 2 times. Furthermore, for those women who had heard about the practice at the MCs, we also asked if the practice had ever been demonstrated to them, since the qualitative research about trial and adoption showed that demonstrations at the MCs encouraged trial of the practice at home.

For women who had never been exposed to the MCs, we asked the same sequence of questions about awareness, trial, and adoption, and reasons for not trying a practice or not practicing it more than a few times. However, instead of asking about whether the practice had been demonstrated to them in the MCs, we asked where they might have heard about the new practices recommended by the program. This latter question was intended to provide some ideas about the potential diffusion of the program messages outside of the MCs.

Box 8.2 From awareness to trial and adoption: Key practices included in the survey

- Leaving expressed breast milk behind for baby when leaving the house
- Using expressed breast milk in gruels
- Adding an egg to the child's portion of gruel
- Preparing enriched gruel with beans and groundnuts
- Preparing mashed plantain with added pumpkin
- Feeding enriched gruel in the evenings
- Feeding an extra meal for two weeks after the child recovered from an illness

8.2.3 Child-feeding and care practices

8.2.3.1 Indicators pertaining to the Guiding Principles for feeding infants and young children

The material covered in the MC learning sessions closely followed current international “Guiding Principles” for infant and young child feeding (PAHO/WHO 2003; WHO 2005). Our baseline and final surveys were designed to capture information reflecting practices taught and promoted in MCs and summarized in the Guiding Principles (see Annex 8.3 for Guiding Principles for feeding breastfed (a) and non-breastfed (b) children). Table 8.1 lists the Guiding Principles, and related recommended practices, as well as data collected and indicators constructed to assess them. Practices related to some recommendations are not easily captured via maternal recall in simple surveys. For example, our survey did not include quantitative data collection on consistency and quantity of food offered/eaten, nor on energy and nutrient density of foods. Survey responses on recall of feeding and care practices may also be biased, when respondents overreport good practices. Data on hygiene practices are particularly susceptible to this type of bias (Ruel and Arimond 2002). For some practices that are difficult to capture via recall, proxy measures are identified in Table 8.1.

8.2.3.2 Constructed indicators for child feeding

Results for most of the indicators listed above and in Table 8.1 are presented as simple percentages reflecting responses to single questions. A few exceptions (constructed indicators) are briefly described below.

Exclusive breastfeeding (last 24 hours). The percent of children exclusively breastfed in the last 24 hours was calculated by considering responses to a series of questions asking caregivers if their child had been given any of a long list of liquids or solid food groups the previous day. If the child was reported to be still breastfeeding but not to consume any of the other liquids/foods listed, s/he was categorized as exclusively breastfed. For those so categorized, we cross-checked against other variables in the data set (recalled age of introduction of various liquids/foods, and frequency of feeding in the past 24 hours). In all cases, the three sets of questions were consistent. Note that exclusive breastfeeding in the last 24 hours is not equivalent to exclusive breastfeeding since birth (which is much harder to ascertain by recall). However, for purposes of comparing between groups, and assessing changes since baseline,

Table 8.1 Guiding Principles^a and recall data collected on recommended practices

Guiding Principle – aspect of feeding	Guidance/recommended practices discussed in MCs	Indicators used in Haiti survey
Duration of exclusive breastfeeding (BF) and introduction of complementary foods	<ul style="list-style-type: none"> - Use good BF practices on day of child's birth (initiate BF in first hour, give colostrum, practice exclusive BF) - BF exclusively for 6 mo; introduce complementary foods (CF) at 6 months of age while continuing to breastfeed 	<ul style="list-style-type: none"> % initiating BF within one hour of birth; % exclusively BF; % using colostrum % of children 0-5.9 mo exclusively BF in past 24 hours Recalled age of introduction of liquids and semi-solid and solid foods
Maintenance of breastfeeding	Continue frequent, on-demand BF until 2 years and beyond	% BF in past 24 hour by age group
Responsive feeding	Feed infants directly, assist older children; if child refuses foods, use different methods of positive encouragement	% reporting use of positive strategies (caressing, playing, offering other choices of food) when child refuses to eat
Safe preparation and storage of foods	Practice good hygiene and proper food handling; this includes hand washing and avoidance of baby bottles	% "usually" using baby bottles
Amount of complementary food needed	"Ensure child energy needs are met." Not measured because requires total energy intake from food and breast milk	No indicator
Food consistency	<ul style="list-style-type: none"> Increase food consistency and variety as infant gets older; from 6 months on, infant can eat pureed, mashed, and semi-solid foods) No direct measurement of consistency; recipes promoted and practiced in Mothers' Clubs were of appropriate consistency 	<ul style="list-style-type: none"> % children given semi-solid foods beginning at 6-6.9 mo % fed recipes promoted in MCs (past 24 hr, 3+ or 7+ times last wk)
Meal frequency and energy density	<ul style="list-style-type: none"> Increase number of times child is fed complementary foods as s/he gets older; BF children, feed: 2-3 times for 6-8 mo; 3-4 times for 9-23 mo; Non BF, feed: 3-4 times for 6-8 mo; 4-5 times for 9-23 mo Energy density: no direct indicators, but promoted recipes were formulated to be of adequate energy density 	<ul style="list-style-type: none"> % children fed at least minimum recommended number of meals of solid/semi-solid foods yesterday Mean number of meals; mean number of snacks % fed recipes promoted in MCs (past 24 hr, 3+ or 7+ times last wk)
Nutrient content of complementary foods	<ul style="list-style-type: none"> Feed children a variety of foods including nutrient-rich food groups (e.g., animal source foods and vitamin A-rich food) daily. Promoted recipes were formulated to be nutrient-dense 	<ul style="list-style-type: none"> % fed foods from 8 food groups yesterday Mean dietary diversity % fed recipes promoted in MCs (past 24 hr, 3+ or 7+ times last wk)
Use of vitamin-mineral supplements or fortified products for infant and mother	<ul style="list-style-type: none"> Use as needed for child; BF mothers may also need supplements In areas where vitamin A deficiency is prevalent or under-five mortality rate is > 50/1000, give high-dose vitamin A supplements twice annually 	<ul style="list-style-type: none"> % eating fortified WSB (past 24 hr, 3+ or 7+ times last wk); age first received WSB and number of months received % women who received prenatal iron and postnatal vitamin A supplements; % children who received vitamin A in past 6 mo
Feeding during and after illness	Increase fluid intake during illness, including more frequent BF. After illness, give more food, more often	% giving more liquids during diarrhea; % giving the same (or more) food during diarrhea; % giving extra meals after diarrhea

^a Guiding Principles for feeding infants and young children are described in PAHO/WHO 2003 (for breastfed children) and WHO 2005 (for non-breastfed children).

exclusive breastfeeding in the last 24 hours is a useful measure and is widely used in nationally representative surveys such as the Demographic and Health Surveys.

Fed minimum recommended number of meals of solid/semi-solid foods. Respondents reported the number of times during the morning, at noon, afternoon, evening, and night that children were fed solid and/or semi-solid foods. These were summed and the total compared to recommendations. For breastfed children aged 6-8.9 mo, 2-3 meals are recommended, plus snacks “as desired,” and for those aged 9-23.9 mo, 3-4 meals are recommended, plus snacks (PAHO/WHO 2003). These recommendations assume an energy density of at least 0.8 kcals/gram for complementary foods, and also assume that children are fed to gastric capacity at meals. Formative research done in the study areas confirmed that the energy density of commonly prepared complementary foods was adequate (Ruel et al. 2004). All promoted enriched recipes also had an energy density of 0.8 or greater. For non-breastfed children, 4-5 meals are recommended, again with 1-2 snacks as desired (WHO 2005). We constructed a dichotomous variable indicating whether or not each child received at least the minimum number of meals for his/her age and breastfeeding status.

Dietary diversity. A dietary diversity score was calculated, identical to one used at baseline, and summing the following eight food groups: grains (and all gruels made from grain); roots and tubers; legumes; vitamin-A rich fruits and vegetables; other fruits and vegetables; dairy products (including infant formula); flesh foods and eggs (including meat, organ meat, poultry, fish, seafood, and eggs); and nuts. Each food group eaten yesterday provided 1 point, yielding a diversity score ranging from 0-8. For selected nutrient-dense foods, and for recipes promoted in MCs, we also created dichotomous indicators reflecting whether or not the child had received these foods at least 3 times in the previous week, and whether they had received them at least 7 times in the previous week.

IYCF indicator: An IYCF indicator was created using data on breast feeding and key aspects of complementary feeding for children between 6 and 23 months of age. The indicator was modeled after the IYCF indicator developed for use in the Demographic and Health Surveys (Mukuria, Kothari, and Abderrahim 2006), with one adaptation. In our survey, data were not gathered on consumption of foods containing fats and oils, and therefore, this variable was not available for inclusion in the index. Nevertheless, an indicator was created that captured the three main components of the IYCF indicator: whether a child received breast milk or other calcium rich foods if not breast fed, whether the child was fed solid/semi solid foods a minimum number of times per day, and whether the child was fed an age-appropriate minimum number of food groups (dietary diversity). The dietary diversity variable was slightly different from the DHS IYCF indicator; in our IYCF indicator, we separated nuts from legumes (in effect substituting the variable on nut consumption for the variable on fats/oils consumption). Thus, the variable on dietary diversity that was created for the IYCF indicator had seven food groups, exactly the same number as the DHS IYCF indicator, but one of the variables was different between the two.

8.2.4 Other care practices: Preventive care, hygiene practices, and care during illness

Families who participate in WV-Haiti programs are encouraged to care for their children by accessing RP preventive health-care services such as immunizations, and by using good hygiene practices. Parents are encouraged to seek help from the health agent or other medical professional when children are ill, and to use the ORS that is distributed at RPs when children have diarrhea. These caregiving and care-seeking practices are promoted within the larger MCHN program, and thus for all children in both preventive and recuperative areas.

The following indicators were used to measure these other care practices:

1. Percent of children 12 mo and older who were fully immunized;
2. Percent of ill children (last two weeks) for whom medical advice or care was sought, by symptom (fever, respiratory symptoms, diarrhea);
3. Percent of children with diarrhea who were given ORS or homemade sugar-salt solutions.

Child fully immunized. The percent of children fully immunized was constructed from information about individual vaccines received. This information was available from health cards and/or from mother's recall. Information from these two sources is presented separately (with recall data reported only for those who did not have health cards) and is then combined for an overall estimate. Children are considered to be fully immunized if, at ages 12 mo and older, they have received the following vaccines: BCG, 4 polio, 3 DTP, and measles (WHO 2002).

Hygiene scale scores. Spot observations of child and maternal cleanliness, as well as cleanliness of the interior and exterior of the house were used as proxy measures of hygiene practices. Observers assessed the cleanliness of the child's hands, face, clothes, and hair, and rated each on a scale of 1-3. For children who were naked at the time of observation, the cleanliness of their body was assessed instead of cleanliness of clothes. The observer also noted yes/no for whether the child had an "unattended" runny nose. A child cleanliness scale was constructed using these variables to be identical to the scale used at baseline. The possible range for the scale was 4-13.

A similar scale was constructed for maternal cleanliness (range 4-12), including four of the same items: cleanliness of hands, face, clothes, and hair, each scored from 1 to 3. Two scales were constructed for cleanliness of the area around the house (general appearance of the compound, was the compound swept, and were garbage, animal feces, or human feces absent from the compound), and the cleanliness of the interior of the house (general appearance, was the floor swept, were drinking water containers covered, and were dirty clothes absent). These two scales ranged from 1 to 7 (exterior) and 1 to 6 (interior).

8.3 Analysis

The analyses presented in this chapter are meant to describe current practices. As the overall study was designed to compare the preventive and recuperative approaches, statistical tests are reported primarily for these comparisons, consistent with the study design and sampling. Most results are presented for index children (12-41 months of age), but for a few variables that are relevant mainly for younger children (e.g., exclusive breastfeeding during the first 6 months), we include results for the sample of younger siblings. Note, however, that the statistical power for this sample of younger siblings is much lower than for the index children because the sample size of younger siblings is only 341. For data on immunization coverage, care-seeking, and hygiene, baseline information is available only from 18 months onward, so for comparability we present this information for children aged 12-41 months.

In order to account for the survey design, tests of differences in means and proportions were performed in STATA using *xtreg* (continuous variables) or *xtlogit* (dichotomous variables). This allows specification of the paired cluster design and provides the most appropriate standard error for these tests. P-values of less than 0.05 are considered to be significant.

For some topics/variables, we also examined differences between program participants and nonparticipants using two measures of participation: (1) whether the mother had ever participated in the MCs or not; and (2) whether the child was currently in the program and therefore receiving food assistance commodities.²³ These comparisons are presented mainly to provide additional insight about the plausibility of our key results; they are not part of our main analysis, which focuses on an “intent to treat” approach.

Finally, comparisons are also made between breastfed and non-breastfed children for key relevant variables.

Comparisons with baseline results are often presented in our results, but they are meant to be interpreted qualitatively—i.e., they are presented to provide a sense of the initial levels of the different variables assessed in the final survey. No statistical testing is done to compare baseline and final survey results.

8.4 Results

8.4.1 Impact of the program on maternal knowledge

We examined differences between the program groups on a variety of maternal knowledge scales, capturing both overall maternal health and nutrition knowledge, and knowledge on specific groups of topics taught at the Mothers’ Clubs.

Infant and Young Child-feeding Knowledge Test: At baseline, the program communities were not different on any of the knowledge variables. At the final survey, respondents in the preventive community scored higher than those in the recuperative group on knowledge tests

²³ These participation variables are slightly different from the ones used in Chapter 6. This is because we viewed mother’s exposure to MCs as the key program exposure for changes in practices. For current participation, we wanted to assess the relationship between current receipt of food and selected practices.

related to feeding frequency and on the overall knowledge scale; differences were statistically significant, but of small magnitude (Table 8.2).

Notably, results for the three knowledge tests for which data were also available at baseline were markedly higher at the final survey, particularly among respondents who had ever participated in the program (Table 8.3). Within program groups, the differences between respondents who had ever been exposed to the program and those who had not were large and statistically significant. Knowledge scores among those never exposed to the program were close to baseline values, while scores for those who had been exposed were higher. This was true for both program groups, which shows that exposure to the BCC strategy included in both program approaches had a large impact on maternal knowledge.

General Nutrition and Health Knowledge Test of BCC topics: The knowledge scores based on a test of topics taught at the MCs are slightly higher (and statistically significant) among preventive community respondents for overall health and nutrition knowledge and knowledge pertaining to childhood illness. All other scores are almost identical between the two groups and not statistically different.

Table 8.3 shows, as would be expected, that overall and topic-specific knowledge is quite different for mothers who never participated in the program compared to those who had participated. Although this was true for both program groups, differences between participants and nonparticipants were larger in the preventive group; participating mothers from preventive communities had higher overall knowledge scores and higher scores to the breastfeeding and complementary feeding knowledge tests (Table 8.3). Trends for proportion of knowledge questions answered correctly were the same as for the knowledge scores.

Multivariate analyses show a statistically significant interaction between having ever participated in the program and program group (preventive versus recuperative) (Table 8.4). The significant interaction confirms findings from bivariate analyses described above that suggest that a larger effect of program participation on overall knowledge score is found in preventive compared to recuperative communities.

Table 8.2 Maternal nutrition and health knowledge, by program group

Knowledge scores	Baseline		Final survey					
	Mean	SD	Preventive	Recuperative		Overall		
			(n = 748)	(n = 752)		(n = 1,500)		
			Mean	SD	Mean	SD	Mean	SD
<i>Infant and Young Child-feeding Knowledge Test</i>								
Knowledge about appropriate introduction of foods (maximum score: 6)	2.4	1.7	4.9	1.7	4.9	1.7		
Appropriateness of feeding frequency knowledge (maximum score: 6)	5.4	0.9	5.7	0.6	5.6**	0.7		
Feeding knowledge, weighted (maximum score: 6)	4.3	0.7	5.4	0.7	5.3**	0.7		
<i>Overall Nutrition and Health Knowledge Test (BCC topics)</i>								
Overall health and nutrition knowledge (maximum score: 28)	n/a		20.1	2.9	19.8**	2.8	19.9	2.9
Breastfeeding knowledge (maximum score: 8)	n/a		4.4	1.3	4.3	1.2	4.4	1.2
Complementary feeding knowledge (maximum score: 9)	n/a		8.4	1.0	8.3*	1.0	8.3	1.0
Knowledge about child illness (maximum score: 4)	n/a		2.4	0.9	2.3**	0.9	2.3	0.9
Knowledge about general health issues (maximum score: 5)	n/a		3.3	1.0	3.2	1.0	3.2	1.0
Knowledge about malnutrition (maximum score: 2)	n/a		1.7	0.5	1.7	0.5	1.7	0.5
<i>Percentage of answers correct on each of the scores</i>								
Percentage of answers correct on overall knowledge test	n/a		71.7	10.6	70.6*	9.9	71.2	10.2
Percentage of answers correct on breastfeeding questions	n/a		55.0	15.8	53.8	14.6	54.4	15.2
Percentage of answers correct on complementary feeding	n/a		92.9	10.6	92.0*	11.3	92.5	11.0
Percentage of answers correct on child illness	n/a		65.2	20.6	63.7	20.8	64.5	20.8
Percentage of answers correct on general health	n/a		58.9	21.7	56.6**	23.0	57.8	22.4
Percentage of answers correct on malnutrition	n/a		85.1	25.4	87.0	23.9	86.1	24.7

Note: * 0.05 < p < 0.1; ** p < 0.05; *** p < 0.01.

Table 8.3 Maternal nutrition and health knowledge, by program group and participation (ever participated)

	(n = 1,524)		Preventive				Recuperative			
			(n = 126)		(n = 622)		(n = 126)		(n = 626)	
			Nonparticipant		Participant		Nonparticipant		Participant	
Nutrition and health knowledge	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
<i>Infant and Young Child-feeding Knowledge Test</i>										
Knowledge about appropriate introduction of foods (maximum score: 6)	2.4	1.7	3.4	2.3	5.2 ^a	1.4	4.1	2.2	5.1 ^a	1.5
Appropriateness of feeding frequency knowledge (maximum score: 6)	5.4	0.9	5.6	0.7	5.7 ^a	0.6	5.5	0.8	5.7 ^a	0.6
Feeding knowledge, weighted (maximum score: 6)	4.3	0.7	4.7	0.8	5.5 ^a	0.6	4.9	0.9	5.4 ^a	0.6
<i>Overall Nutrition and Health Knowledge Test (BCC topics)</i>										
Overall health and nutrition knowledge (maximum score: 28)	n/a		16.6	3.5	20.8 ^a	2.2	17.9	3.4	20.2 ^a	2.5
Breastfeeding knowledge (maximum score: 8)	n/a		2.7	1.5	4.7 ^a	0.9	3.3	1.4	4.5 ^a	1.0
Complementary feeding knowledge (maximum score: 9)	n/a		7.5	1.3	8.5 ^a	0.8	7.9	1.3	8.4 ^a	0.9
Knowledge about child illness (maximum score: 4)	n/a		1.9	0.9	2.4	0.8	2.1	1.0	2.3	0.9
Knowledge about general health issues (maximum score: 5)	n/a		2.9	1.2	3.3	1.0	3.0	1.1	3.2	1.0
Knowledge about malnutrition (maximum score: 2)	n/a		1.5	0.7	1.7	0.5	1.6	0.6	1.8	0.4
<i>Percentage of answers correct on each of the scores</i>										
Percentage of answers correct on overall knowledge test	n/a		59.2	12.6	74.3 ^a	8.0	63.9	12.2	72.0 ^a	8.8
Percentage of answers correct on breastfeeding questions	n/a		34.2	18.4	59.3 ^a	11.2	41.6	18.0	56.4 ^a	12.4
Percentage of answers correct on complementary feeding	n/a		83.7	14.9	94.8 ^a	8.4	87.4	14.6	92.9 ^a	10.3
Percentage of answers correct on child illness	n/a		57.9	24.0	66.7	19.6	60.0	21.3	64.5	20.6
Percentage of answers correct on general health	n/a		48.0	23.4	61.2	20.7	51.7	25.3	57.6	22.4
Percentage of answers correct on malnutrition	n/a		74.2	33.5	87.3	22.9	81.5	31.5	88.2	21.9

^a Difference between participants and nonparticipants (within program group) is significant at $p < 0.05$ (random effects regression analysis).

Baseline

Table 8.4 Program impact on maternal nutrition and health knowledge (regression analysis)

	(1) Bivariate model Knowledge	(2) Main effects, adjusted Knowledge	(3) Main effects, ever participated Knowledge	(4) Interaction model, ever participated Knowledge
Program	-0.31* (0.15)	-0.32* (0.16)	-0.34* (0.14)	1.06** (0.36)
Respondent's education (years)		0.12** (0.03)	0.12** (0.03)	0.12** (0.03)
Partner's education (years)		0.03 (0.03)	0.04 (0.02)	0.04 (0.02)
Ever participated			3.31** (0.20)	4.13** (0.28)
Ever participated x Program				-1.66** (0.40)
Constant	20.09** (0.15)	19.86** (0.17)	17.07** (0.24)	16.37** (0.29)
Observations	1,500	1,250	1,250	1,250
Number of pairs	10			

Notes: Standard errors in parentheses. * significant at $p < 0.05$; ** significant at $p < 0.01$.

In summary, results of the impact of the two program approaches on maternal knowledge show that:

For Infant and Young Child-feeding Knowledge Test:

- ❖ There were no differences between program communities in maternal knowledge at baseline.
- ❖ At the final survey, mothers from the preventive program communities had significantly higher scores on the overall feeding knowledge scale and the knowledge about feeding frequency tests than mothers from the recuperative program communities; all differences were of small magnitude, however.
- ❖ Both preventive and recuperative groups showed markedly greater nutrition knowledge at final survey compared to baseline.
- ❖ Attained knowledge at final survey was significantly higher among women who had participated in the program at some point in the previous 3 years, compared to those who had never participated.

For the General Nutrition and Health Knowledge Test – based on MC and RP topics

- ❖ Maternal knowledge about health and nutrition topics taught at the MCs was good, with respondents getting about 71% correct answers on the general nutrition and health knowledge test.

- ❖ Respondent knowledge about BCC topics was statistically significantly better in preventive communities compared to recuperative communities in overall scale as well as subscales related to breastfeeding and complementary feeding, but differences were of small magnitude.
- ❖ More meaningful and significant differences in knowledge were seen between respondents ever exposed to the program versus those never exposed to the program. Differences between exposed and nonparticipant respondents were greater in the preventive communities than in the recuperative communities.

8.4.2 Trial and adoption of key recommended practices

Figures 8.1a through 8.1g present results related to awareness, trial, and adoption of key recommended practices; Table 8.5 presents comparisons between preventive and recuperative communities in trial of recommended practices and the reasons reported for non-trial of the practices. Data are presented separately for women who had ever participated versus those who had never participated because data collection instruments were slightly different for these two groups of women. Although statistical significance is not depicted in Figures 8.1a through 8.1g, significant results are discussed below, and Annex 8.4 presents the same results in tabular format, with statistical significance indicated in the table.

Differences between program groups among participants in awareness, trial, and adoption were statistically significant for almost all practices. Notably, for adoption rates, the differences between participants in preventive and recuperative groups were seen mainly for complementary feeding-related practices, with no significant differences in adoption of the use of expressed breast milk. This conforms to expectations because pregnant and lactating women were targeted in both program approaches, and participation rates were remarkably similar for pregnant and lactating women in the two program groups.

Overall, the magnitude of differences in awareness, trial, and/or adoption between the preventive and the recuperative program are of a smaller magnitude than the differences between respondents who had ever participated versus those who had never participated in each of the program groups (also statistically significant for almost all of the practices). The overall pattern suggests that in both program groups, awareness, trial, and adoption are substantially higher among respondents who had ever participated in the program compared to those who had never participated.

Practices that had been tried by at least half of the participant mothers were the use of expressed breast milk, adding an egg to the child's gruel, feeding enriched gruel in the evening, preparing special recipes such as mashed plantain with pumpkin, and feeding an extra meal a day after an illness (Table 8.5). The practices that had been tried by only about one-third of participant respondents were adding breast milk to gruel, and preparing gruel with added beans and nuts.

Table 8.5 Differences in trial of recommended practices between preventive and recuperative groups and reasons for non-trial of practices, by program exposure

Recommended practice	Ever participated in MC			Never participated in MC		
	Recup.		Reasons for non-trial (among those who did not try the practice)	Recup.		Reasons for non-trial (among those who did not try the practice)
	(N = 622)	(N = 622)		(N = 122)	(N = 130)	
	<i>Ever tried</i>			<i>Ever tried</i>		
		%		%	%	
Leaving expressed breast milk behind when going out	65.1 ^{a b}	59.3 ^a	❖ Don't leave home for long (~75%)	22.2	24.6	❖ Don't know how to do it (~30%) ❖ Don't leave home for long (~50%)
Adding breast milk to gruel	31.7 ^{a b}	24.0 ^a	❖ Don't know how to do it (~33%) ❖ Afraid child won't eat it (~10%) ❖ It's disgusting (~33%)	3.2	4.0	❖ Don't know how to do it (~25%) ❖ Afraid child won't eat it (~15%) ❖ It's disgusting (~45%)
Adding an egg to child's gruel	72.5 ^b	53.7 ^a	❖ Don't know how to do it (~23%) ❖ No ingredients (~25%) ❖ Expensive ingredients (~20%) ❖ Negligence/don't care (~25%)	76.2	70.6	❖ Don't know how to do it (~25%) ❖ No ingredients (~28%) ❖ Expensive ingredients (~25%) ❖ Negligence/don't care (~15%)
Prev.						
Preparing gruel with beans and nuts	45.5 ^{a b}	35.9 ^a	❖ Don't know how to do it (~35%) ❖ No ingredients (~235%) ❖ Expensive ingredients (~20%) ❖ Negligence/don't care (~10%)	7.9	4.0	❖ Don't know how to do it (~28%) ❖ No ingredients (~23%) ❖ Expensive ingredients (~25%) ❖ Negligence/don't care (~15%)
%						
Feeding enriched gruel in the evening	76.7 ^{a b}	67.1 ^a	❖ No ingredients (~40%) ❖ Expensive ingredients (~10%) ❖ Negligence/don't care (~15%)	14.3	15.9	❖ No ingredients (~50%) ❖ Expensive ingredients (~12%) ❖ Negligence/don't care (~12%)
Preparing mashed plantain with pumpkin	79.7 ^{a b}	67.3 ^a	❖ Don't know how to do it (~15%) ❖ Pumpkin difficult to find (~47%) ❖ Pumpkin is expensive (~13%)	36.5	29.4	❖ Don't know how to do it (~12%) ❖ Pumpkin difficult to find (~50%) ❖ Pumpkin is expensive (~12%)
Feeding an extra meal after illness	90.7 ^{a b}	84.0	❖ Child has no appetite (~35%) ❖ Not enough food at home (~35%) ❖ Not enough time (~10%)	77.8	81.4	❖ Child has no appetite (~35%) ❖ Not enough food at home (~35%) ❖ Not enough time (~10%)

^a Difference between participants and nonparticipants (within program group) is significant at $p < 0.05$ (random effects regression analysis).

^b Difference between program groups is significant at $p < 0.05$ for participants (random effects regression analysis).

Figure 8.1a From awareness to adoption: Leaving expressed breast milk behind when going out




Figure 8.1b From awareness to adoption: Adding breast milk to gruel




Figure 8.1c From awareness to adoption: Adding an egg to child's gruel




Figure 8.1d From awareness to adoption: Preparing gruel with beans and nuts




Figure 8.1e From awareness to adoption: Feeding enriched gruel in the evening




Figure 8.1f From awareness to adoption: Preparing mashed plantain with pumpkin





Figure 8.1g From awareness to adoption: Feeding an extra meal after illness



As expected, trial of recommended practices was markedly lower among nonparticipating mothers. Only two of the recommended practices were reported by more than half of nonparticipant mothers: the addition of an egg to the child's gruel and feeding an extra meal after an illness. Between 25% and 37% of nonparticipant mothers had tried leaving expressed milk behind for the child and preparing mashed plantain with pumpkin, but for the remaining practices, the trial rates were extremely low. This was likely driven - at least partly - by low awareness rates, which was observed among nonparticipants for most practices, except leaving expressed breast milk when mother is absent and feeding an extra meal after recovery from an illness.

Patterns of adoption (defined as engaging in the recommended practice at least 3 or more times) followed the patterns of trial. On average, 2.5 practices had been adopted by participants in the preventive group, compared with 2.1 practices in the recuperative group and less than 1 practice among nonparticipants in the preventive group and 0.5 among nonparticipants in the recuperative group ($p < 0.05$ for all differences).

Figures 8.1a to 8.1g (and Annex 8.4) present results that show the drop-off from awareness to trial and adoption, by exposure to the BCC program and by program approach. Results are shown for seven key practices promoted by the program, and demonstrate that depending on the practice, exposure to the program can either impact the translation from awareness to trial (e.g., for using expressed breast milk), or have a direct influence on awareness itself, as well as on subsequent trial and adoption.

For example, the use of enriched gruels in the evening was a completely new practice recommended by the program. In this case it is apparent that exposure to the program led to greater awareness of the practice among participants compared to nonparticipants; for a substantial proportion of participants, this led in turn to trial and adoption. On the other hand, for a practice where the difference in awareness between exposed and nonparticipant groups was not as large, e.g., the use of expressed breast milk, participation in the program appears to better enable the translation of knowledge into action among those exposed to it. Interestingly, nonparticipants were more likely to report *trial* than *awareness* of two practices (addition of an egg to the child's gruel and feeding extra meals after an illness (Figures 8.1c and 8.1g)). The reasons for this are not immediately clear.

Reasons for non-trial of practices: As demonstrated in the qualitative study done as part of the operations research in 2004 (Menon et al. 2005), the translation of awareness into trial of a recommended practice is influenced by a variety of social, cultural, and economic factors. Table 8.5 presents findings on the reasons reported for *not* trying the recommended practices. As was revealed in our previous qualitative work, the reasons for non-trial vary among practices. Since the reasons for non-adoption are largely the same as reasons for non-trial, we do not present those results separately.

For practices that related to the use of expressed breast milk, the major reason for not trying the practice was that 75% of participant and 50% of nonparticipant mothers who had not tried the practice reported not leaving the house for long periods of time. Among the nonparticipant mothers, lack of knowledge was also mentioned as a barrier among 30% of those women who did not try the practice. Perceptions that the practice of using expressed breast milk was unclean were a deterrent to trial of adding breast milk to gruel among over a third of participant mothers, and 45% of nonparticipant mothers who had not tried the practice. Very few nonparticipant mothers had ever tried adding breast milk to gruel.

For those practices that required the use of resources to purchase nutrient-rich foods, it is apparent that economic conditions and access to ingredients were considerable barriers to moving from awareness to trial. For example, mothers reported not trying recipes that encouraged them to add an egg to the child's gruel or to use pumpkin along with mashed bananas because they did not have the ingredients at home, or because the ingredients were expensive. Similarly, they reported not feeding the child enriched gruels in the evening or giving an extra meal after the child recovered from an illness because of inadequate resources at home. In addition to resource constraints, poor child appetite was mentioned as a reason for not trying to feed an extra meal per day to the child.

An interesting reason for the lack of trial of recommended practices is "negligence," which is reported by about 10% of women who did not try recommended practices. The word "negligence" is translated literally from the Creole word and since it is a self-reported reason by respondents, it is not clear if it should be taken in the literal sense of the word in English. Without a deeper understanding of the meaning of this word in Haitian culture, we interpret it to mean that these women who reported "negligence" as a reason for not trying a practice simply did not pay attention to the recommendations, or were not motivated to try them at home. Further examination of the characteristics of women who report "negligence" as a reason for not trying practices, combined with future qualitative research, could help explain this construct.

Lastly, it should be noted that there were no differences in reasons for not trying recommended practices between those who had ever participated and those who had never participated in the program. A difference was seen only in the case of using expressed breast milk, where about a third of nonparticipant mothers said they had not tried the practice because they did not know how to do it.

In summary, results of the impact of the two program approaches on trial and adoption of key practices show that:

- ❖ For most key practices, respondents in preventive program areas were more likely to report awareness, trial, and adoption than were respondents in recuperative areas. In most cases, however, differences were of relatively small magnitude.
- ❖ Differences between those exposed to the program and those not exposed (in each area) were much larger than differences between program areas.
- ❖ Reasons for non-trial and non-adoption of practices were practice-dependant and ranged from economic and sociocultural to child-focused reasons.

8.4.3 Impact of intervention on infant and young child-feeding and care practices

Child-feeding and care practices were measured by maternal recall in the final survey. One important caveat of this approach to assessing the impact of a BCC program on practices is the possibility that mothers systematically report what they have learned, which results in an overestimate of the true impact of the program on behavior change. Although this type of error is inevitable, the main thrust of our study is *not* on assessing change in practices since baseline, but rather on comparing the impact of two different program approaches on outcomes at the end of the study.

As in previous chapters, the presentation of results in this section emphasizes differences between preventive and recuperative program communities. While mention is made of baseline results and “changes” since baseline, differences between baseline and final survey should be interpreted as *qualitative* information - and are not tested for statistical significance.

Early infant feeding practices

At baseline, there were no differences between program communities in maternal report of early feeding practices (early initiation of breastfeeding; exclusive breastfeeding on the first day; using colostrum). These recommended practices, however, were reported by a relatively small percentage of mothers—less than half of the mothers reported early initiation of breastfeeding and exclusive breastfeeding on the first day, and approximately two-thirds reported giving colostrum to the newborn infant (Table 8.6 and Figure 8.2). In the final survey, the percentage of mothers who reported optimal early infant feeding practices was much higher than at baseline in both the preventive and recuperative program communities, and no differences between the program communities were found for early initiation of BF (65%) and exclusive BF on the first day (90%). There was a small difference in the proportion reporting that they gave colostrum, with more mothers in the recuperative areas reporting this practice compared to preventive areas (88% versus 84%; $P < .05$).


Table 8.6 Child-feeding and care practices, by program group and time of survey

Feeding/care practice	Baseline	Final Survey		
		Preventive	Recuperative	All children
<i>Early child-feeding practice (index child)</i>	(n = 1,602)	(n = 748)	(n = 752)	(n = 1,500)
- Initiated breastfeeding within 1 hour	36.4	64.6	65.8	65.2
- Only breast milk on first day	47.1	90.0	90.4	90.2
- Gave colostrum	63.9	84.4*	88.3*	86.3
<i>Early child-feeding practices (younger siblings 0-11 mo)</i>	(n = 431)	(n = 159)	(n = 185)	(n = 344)
- Breastfed within 1 hour	48.2	84.1	81.9	83
- Only breast milk on first day	55.8	98.1	93	95.3
- Gave colostrum	73.3	95	94.1	94.5
<i>Exclusive breastfeeding in past 24 hour (if < 6 mo)</i>	(n = 255)	(n = 100)	(n = 129)	(n = 229)
	42.0	93	91.5	92.1
<i>Child age when liquids foods were introduced (index child)</i>				
- % who started liquid at 6-8.9 mo	22.2	78.7	78.9	78.8
- % who started semi-solid food at 6-8.9 mo	30.9	83.4	82.4	82.9
- % who started solid food at 6-8.9 mo	42.5	67.9	69.1	68.5
- % who started meat at 6-8.9 mo	15.9	65.9	63.4	64.7
- % who started eggs at 6-8.9 mo	38.1	73.4	60.3	71.9
<i>Maintenance of breastfeeding (index child)</i>				
- % still breastfeeding at 12-17 mo	87.2	89.8	89.8	89.8
- % still breastfeeding at 18-23 mo	36.2	47.1	44.9	46.1
- % still breastfeeding at 24-42	5.4	6.1	4.9	5.5
<i>Responsive Feeding (index child)</i>				
- Child normally feeds himself (12-41 mo)	83.9	62.0*	68.1*	65.0
- 12-23 mo	57.3	34.9	39.5	37.0
- 24-42 mo	93.5	85.9	87.6	86.8
- Mothers who mention 2+ positive strategies	23.6	64.9*	59.7*	62.3
- Caress him/her	53	55.1	66.5	60.7
- Play with him/her	2.2	25.1	12.4	18.8
- Offer other foods	6.5	4.7	4.5	4.6
- Mothers who mention negative strategies				
- Force him/her	35.2	14.1	14.8	14.4
<i>Safe preparation and storage of complementary foods - use of baby bottles</i>				
- % mothers who used baby bottles	39.9	17.5	16.2	16.9
Breastfed children				
- 12-17 months				
% fed minimum no. of meals	38.2	60.8	54.2	57.6
Mean no. of meals	2.4	2.8	2.6	2.7
Mean no. of snacks	1.8	2.1	1.9	2.0
- 18-23 months				
% fed minimum no. of meals	53.9	66.7	56.6	62.4
Mean no. of meals	2.6	2.9	2.6	2.8
Mean no. of snacks	2.2	2.2	2.3	2.2
- 24-42 months				
% fed minimum no. of meals	--	--	--	--
Mean no. of meals	2.6	2.7	2.8	2.7
Mean no. of snacks	2.0	2.8	1.7	2.3
Non-breastfed children				
	(n = 1,373)	(n = 475)	(n = 509)	(n = 984)
- 12-17 mo				
% fed minimum no. of meals	0.0 (n=17)	14.3	10.5	12.5
Mean no. of meals	2.7	2.9	2.7	2.8
Mean no. of snacks	2.0	2.4	2.6	2.5
- 18-23 mo				
% fed minimum no. of meals	14.1	17.3	12.1	15.0
Mean no. of meals	2.7	2.8	2.8	2.8
Mean no. of snacks	2.5	2.4	2.2	2.3

(continued)

Feeding/care practice	Baseline	Final Survey		
		Preventive	Recuperative	All children
- 24 mo and older				
% fed minimum no. of meals	--	--	--	--
Mean no. of meals	2.6	2.7	2.7	2.7
Mean no. of snacks	2.3	2.3	2.1	2.2
<i>Feeding patterns - evening meal (index child)</i>				
- Will wake sleeping child for evening meal	20.6	18.6	19.7	19.1
<i>Percent of children who consumed selected food groups in previous 24 hrs (index child)</i>				
- Cereals	96.3	98.0	97.1	97.5
- Roots, tubers, starchy vegetables	35.0	33.2	37.1	35.2
- Legumes	75.8	87.3	84.2	85.7
- Vitamin A-rich vegetables	95.1	72.2*	67.3*	69.7
- All other fruits and vegetables	49.9	38.4***	30.3***	34.3
- Milk and formula	18.9	27.2**	21.1**	24.1
- Meat, fish and egg	88.3	62.5***	53.6***	58.1
- Nuts	48.3	23.0	24.1	23.5
<i>Mean dietary diversity (index child)</i>				
- 12-17 mo	4.8	4.4*	4.0*	4.2
- 18-23 mo	5.0	4.5	4.2	4.3
- 24-35 mo	5.1	4.4	4.2	4.3
- 36-41 mo	5.1	4.4	4.1	4.2
- All index children: 12-41 mo	5.1	4.3***	4.1***	4.3
<i>Percent of children who consumed selected animal source foods (index child)</i>				
- Eggs	42.6	20.7*	16.2*	18.5
- Chicken	25.1	7.8	5.6	6.7
- Fish and seafood	65.8	34.7	30.2	32.4
- Beef and pork	64.9	28.4***	21.0***	24.7
- Heart and liver	18.3	5.0	4.1	4.5
<i>Frequency of consumption of nutrient-rich foods (index child)</i>				
- Food/groups consumed 3 or more times in the last 7 days				
Eggs	8.4	14.3	12.1	13.2
Flesh food	34.8	72.6***	60.1***	66.3
Vitamin A-rich orange/red fruits/vegetables	67.2	47.3	46.8	47.0
<i>IYCF indicator</i>				
- Score	Not available	2.12*	1.99*	2.1
- % meeting minimum recommendations for all 3 practices	Not available	43.0*	36.3*	39.8
<i>Consumption of WSB (only among index children currently in program)</i>		(n = 280)	(n = 106)	(n = 320)
- Gruel made with WSB in past 24 hr	n/a	53.0	48.1	51.7
- Gruel made with WSB 3+ times in past 7 d	n/a	58.6	58.5	58.5
- Gruel made with WSB 7+ times in past 7d	n/a	5.4	6.6	5.7
- Other food made with WSB in past 24 h	n/a	16.7	17.9	17.1
- Other food made with WSB 3+ times in past 7 d	n/a	16.0	17.9	16.5
- Other food made with WSB 7+ times in past 7d	n/a	0.7	0.9	0.8
<i>Vitamin and Mineral Supplements (index child)</i>				
- % children receiving vitamin A capsule in last 6 mo	30.2	51.6*	45.7*	48.7
- % women who received postpartum vitamin A	7.84	63.5***	54.1***	58.8
- % women who received prenatal iron supplements	65.3	82.3	84.0	83.2
<i>Feeding during and after diarrhea (index child)</i>				
- % who gave more liquid	52.5	50.0	44.8	47.4
- % who gave more semi-solids/solids	6.4	30.6	28.2	29.4
- % who gave extra meal after recovery	n/a	66.7*	55.7*	61.4

Figure 8.2 Percent of mothers who reported optimal early feeding practices during first day of life, by program group and baseline and final survey—Index child



* Statistically significant at $p < 0.05$.


Since many of our index children were born before their mother was exposed to the program BCC, we also look at whether early infant practices had improved for their younger sibling (see Table 8.6 and Figure 8.3). We found no differences between program groups, but younger siblings from the final survey were much more likely to have received optimal early infant practices than those from the baseline survey. As was expected, younger siblings were also more likely than index children to have been breastfed within the first hour (83% versus 65% of index children), and to have been given colostrum (95% versus 86%). These findings suggest that improved maternal knowledge may indeed have translated into improved practices related to early infant feeding practices.

Exclusive breastfeeding up to 6 months of age is a key practice and one that received great emphasis in the MC's learning sessions. This practice was assessed only among younger siblings, because our index children were between 12-41 months of age. At baseline, 42% of mothers with infants younger than 6 months of age reported exclusively breastfeeding the previous day, with no difference between program areas. At the time of the final survey, 92% of women with an infant younger than 6 months reported exclusively breastfeeding in the previous 24 hours, again with no differences between program areas (Table 8.6).

While these results, as noted, could reflect women's knowledge of the "desirable" response, this is unlikely because the categorization of children into the "exclusive breastfeeding" category required that mothers respond consistently to several feeding practices questions (see variable creation in methods section 8.2). For instance, mothers were asked how many times the infant had been fed solids/semi-solids the previous day (at different times during

the day); they were also asked about a long list of liquids and foods that their infant might have consumed yesterday, and about the timing of introduction of various foods in their infant's diet. Responses were consistent across all these questions.

Figure 8.3 Percent of mothers who reported optimal early feeding practices during first day of life, by program group and baseline and final survey—Younger siblings



* Difference between preventive and recuperative at final survey is statistically significant ($p < 0.05$).

We also explored whether the proportion of infants reported to be exclusively breastfed fell off sharply in the first 6 mo, as had been the case at baseline. In the final survey, this was not the case (see Figure 8.4). Median duration of breastfeeding, as assessed by Kaplan-Meier survival tables,²⁴ was 5.9 months and did not differ between program areas.


Timing of introduction of liquids, and semi-solid and solid foods

Mothers were asked to recall when they had first given their child various liquids and foods. Specifically, they were asked to recall at what age they had first given their infant liquids other than breast milk, semi-solid foods, solid foods, meat products, and eggs. A large proportion of women reported introducing liquids and semi-solid foods at the recommended age of 6-8.9 months (e.g., 79% for liquids, 83% for semi-solid foods; 69% for solid foods), with no difference between the preventive and recuperative program areas. Compared to baseline, a much lower percentage of mothers reported introducing liquids and semi-solid foods earlier than at 6 months of age (21% for liquids and 17% for semi-solid foods); at baseline 81% reported giving liquids early and 73% reported giving semi-solids early (Table 8.6; Figure 8.5). For some

²⁴ Survival tables allow analysis of “censored” data, where all observations can be incorporated into the analysis whether or not the “event” (in this case, move from exclusive to nonexclusive breastfeeding) has occurred.

infants, meat (29%) and eggs (21%) were introduced late (after 9 mo), with fewer children (7%) given these foods too early. The percentages of mothers who introduced these nutrient-rich foods at the recommended age (6-8.9 months) were much higher in the final survey compared to baseline (Table 8.6).

Figure 8.4 Comparing baseline and final sample exclusive breastfeeding in past 24 hours for infants < 6 months: Kaplan-Meier survival functions (preventive/recuperative groups combined)




Maintenance of breastfeeding

Table 8.6 shows the proportion of index children still breastfed, by age group. The pattern is similar in preventive and recuperative areas, and is also quite similar to the pattern seen at baseline, with most children being breastfed up to approximately 17 months, followed by a sharp decline thereafter.

Responsive feeding

Responsive feeding is a complex behavior and one that is difficult to capture in simple surveys. In the survey, respondents were asked if their child ate by him/herself, or if s/he was fed. The appropriateness of a child feeding him/herself is clearly related to age: as children develop into their second and third years, it is expected that children will gain experience with feeding themselves, but supervision by an adult caregiver is important to ensure that the child eats adequate amounts of food. Mothers were also asked if they would take any action when

Figure 8.5 Maternal recall of timing of introduction of liquids and foods, at baseline and final survey (preventive/recuperative groups combined)



the child refused to eat. If they responded “yes,” they were prompted for three actions or strategies they might use when the child refused. Women in recuperative communities were slightly more likely to report that their child fed him/herself (68% versus 62% in preventive areas) (Table 8.6). Conversely, women in preventive communities were slightly more likely to be able to name two or more positive actions that they took when the child refused to eat (65% versus 60% in recuperative areas). There were, however no significant differences between program communities in the percentage of mothers who reported using any given positive or negative strategy when the child refused to eat.

Compared to baseline, final survey results suggest positive changes in recommended practices related to responsive feeding. The proportion of children in both program communities who were reported to feed themselves was lower in the final survey compared to baseline. The proportion of women who reported using two or more positive strategies (caressing, playing, offering different food) increased from 24% to 62%, while the proportion reporting using one or more negative strategy decreased markedly (forcing the child decreased from 35% at baseline to 14% at the final survey).

Safe preparation and storage of complementary foods - Use of baby bottles


Our survey did not include data on the safe preparation and storage of complementary foods. However, we did gather data on the use of baby bottles. We found no difference between preventive and recuperative communities in the use of baby bottles at the time of the final survey (about 17% in both groups). This was substantially lower than the proportion of mothers who reported using baby bottles at baseline (40%).

Feeding frequency

Infants and young children need to be fed frequently throughout the day, due to their small gastric capacity and their high energy and nutrient requirements. Non-breastfed children must be fed more often than breastfed children because they do not benefit from the energy and nutrients contained in breast milk. Current recommendations for feeding frequency are that breastfed children 6-8 months old should receive 2-3 meals of complementary foods/day and 9-23 months old, 3-4 meals a day (PAHO/WHO 2003). Non-breastfed children should receive one additional meal a day, and thus the recommendation is 3-4 meals/day for 6-8-month-old children and 4-5 meals a day for 9-23-month-old children. There are no specific recommendations for children 24 months and older, who are expected to consume the family diet; results for this age group are more difficult to interpret.

At baseline, frequency of feeding was similar between the two program groups for breastfed children 12-41 months old (Table 8.6). The same was true at the final survey. Although children in the preventive program communities were slightly more likely to have met the minimum recommended number of meals between 12-23 months of age and had consumed, on average, a slightly higher number of meals, the differences between program groups were not statistically significant. Compared with baseline, a much higher proportion of 12-17-month-old breastfed children in both program groups had consumed the minimum recommended number of meals (58% compared to 38% at baseline) (Figure 8.6) and the average number of meals had increased by 0.3 and 0.2 among 12-17 and 18-23-month-old children, respectively.

Figure 8.6 Percent of breastfed children having received complementary foods minimum recommended times yesterday, by age, program group, and time of survey




* Differences between preventive/recuperative statistically significant ($p < 0.05$).

Analysis of younger siblings (children 6-12 months of age) showed similar findings: i.e., no significant difference between program groups at final survey and a generally larger proportion of children having been fed the minimum recommended number of times compared to baseline (not shown).

Non-breastfed children were much less likely than breastfed children to have received the minimum recommended number of meals (Figure 8.7). Overall, 86% of non-breastfed children aged 12-23 months were reported to have had solid/semi-solid foods²⁵ fewer than the recommended 4 times the previous day, with no change since baseline (87%), and no difference between program areas. The number of meals at different ages also increased only slightly between baseline and the final survey among non-breastfed children (average increase of 0.1 meal). The results should be interpreted with caution because the data on number of meals does not include information on consumption of milks other than breast milk. However, only 23% of non-breastfed children were reported to have had dairy products the previous day. An additional caveat is the contribution of snacks to daily energy intake, which could be more important for non-breastfed than breastfed children. This information is not available from our survey data, but data on the number of snacks suggest a slightly higher number of snacks among non-breastfed 12-23-month-old children compared to breastfed children. The learning sessions did not emphasize different feeding frequencies for non-breastfed children, and this could have influenced the results for this subset of children.

²⁵ About 28 percent of children between 12 and 23 months of age were not being breastfed any longer.

Figure 8.7 Percent of non-breastfed index children fed complementary foods a minimum recommended times, by age, program group, and time of survey



Note: Percent of non-breastfed children fed minimum number of meals was 0% at baseline for 12-17-month-old children.

* Differences between preventive/recuperative statistically significant ($p < 0.05$).


In the formative research conducted early in this project (Menon et al. 2002b), it was noted that young children were rarely fed in the evening, for several reasons. Both lack of food, and cultural beliefs that late feeds could lead to indigestion were cited as reasons for this. The importance of giving children a meal late in the day was one program message. However, meal patterns remained similar to baseline in that respect (Table 8.6), with few children being fed in the evening. Only approximately one-fifth of mothers at baseline and at the final survey reported waking a sleeping child to feed an evening meal, without any difference between program groups. Information from a 24-hour recall suggests an even lower percentage - 10% - of young children being fed an evening meal (not shown), again, without any differences between program groups.

Nutrient content of complementary foods: Dietary quality and diversity

A diverse diet that includes micronutrient-rich food groups can help ensure adequate intakes of all nutrients for infants and young children. There were no differences at baseline between the two program groups in the proportion of children who had consumed different food groups, or in mean dietary diversity. Results of the final survey (Table 8.6; Figure 8.8) show that children in the preventive areas were more likely than those in recuperative areas to have eaten vitamin A-rich fruits and vegetables, other fruits and vegetables, dairy, and other animal-source foods (meat, fish, and eggs) (all had $p < 0.05$) on the previous day. Consumption of cereals, roots/tubers/starchy vegetables, and nuts did not differ between program communities at final

survey. Mean dietary diversity was consistently greater among preventive compared to recuperative program children, but differences were statistically significant only for the 12-17-month age group and for all ages grouped combined.

Figure 8.8 Percent of index children who consumed selected food groups in previous 24 hours, by program group, and baseline and final survey



* Differences between preventive/recuperative statistically significant ($p < 0.05$).

Overall dietary diversity and the proportion of children consuming fruit and vegetables and meat/fish/eggs were lower at the final survey compared to baseline (Table 8.6; Figure 8.8). This may reflect the deteriorating economic situation, increased food prices and continued extreme food insecurity in the region during the survey period (see Chapter 9 on food security).

Because eggs and flesh foods are high in a number of bioavailable micronutrients, we also looked more closely at these food groups (Table 8.6). In the final survey, index children in preventive areas were more likely to have had eggs, and beef or pork than children from recuperative program areas. Children in the preventive program communities were also more likely to have consumed flesh foods 3 times or more in the previous 7 days (73%) than were children in the recuperative group (60%).

Finally, the WV-Haiti program had developed and promoted a series of recipes based on donated food commodities (wheat-soy blend (WSB)) complemented with nutrient-rich locally available foods. Among *current* program beneficiaries²⁶ at the time of the final survey,

²⁶ This analysis is done only among children who are current program beneficiaries because children not currently in the program do not receive WSB.

approximately half of the children had consumed a WSB-gruel on the day prior to the survey and 17% had consumed a complementary food preparation containing WSB and other foods with no difference between the two program areas (Table 8.6). We also assessed frequency of consumption of these foods over the past week; 59% of mothers reported having given the child a WSB-gruel and 17% gave another preparation containing WSB three or more times in the previous week, again, with no difference among participants from the preventive versus the recuperative areas.

Energy and nutrient-dense foods are particularly important for non-breastfed children because they do not benefit from all the nutrients contained in breast milk. In spite of their great need for nutrient-dense foods, we found that non-breastfed index children were *not* more likely than breastfed children to have had any of the animal source foods in the previous 24 hours, nor were they more likely to have had these foods more frequently over the past week (results not shown).

IYCF indicator

The score on the IYCF indicator was slightly higher and statistically significantly different in the preventive compared to recuperative approach (2.13 versus 1.99) (see Table 8.6). The proportion of children between 6 and 23 months of age for whom the 3 practices met the minimum recommendations was also higher in the preventive group (43% versus 36%, $p < 0.05$ using random effects logit regression models to compare program groups).

Food consistency

As noted above, food consistency was not directly observed or measured. However, several indirect indicators related to food consistency were measured. Results related to recipes promoted in MCs are presented above: in addition to being nutrient-dense, these recipes were designed to help mothers prepare foods of appropriate consistency for their child. As reported above, a large proportion of children were fed these recipes on a regular basis while participating in the program, and there were no differences between the groups.

Second, infants should be offered pureed, mashed, and semi-solid food starting at 6 months of age. The percentage of children first offered semi-solid food at 6-6.9 months serves as an indicator of both appropriate timing *and* consistency of food. As reported above, overall, 83% of mothers of index children reported first giving semi-solid foods to index children at 6-6.9 months, without any difference between the two program groups. However, as noted previously, it is unclear whether these results truly represent improved practices or whether they reflect major improvements in maternal knowledge.

Use of fortified foods and vitamin-mineral supplements

Fortified foods and vitamin-mineral supplements can help ensure nutrient adequacy, especially where access to and intakes of animal-source foods are limited. In our study area, we determined that fortified commodities such as CSB and WSB could help meet micronutrient needs for children in the second year of life and beyond, but for infants 6-11 mo, even these fortified commodities were insufficient to allow infants to meet their daily iron and zinc

requirements (Ruel et al. 2004). Therefore additional micronutrient supplements may be required.

Results related to intake of fortified WSB were presented above and showed no major differences between program groups when children were in the program.

Traditional supplementation with vitamin A capsules (children and mothers) and with iron (mothers during pregnancy) was part of the range of services offered in both program areas. These services, however, were not offered only to program participants; they were available at the Rally Posts for the whole community.

Coverage with vitamin A capsules, both for index children in the last 6 months and for mothers immediately postpartum, was higher in the preventive than in recuperative areas, though differences were not large (differences of 6-9 percentage points in favor of the preventive group) (Table 8.6). Coverage with iron supplements during pregnancy was high in both program areas and did not differ between areas.

Compared to baseline, overall coverage for women had improved dramatically, particularly for vitamin A capsules. At baseline, coverage with iron pills during pregnancy was 65% compared to 83% in the final survey. At baseline, only 8% had received a postpartum vitamin A capsule, as compared to 59% in the final survey.

Finally, vitamin A capsule coverage for index children also increased dramatically, from 30% at baseline to 49% in the final survey, but still falls far short of universal coverage even among children who had ever participated in the program (52%, not shown).

Feeding during and after diarrhea

Recommended practices for feeding during and after diarrhea include giving more liquids and continuing to offer solid food while the child is ill, and giving an extra meal each day once appetite returns (as the child recovers). Results of the final survey show that overall, approximately half of mothers report giving more liquids, and 30% offering more semi-solid or solid foods when their child was ill (Table 8.6). There were no differences between program areas. For liquids, these results are very similar to baseline, but for semi-solid and solid foods, a marked increase over baseline is observed (6% at baseline compared to 30% at final survey). After the child recovered, 61% reported that they were able to give their child an extra meal each day, for a number of days ranging from 1-15 (median 7). Mothers in preventive areas were more likely to report giving an extra meal to the child during convalescence (67% versus 56% in recuperative areas).

8.4.4 Impact of interventions on preventive care, care during illness, and hygiene

Immunization

Approximately one-quarter (27%) of the index children were fully immunized - based on their health card or maternal recall - at the time of the final survey, with no differences between program groups (Table 8.7). Although these numbers are higher than at baseline (11%), they reflect very low coverage of immunization. Even among children who had participated in the

program at some point in the previous three years, the percent of fully immunized children was very low (31% for preventive and 26% for recuperative). These results, however, are consistent with the program's reported problems with supply of vaccines (see Chapter 5).

Table 8.7 Preventive care, care during illness, and hygiene, by program group and time of survey, for index children 12 to 42 months of age^a

Feeding/care practice	Baseline (n = 1,462)	Final survey		
		Preventive (n = 550)	Recuperative (n = 564)	All children (n = 1,114)
<i>Immunization status</i>	(n = 1,468)	(n = 551)	(n = 565)	(n = 1,116)
- Fully immunized (according to card)	12.9	28.9	25.0	26.8
- Fully immunized (from recall)	6.6	27.2	28.9	28.0
- Fully immunized (card or recall)	10.5	28.5	25.7	27.1
<i>Curative health-care-seeking</i>				
Sought treatment when ill with:				
- fever	82.2	66.9*	57.9*	62.0
- cough	81.0	51.4	49.2	50.3
- fast breathing	80.9	67.2	68.5	67.9
- diarrhea	72.0	45.0	36.8	40.9
Used ORS when child had diarrhea (denominator: children who had diarrhea in previous 2 weeks)	40.0	45.0	46.8	45.9
Used home-made sugar-salt-solution when child had diarrhea	10.0	27.2	19.9	23.5
Used either ORS or homemade sugar-salt-solution when child had diarrhea	45.4	41.4	40.9	41.2
<i>Hygiene scales (index child)</i>				
- Overall child cleanliness score (mean); max = 13	7.7	8.6	8.5	8.5
- Maternal cleanliness score (mean); max = 12	9.8	8.4	8.3	8.4
- House interior cleanliness score (mean); max = 6)	4.8	4.5	4.6	4.5
- House exterior cleanliness score (mean); max = 7)	5.8	5.4	5.6	5.4

* Significant at $p \leq 0.05$ (comparison of preventive and recuperative areas; results from xtlogit for dichotomous variables).

^a Baseline data for these variables were only available for children aged 18 mo and older; final data are shown for the same age group for comparison.

Health seeking practices for fever, respiratory infections, and diarrhea, and use of oral rehydration solution (ORS) for diarrhea

Table 8.7 also presents results on health-care-seeking practices for basic childhood illnesses such as fever, upper respiratory infections, fast breathing, and diarrhea. There are no striking differences between program groups for care-seeking for any of the illnesses. Compared to baseline, care-seeking for all four illnesses was markedly lower. Since health-care service provision had improved since baseline (with the availability of the WV MCHN program services), the lower care-seeking at the follow-up survey could indicate either lower severity of illness or poor availability of economic resources to use services. However, since most WV

services were available free of cost, the latter was less likely to be of an explanation for lower rates of care-seeking.

Use of ORS for diarrhea was also not different between the two program groups, and there was little difference from baseline. However, in the final survey women were more likely to report having used a home-made sugar-salt solution for diarrhea.

Hygiene

We collected data on proxies for hygiene practices, which could be assessed by spot-check observations (see description of method in section 8.2). Results show no difference in any of the hygiene scales between program groups (Table 8.7).

In summary:

- ❖ At baseline there were no differences between program groups in feeding practices, care-seeking, or proxies for hygiene practices.
- ❖ In the final survey, there were no differences between program groups in early feeding practices (initiation of BF and feeding of colostrum). Large improvements were seen since baseline.
- ❖ Similarly, there were no differences between program groups in the timing of introduction of liquids and complementary foods, but large improvements were seen since baseline, with a majority of women reporting initiating feeding of both liquids and semi-solids at approximately 6 months of age.
- ❖ There were no differences between program groups in continued breastfeeding up to 24 months, and no differences since baseline either. Haitian women's breastfeeding practices are generally positive, with children being breastfed until between 18 and 24 months of age.
- ❖ Slightly more young children in the preventive group received assistance while eating than in the recuperative group; there was a moderately large increase in this practice over baseline.
- ❖ Use of baby bottles was not different between program groups, but rates of use were halved since baseline.
- ❖ There were no differences between program groups in the frequency of feeding, but meal frequency for children in the vulnerable age of 12-17 months of age had increased since baseline.
- ❖ There were no differences between program groups in the practice of feeding the child an evening meal, and no difference since baseline either.
- ❖ Diet quality was slightly better in the preventive group overall, as assessed by several proxies. Children in preventive areas were more likely to have had nutrient-rich food

groups, including eggs and beef/pork in the previous day or week, and mean dietary diversity was higher. The overall consumption of ASF was lower than at baseline, possibly due to the economic crisis in Haiti, which had led to increased food prices over the study period.

- ❖ Scores on the IYCF indicator – which combines information on breastfeeding, feeding frequency, and dietary diversity - were slightly higher in the preventive group than in the recuperative group.
- ❖ Vitamin A supplementation rates were higher in the preventive group, with large differences seen since baseline.
- ❖ Feeding practices during and after diarrhea were also better in the preventive group, particularly for reported increases in meal frequency after an episode of diarrhea. Large improvements were also seen since baseline.
- ❖ Immunization rates were not different between the program groups, but had almost tripled since baseline. However, rates remain very low.
- ❖ There were no differences in markers of hygiene practices between program groups.
- ❖ Care-seeking for fever, cough, fast breathing, and diarrhea were not different between program groups. Care-seeking rates were lower at the follow-up survey than at the baseline survey.
- ❖ Use of ORS when the child had diarrhea did not differ between program groups and had not changed markedly since baseline; use of home-prepared solutions had, however, increased.

8.4.5 Comparisons between participants and nonparticipants on child-feeding and care practices

As noted in the methods section, additional analyses for some key outcomes were done to compare child-feeding and care practices between participants and nonparticipants, within the preventive and recuperative program communities. The purpose of this analysis was to assess whether participation in the program was associated with a meaningful change in practices. Except when specified otherwise, the indicator used for participation here is “ever participated” in the program (and therefore exposed to the BCC delivered in the MCs). A summary of the differences by participation is presented in Table 8.8, and detailed results are presented in Annexes 8.5 to 8.12. We provide here a brief summary of the practices where marked differences were found between participant and nonparticipant mothers.

Table 8.8 Summary of findings on behavior change outcomes

	Preventive vs. Recuperative ^a	Change compared to baseline ^b	Difference between ever participated vs. never participated ^c
Knowledge			
<i>Child feeding knowledge tests</i>			
- Timely introduction of foods	- ^d	↑	Large diff
- Feeding frequency	Prev > (small diff.)	↑	Large diff
- Feeding knowledge (weighted)	Prev > (small diff.)	↑	Large diff
<i>Overall nutrition and health knowledge test (BCC topics)</i>			
- Breastfeeding knowledge	-	n/a	Large diff
- CF knowledge	Prev > (small diff.)	n/a	Large diff
- Child illness	Prev > (small diff.)	n/a	-
- General health issues	-	n/a	-
- Malnutrition	-	n/a	-
Awareness, trial and adoption of key practices			
- Awareness	Prev > (small diff.)	n/a	Large diff
- Trial	Prev > (small diff.)	n/a	Large diff
- Adoption	Prev > (small diff.)	n/a	Large diff
Care and feeding practices			
<i>Early infant practices</i>			
BF within 1 hour	-	Large ↑	Large diff.
EBF on 1 st day	-	Large ↑	Large diff.
Gave colostrum	Recup >	Large ↑	Large diff.
<i>Timing of introduction of liquids and foods</i>			
Continued BF	-	Large ↑	Large diff.
Responsive feeding	-	-	-
	Prev > (for helps child feed)	↑	More positive strategies (prev.)
Use of baby bottles	-	↓ by half	Large diff.
Meal frequency	-	↑ (for 12-17 mo, BF)	> if child currently in program
Evening meal	-	-	-
<i>Dietary diversity (DD)</i>			
- Food group consumed	Prev > for nutrient-rich foods (small differences)	↓ for nutrient-rich foods	-
- Mean dietary diversity	Prev > at 12-17 mo	↓	Small diff. in some age groups
- Animal source food consumed	Prev > for eggs, beef/pork	↓ by half	Small diff.
- Consumption of WSB while in program	-	n/a	Large diff. (as expected)
<i>IYCF indicator</i>			
- Score	Prev > (small diff)	Not available	No diff in prev.; partic<nonpartic. in recup.
- % meeting recommendations for all 3 practices	Prev >	Not available	No diff in prev.; partic<nonpartic. in recup.
<i>Vitamin and mineral supplements</i>			
Feeding during and after diarrhea	Prev > for VA Prev > for extra meals after recovery	Large diff. Large ↑ in giving more food during diarrhea	Large diff. Large diff for extra meals after recovery

(continued)

	Preventive vs. Recuperative ^a	Change compared to baseline ^b	Difference between ever participated vs. never participated ^c
<i>Preventive care, hygiene, and curative care seeking</i>			
<i>Immunization: fully immunized</i>	-	Large ↑	Diff. only in prev.
<i>Hygiene scores</i>	-	↓	Small diff. in recup for child cleanliness
<i>Curative care seeking</i>	-	↓	Small diff for fever
<i>Use of ORS when child has diarrhea</i>	-	-	No diff.
<i>Use of home-made SSS when child has diarrhea</i>	-	↑	Diff. only in prev.

^a Only statistically significant differences reported for differences between preventive and recuperative program groups.

^b Only large differences are reported – no statistical testing.

^c Only large differences are reported – no statistical testing.

^d “-” means no difference; “n/a” means not applicable.

Early infant feeding practices and timing of introduction of liquids and semi-solid and solid foods

In both program groups, a much larger proportion of mothers who had been exposed to the BCC intervention reported optimal early infant feeding practices compared to nonparticipant mothers (Annex 8.5). The same was true for the timing of introduction of liquids and semi-solid and solid foods. Among nonparticipant mothers, 30% reported giving liquids by 2 months and 56% by 6 months, compared to those who had been exposed to the MCs (4% reported giving liquids by 2 months and 14% did so before 6 months). Nonparticipant mothers were also more likely to have introduced semi-solid foods before the age of 6 months (close to 50%) compared to participant mothers (approximately 10%) (Annex 8.6).

Use of baby bottles

There were large differences in the use of baby bottles by program participation, with high use of baby bottles among mothers who had never been exposed to the program compared to those who had. In the preventive group, 46% of nonparticipants reported using baby bottles compared to 12% among participants. In the recuperative group, differences were of smaller magnitude but showed a similar pattern: 24% among nonparticipants compared to 15% among participants.

Meal frequency

Children whose mothers had ever participated in the MCs were no more likely to be fed (at least) the recommended minimum number of times the day before the survey. However, if the child was currently receiving a food ration, the mean number of meals was higher both for 12-23-month-old children and for younger siblings (6-11 months old). This, however, was not true for older children. Most strikingly, non-breastfed children aged 12-23 months were nearly 3 times as likely to be given at least the minimum recommended number of meals (four) if they were currently receiving a food ration (23% versus 8% for those not receiving a ration) (Annex 8.7). These results suggest that food assistance, which at least temporarily improves household food security, may have short-term beneficial effects on frequency of feeding for non-breastfed

children. This is particularly important because as discussed above, very few non-breastfed children were fed as frequently as recommended.

Nutrient content of complementary foods: Dietary quality and diversity

There were few differences in diet quality related to program participation, when participation was defined as “ever participated.” Children whose mothers were never exposed to the program were much less likely to have consumed eggs in the previous day, especially in the preventive program areas (Annex 8.9); eggs were specifically recommended in the recipes promoted and demonstrated at the MCs.

Also, as could be expected, the use of fortified foods—which in this population was largely limited to WSB—was much more common among *current* program participants (in both program groups) than among nonparticipants (Annex 8.11). This is expected as WSB is only available through food assistance programs in Haiti.

Overall dietary diversity was also slightly higher among participant compared to nonparticipant children in both program groups in certain age groups (children aged 18-23 months in preventive areas, and children aged 24-29 months in recuperative areas) (Annex 8.12).

IYCF indicator: Mothers who had been exposed to the program in the preventive group had better scores, and were more likely to have met minimum recommendations for all 3 practices, than mothers who were exposed in the recuperative group. There was little difference between exposed and nonexposed mothers in the preventive group. In the recuperative group, however, the IYCF indicator scores and the percent meeting minimum recommendations were higher among *nonexposed* mothers than among exposed mothers. This could be due to reverse causality; children who are fed poorly are more likely to be malnourished and thus be eligible for enrollment in the recuperative program (i.e., be “exposed”).

Use of vitamin-mineral supplements. Mothers who had been exposed to the program were much more likely to have received prenatal iron supplements and postpartum vitamin A in both program groups. Children were also more likely to have received vitamin A supplements in the previous 6 months (Annex 8.13). These findings suggest that, although these services are available at RPs for the entire population, they appear to be used more by women who have participated in the MCs, and their children.

Feeding during and after diarrhea. Preventive area mothers who had participated in the MCs were more likely to offer more extra food to their child during a diarrheal episode. Participating mothers in both areas were more likely to feed their child an extra meal after recovery from diarrhea (Annex 8.14).

Preventive care, care during illness, and hygiene

When compared to nonparticipants, immunization rates were higher among participants in the preventive group (Annex 8.15). There were no differences in proxies for hygiene practices between participants and nonparticipants in either program group, except for child cleanliness where participants in the recuperative group had better scores than nonparticipants (results not

shown). Care-seeking for fever, cough, fast breathing, and diarrhea were also not very different between participants and nonparticipants.

However, use of ORS when the child had diarrhea was higher by about 10 percentage points among participants when compared to nonparticipants but differences were not significantly different. Use of salt-sugar solution (SSS) was marginally significantly higher ($p = 0.06$) among participants in the preventive group but not in the recuperative group.

In summary:

- ❖ There were large differences in early feeding practices between ever-participants and never-participants, i.e., initiation of BF and feeding of colostrum.
- ❖ There were large differences between ever-participants and never-participants in the timing of introduction of liquids and complementary foods.
- ❖ There were no differences between ever-participants and never-participants in continued breastfeeding up to 24 months.
- ❖ Use of baby bottles was significantly lower among ever-participants compared to never-participants.
- ❖ Meal frequency was higher for children who were currently enrolled in the program; non-breastfed children currently receiving rations were much more likely to be fed with recommended frequency.
- ❖ There were no differences between ever-participants and never-participants in the practice of feeding the child an evening meal.
- ❖ Among children 6-23 months of age, there were no differences between ever-participants and never-participants in the preventive group on the IYCF indicator. Among the recuperative group, however, ever-participants had *lower* scores on the IYCF indicator than never-participants.
- ❖ Consumption of fortified WSB was markedly higher among ever-participants, as expected.
- ❖ Vitamin A supplementation rates were higher among ever-participants (both women and children) than among never-participants. Women who had participated were also more likely to have received iron supplements during pregnancy.
- ❖ Feeding practices during (preventive area) and after diarrhea (both areas) were different between ever-participants and never-participants.
- ❖ Immunization rates were also different between ever-participants and never-participants, but only in the preventive group.

- ❖ There were no differences in markers of hygiene practices between ever-participants and never-participants in either program group, except for child cleanliness where participants in the recuperative group had better scores than never-participants.
- ❖ Care-seeking for fever, cough, fast breathing, and diarrhea were not very different between ever-participants and never-participants.
- ❖ Use of ORS when the child had diarrhea was higher among ever-participants than never-participants but differences were not significant. Use of salt-sugar solution (SSS) was marginally significantly higher among ever-participants.

8.5 Conclusions

This chapter presented a wide range of results related to behavior change outcomes. The BCC activities, delivered through the MCs (and described in Chapters 3, 4, and 5) comprised the primary program inputs aimed at influencing behavior change. These inputs were expected to lead to improvements in maternal knowledge and skills and thereby to improved infant feeding practices (see program theory description in Chapter 2). Differences in practices between program groups were expected due to two specific aspects of program design: (1) the timing of delivery of messages regarding complementary feeding, which was designed to be provided *before* the child reaches complementary feeding age for the preventive group; and (2) the duration of exposure to the BCC, which was longer for the preventive (up to 18 months) compared to the recuperative group (maximum 9 months). As highlighted in Chapters 5 and 6, both of these critical design characteristics were achieved and mothers in preventive areas were thus more likely to receive the education/BCC in a timely fashion and to be exposed to the messages for a longer duration than mothers in the recuperative areas. In addition to the differences in timing and duration of exposure, the results on program participation also showed greater program participation rates after the child was 6 months old in preventive compared to recuperative areas (Chapter 6), thus suggesting an additional mechanism by which the preventive program may have had a greater impact on behavior change than the recuperative program.

The results presented here demonstrate that the program pathway and the expectations based on program design were generally achieved as intended, and differences between program groups were consistent with expectations based on program targeting and exposure. For instance, there were no differences between the groups in knowledge and practices related to early breastfeeding and exclusive breastfeeding. This was expected, given that exposure to the program was similar in both groups during pregnancy and the first 6 months of lactation. By contrast, differences in favor of the preventive group were observed in several of the complementary feeding behavior change outcomes; again this was expected, given that the preventive approach was intended to provide greater intensity and more timely delivery of education outcomes. In general, however, the magnitude of differences between the groups was quite small for most behavior change outcomes.

Although overall differences between program groups were generally small, differences since baseline in both groups were quite large; this suggests that the program was successful in raising overall child-feeding and care knowledge and practices in the communities where it operated. Differences between participants and nonparticipants were also large and statistically

significant for many behavior outcomes. Participants fared generally much better than nonparticipants for most BCC outcomes, from knowledge, trial, and adoption to the full spectrum of reported child-feeding practices. This was true in both the preventive and recuperative program groups, and can be interpreted as reflecting programmatic success. It also suggests that the process used to develop the BCC program, the technical content and staff development intrinsic to the program, was successful, as was the mechanism used to deliver the BCC program (i.e., the small peer-group approach). Finally, since improvements in practices were seen for younger siblings in both program groups, the results also suggest that the positive benefits of a solidly designed and well-implemented BCC program extend beyond immediate effects. It will be important to continue to monitor knowledge and practices related to infant and young child feeding in these Haitian communities to evaluate the true long-term impact of the investments in the development of such a strategy.

9. IMPACT OF THE PROGRAM ON FOOD SECURITY

9.1 Introduction

This chapter discusses the impact of the WV-Haiti MCHN program on food security outcomes, particularly those outcomes related to food access (rather than utilization or availability). The WV MCHN program, by virtue of providing food rations to households and supportive learning environments (via Mothers' Clubs), could directly impact household resources such as food security, and such impacts could be greater in the preventive than in the recuperative program communities, since program benefits are provided for longer durations and to a large proportion of households. This chapter therefore examines differences between the two program groups on household food security outcomes. It also compares the food security indicators at baseline in 2002 and at the final survey in 2005.

9.2 Data and Analysis

9.2.1 Data and variables

The final impact survey gathered data on two dimensions of household food insecurity:

(1) *Household experiences* related to food insecurity. Data were collected on 11 types of food-insecurity experiences. These experiences of food security covered a range from less extreme to more extreme, e.g., "cooking without beans" (a preferred food), which was at the less extreme end of the experiences, compared to "going to bed hungry," which was considered a more extreme experience of food insecurity.

Data on these 11 individual experiences related to food insecurity were used to develop a composite food-insecurity scale that summarized the information from the individual variables into a meaningful composite measure. The scale was constructed by summing the total number of food-insecurity-related experiences that a household had faced in the past 30 days. All the variables included contributed a maximum of 1 point to the scale, and variables with multiple categories were recoded so that the highest category contributed 1 point, the lowest category, 0 points, and categories in between contributed between 0 and 1 point. Thus, the scale ranged from a minimum of 0 to a maximum of 11, where households with a score of 11 would have experienced extreme levels of food insecurity. The distribution of the food-insecurity experiences scale was divided into terciles to create three groups that represented low, moderate, and severe food insecurity.

The detailed scoring of the variables that were included in the scale is presented in Annex 9.1. The reliability of the household food-insecurity experiences scale was tested using Cronbach's alpha. The scale had a reliability of 0.77.

(2) *Months of inadequate household food provisioning* (MIHFP). The MIHFP measure was adapted from an indicator developed by FANTA (Bilinsky and Swindale 2005). Respondents were asked to indicate whether they had enough food for household consumption in each individual month in the previous 12 months. Responses were on a 3-point scale and ranged

from “Yes, enough food,” to “No, not enough at all.” Responses to these questions were summed up to create two summary variables:

Months of inadequate household food provisioning (MIHFP): This summary variable captured the number of months that a respondent reported any food inadequacy in the household. In order to create it, the 3-point scale was recoded such that any kind of insufficiency (either “not enough at all” or “somewhat enough”) was coded “yes.” Although this could overestimate the number of months of food inadequacy, we used this coding because it is closer to the responses we would have received from respondents had we used the original FANTA yes/no response sets for these questions (Bilinsky and Swindale 2005).

Severity of food inadequacy: A summary variable was created by adding up responses on the 3-point scale for all 12 months in the MIHFP measure. A higher score indicated more severe food inadequacy. The distribution of the score for the entire survey sample was also divided into terciles to enable categorical comparisons. The terciles were labeled “low,” “medium,” and “high,” and the proportion of households in the “high” group was compared between program approaches.

9.2.2 Analysis

A systematic stepwise approach focusing on differences between program communities at the time of the final survey was used for the analysis. It included the following steps:

- 1) First, we assessed the unadjusted difference between program communities using random effects regression methods that adjusted for the paired design.
- 2) Next, the difference between program communities was examined after adjusting for common confounding factors like respondent’s education level and partner’s education level, depending on the outcome of interest.
- 3) Third, we examined benefits of program participation by separating participants and nonparticipants, and examining the differences between participants and nonparticipants in the two program groups. We also note any relevant differences between nonparticipants in each of the program groups. In examining differences between program participants and nonparticipants within program group, we used two measures of participation: (1) whether a household/mother had ever received food rations; (2) whether a household/mother was receiving program benefits at the time of the survey. We evaluated the significance of differences between participants and nonparticipants using interaction terms between participation and program group, evaluating separately the interactions between “ever participation” and program group and between “current participation” and program group.
- 4) Finally, where possible, we draw informal comparisons with the baseline results to evaluate the extent of change from baseline.

In general, in evaluating the differential influence of the program approaches on these outcomes of interest, we do not adjust for any variables that could be influenced by the program. Adjusting for such variables would underestimate the impact of the program.

9.3 Results

9.3.1 Impact of the Program on Household Food Insecurity

Descriptive results for the three measures of food insecurity - household food insecurity experiences scale, number of months of food inadequacy, and severity of food inadequacy scale - are presented in Table 9.1. Note that there were no differences in food insecurity at baseline and, therefore, the baseline results are presented for the two programs combined. Table 9.1 shows that preventive program communities generally fared better in terms of household food security at the time of the final survey than the recuperative communities, where food insecurity was consistently higher.

Table 9.1 Household food insecurity, by program and time of survey

		Baseline (2002)	Final survey (2005)		
		Both groups (n = 1,514)	Preventive (n = 748)	Recuperative (n = 752)	Both groups (n = 1,500)
Food security characteristics	Variable		Mean (SD)	Mean (SD)	Mean (SD)
Household food insecurity experiences (scale)	<i>fdinsec</i>	7.8 (1.8)	7.7 (2.4)	7.9 (2.3)*	7.8 (2.3)
Number of months of food inadequacy	<i>Mo_insuf</i>	n/a	8.8 (2.8)	9.2 (2.7)*	9.0 (2.8)
Severity of food inadequacy	<i>sevinsuf</i>	n/a	25.7 (5.4)	26.5 (5.4)*	26.1 (5.4)

* $p < 0.05$ (differences tested using unadjusted random effects regression models).

Compared to baseline, food insecurity continues to be extremely high in these communities: scores on the food insecurity experiences scale for both groups combined are almost identical between the baseline survey in 2002 and the combined final survey in 2005.

An examination of the individual household food insecurity experiences included in the household food insecurity experiences scale (Table 9.2) shows that the preventive program likely better mitigated some of the more severe food insecurity experiences. Overall, the prevalence of experiences reflecting severe food insecurity in this context, i.e., respondents going to bed hungry, children going to bed hungry, and cooking the same food every day, are significantly lower in the preventive group than in the recuperative group. They are also lower in 2005 compared to 2002 (both groups combined), suggesting that, overall, the program has likely relieved some of the more severe food insecurity experiences.

The impact of the program on food insecurity experiences appears to be driven by current participation in the program. Comparing current program participants within program group with nonparticipants suggests that current participants are slightly less food insecure than nonparticipants, and especially so among preventive communities (Table 9.3 and Figures 9.1 - 9.3). Regression models that include current participation show that the magnitude and significance of the program approach variable is diminished with the introduction of the current participation variable (Table 9.4 - 9.6). This is true for all three measures of food insecurity.

Table 9.2 Selected household food insecurity experiences, by program and time of survey

Food security characteristics	Variable	Baseline	Final survey		
		Both groups (n = 1,514)	Preventive (n = 748)	Recuperative (n = 752)	Both groups (n = 1,500)
			(%)	(%)	(%)
Cooked with less beans	<i>Q713</i>	98.0	97.6	98.0	97.8
Cooked without any beans	<i>Q711</i>	96.4	94.5	94.9	94.7
Ate less food because of hardship	<i>Q708a</i>	89.7	87.0	88.3	87.7
Worried about not having enough food	<i>Q719a</i>	88.2	86.5	89.2	87.9
Cooked without herring head	<i>Q715a</i>	87.1	77.5	77.4	77.5
Ate same food day after day	<i>Q716</i>	85.0	64.6	72.2**	68.4
Went to bed hungry	<i>Q717a</i>	83.9	73.8	78.5*	76.1
Children went to bed hungry	<i>Q718a</i>	75.5	59.9	66.8**	63.3
Cooked less because of lack of fuel	<i>Q709a</i>	50.2	48.5	48.5	48.5

* $p < 0.05$ (differences tested using unadjusted random effects regression models).

Table 9.3 All household food insecurity measures at final survey, by participation (ever and current)

Food security characteristics	Variable	Preventive				Recuperative			
		Non-participant		Participant		Non-participant		Participant	
		Mean	SD	Mean	SD	Mean	SD	Mean	SD
(1) Ever participated									
		(n = 126)		(n = 622)		(n = 130)		(n = 622)	
Household food insecurity experiences (scale)	fdinsec	7.8	2.2	7.6	2.4	7.9	2.5	8.0	2.3
Months of inadequate household food provisioning (MIHFP)	Mo_insuf	8.9	2.9	8.8	2.8	9.2	2.8	9.2	2.7
Severity of food inadequacy	sevinsuf	25.8	5.6	25.8	5.4	26.6	5.6	26.5	5.3
(2) Current participation									
		(n = 423)		(n = 325)		(n = 621)		(n = 131)	
Household food insecurity experiences (scale)	fdinsec	7.9	2.3	7.4	2.4	8.0	2.3	7.7	2.3
Number of months of food inadequacy	Mo_insuf	9.2	2.7	8.4	2.9	9.2	2.7	9.1	2.7
Severity of food inadequacy	sevinsuf	26.3	5.3	25.0	5.4	26.6	5.4	26.3	5.4

Figure 9.1 Household food insecurity experiences score, by program group and current participation at final survey




Figure 9.2 Prevalence of severe food insecurity (based on terciles of the household food insecurity experiences score), by program group and current participation at final survey




Figure 9.3 Months of inadequate household food provisioning (MIHFP), by program group and current participating at final survey




Table 9.4 Program impact on household food insecurity experiences at final survey (regression analysis)

	(1) Bivariate model	(2) Main effects, adjusted	(3) Main effects, ever participated	(4) Interaction model, ever participated	(5) Main effects, currently participating	(6) Interaction model, currently participating
Household food insecurity experiences score						
Program	0.27* (0.12)	0.28* (0.13)	0.28* (0.13)	0.15 (0.33)	0.15 (0.14)	0.13 (0.16)
Respondent's education (years)		-0.11** (0.03)	-0.11** (0.03)	-0.11** (0.03)	-0.11** (0.03)	-0.11** (0.03)
Partner's education (years)		-0.07** (0.02)	-0.07** (0.02)	-0.07** (0.02)	-0.07** (0.02)	-0.07** (0.02)
Ever participated			-0.06 (0.18)	-0.13 (0.25)		
Currently participating					-0.48** (0.15)	-0.51** (0.19)
Ever participated and Program				0.15 (0.36)		
Currently participating and Program						0.09 (0.30)
Constant	7.68** (0.11)	8.06** (0.12)	8.10** (0.19)	8.17** (0.24)	8.26** (0.13)	8.28** (0.14)
Observations	1,500	1,250	1,250	1,250	1,250	1,250
Number of pairs	10					

Notes: Standard errors in parentheses. * significant at 5%; ** significant at 1%.

Table 9.5 Program impact on months of inadequate household food provisioning (MIHFP) at final survey (regression analysis)

	(1) Bivariate model	(2) Main effects, adjusted	(3) Main effects, ever participated	(4) Interaction model, ever participated	(5) Main effects, currently participating	(6) Interaction model, currently participating
Months of inadequate household food provisioning (MIHFP)						
Program	0.34* (0.14)	0.45** (0.16)	0.45** (0.16)	0.30 (0.39)	0.33* (0.16)	0.14 (0.19)
Respondent's education (years)		-0.11** (0.03)	-0.11** (0.03)	-0.11** (0.03)	-0.11** (0.03)	-0.11** (0.03)
Partner's education (years)		-0.02 (0.03)	-0.02 (0.03)	-0.02 (0.03)	-0.02 (0.03)	-0.02 (0.03)
Ever participated			-0.04 (0.22)	-0.13 (0.30)		
Currently participating					-0.45** (0.18)	-0.71** (0.22)
Ever participated and Program				0.17 (0.43)		
Currently participating and Program						0.69 (0.36)
Constant	8.83** (0.19)	8.98** (0.18)	9.02** (0.26)	9.10** (0.32)	9.18** (0.20)	9.29** (0.21)
Observations	1,500	1,250	1,250	1,250	1,250	1,250
Number of pairs	10	10	10	10	10	10

Notes: Standard errors in parentheses. * significant at 5%; ** significant at 1%.


Table 9.6 Program impact on severity of months of inadequate household food provisioning (MIHFP) at final survey

	(1) Bivariate model	(2) Main effects, adjusted	(3) Main effects, ever participated	(4) Interaction model, ever participated	(5) Main effects, currently participating	(6) Interaction model, currently participating
Severity of months of inadequate household food provisioning (MIHFP)						
Program	0.75** (0.27)	0.96** (0.30)	0.96** (0.30)	0.55 (0.76)	0.76* (0.31)	0.41 (0.36)
Respondent's education (years)		-0.27** (0.06)	-0.27** (0.06)	-0.27** (0.06)	-0.27** (0.06)	-0.27** (0.06)
Partner's education (years)		-0.00 (0.05)	-0.00 (0.05)	-0.00 (0.05)	-0.00 (0.05)	-0.00 (0.05)
Ever participated			0.04 (0.41)	-0.20 (0.59)		
Currently participating					-0.75* (0.34)	-1.23** (0.42)
Ever participated and Program				0.49 (0.83)		
Currently participating and Program						1.30 (0.70)
Constant	25.76** (0.44)	26.02** (0.43)	25.99** (0.56)	26.19** (0.66)	26.35** (0.45)	26.56** (0.49)
Observations	1,500	1,250	1,250	1,250	1,250	1,250
Number of pairs	10	10	10	10	10	10

Notes: Standard errors in parentheses. * significant at 5%; ** significant at 1%.

Figure 9.4 maps the responses to the questions related to months of food insecurity experienced by households, by month. It shows a consistent pattern of a slightly lower percentage of households reporting insufficient food at each month in the preventive group, but differences are of small magnitude. As with the previous analyses, however, Figure 9.4 also shows that, in fact, a large proportion of households report food inadequacy throughout the year, ranging from a low of around 50% in January to a high of around 90% in May and June. This pattern corresponds to the cropping and harvest season in the Central Plateau of Haiti. The results confirm the severity of food inadequacy and food insecurity in these Haitian rural communities, while demonstrating the potential for food assistance programs to make a difference.

Figure 9.4 Proportion of households with insufficient food in each month of the calendar year, by program group (derived from the months of inadequate household food provisioning (MIHFP) measure)



In summary, using all three measures of food insecurity, our results demonstrate that

- ❖ There were no differences in food security experiences between groups at baseline.
- ❖ At the end of the study, the three indicators of food insecurity used (household food insecurity experiences, months of inadequate household food provisioning (MIHFP), and severity of MIHFP) were statistically significantly different between the groups showing greater food insecurity among recuperative compared to preventive

communities; experiences of severe food insecurity (e.g., child going to bed hungry, etc.) were also statistically significantly higher among recuperative compared to preventive communities. In general, however, differences between program groups were of small magnitude.

- ❖ At final survey, there were no major improvements from the baseline survey in any of the food insecurity measures for either program group; food insecurity continued to be severe in this region in Haiti.
- ❖ Multivariate models show that current participation in either program approach (preventive or recuperative) is positively associated with reduced food insecurity, number of months of food inadequacy, and severity of months of food inadequacy. This suggests that both program approaches have a short-term impact on reducing household food insecurity.
- ❖ The lack of difference between households who had ever participated in the program and those who had not, however, suggests that the program does not have a long-term effect on food insecurity.

9.4 Conclusions

The results presented in this chapter suggest that, overall, food insecurity continues to be severe and intractable in the Central Plateau of Haiti. The WV MCHN program was able to alleviate severe experiences of household food insecurity, primarily for those households who participated in the program. The differences in food insecurity between program approaches are generally small, but differences between current participants and nonparticipants are significant, suggesting a short-term impact on food insecurity while households are in the program. In spite of reasonably long durations of participation in the food assistance program, there do not appear to be long-term impacts on food insecurity of having ever participated in the program. This reinforces the need for long-term community food security, agricultural, and economic development programs that can generate sustainable and longer term solutions to the severe food insecurity in this region.

10. INDIRECT BENEFITS OF THE PROGRAM

10.1 Introduction

This chapter assesses the potential indirect benefits of participation in the WV MCHN program. More specifically, it examines the impact of the program on household assets, women's assets, and maternal mental and physical well-being, all of which could be positively affected via increased food security. In the baseline study, we documented strong associations between maternal stress and other indicators of poor mental health and food insecurity (Menon et al. 2004). This suggested the possibility that the program inputs, which were intended to improve food security, could also have had spin-off benefits that led specifically to improved women's well-being.

10.2 Data and Analysis

10.2.1 Data and variables

10.2.1.1 Household assets

The food assistance provided by the program constitutes a significant transfer of food on a monthly basis. It could be hypothesized that such transfers of food assistance could free up household resources for other expenditures, which may or may not be reflected in changes in household asset ownership.

The survey questionnaire gathered household-level data on ownership of 17 durable goods (e.g., cooking utensils, furniture, electronics, and so forth), 9 productive assets (agricultural tools), and 6 types of small animals and livestock. Respondents were asked whether or not they owned these individual assets; for the animals and livestock, they were also asked how many of each one they owned. From this data, three simple count indices were created to capture the number of assets owned: these indices consisted of a simple count of whether or not the households owned the assets listed (using Yes = 1 and No = 0). The three indices created are referred to as *durable goods count*, *productive assets count*, and *livestock count*.

10.2.1.2 Women's ownership of assets

Women's ownership of assets was assessed by asking respondents if they owned any of four assets (land, the house they lived in, another house, and livestock) either on their own or jointly with someone. For each asset that a woman owned either alone or jointly, a score of 1 was assigned. A score of 0 was assigned if she did not own the asset. All four asset-ownership variables were then summed to create a scale of ownership of assets that ranged from 0 to 4.

10.2.1.3 Women's physical well-being: Self-rated health visual analogue scale

Women's physical health and nutritional status were assessed by asking respondents to self-rate their health using a visual analogue scale. The scale comprised of a line, 10 centimeters long, where the left end of the line symbolized extremely poor health compared to other women of the respondent's age, and the right-hand end of the line stood for better health than other

women her age. The respondent was asked to mark the spot on the line where she felt her health stood, thus generating data on self-perceived health status.

10.2.1.4 Maternal stress and mental well-being

Women's mental well-being and stress have been hypothesized to influence the quality of childcare, particularly psychosocial care. In our baseline survey of 2002, we found that women's well-being measures were positively associated with child feeding practices such as dietary diversity and consumption of nutrient-rich animal source foods (Menon and Ruel 2003).

In our baseline and final surveys, to characterize maternal mental health and stress levels, we assessed the presence and absence of various symptoms associated with poor mental or emotional well-being (e.g., feeling sad or unhappy, difficulty sleeping, difficulty in enjoying daily activities, etc.), as well as the frequency with which women reported experiencing various symptoms of poor mental health. Data were also obtained on women's satisfaction with their current life situation and their perceptions about the amount of time they had for their daily activities. These data were further combined to create summary scales of overall mental stress, frequency of stress symptoms, time stress, and life satisfaction.

- 1) Mental stress. In order to assess overall mental stress, the variables that measured the presence (coded as 1) or absence (coded as 0) of different symptoms were summed up to create an overall scale of maternal stress and poor mental well-being. Six variables were used to create the mental stress scale, and the scale had a minimum of 0, which indicated that the respondent experienced none of the mental health symptoms, and a maximum of 6, which indicated that the respondent suffered from all six symptoms, or a high level of stress.
- 2) Frequency of stress symptoms. Variables that assessed the frequency of occurrence of different symptoms related to anxiety and stress (such as poor appetite, headaches, poor digestion, fatigue, etc.) were combined by recoding each variable such that for each symptom, the lowest frequency of "never" received a score of 0 and the highest frequency of "often" received a score of 2. Following this, the scores on each of the variables were summed to create an overall scale that had a minimum value of 0 and a maximum of 20. A high score on the scale indicated higher levels of stress, in terms of the number of symptoms a woman suffered from, and the frequency with which she experienced them.
- 3) Time stress. Women's time and workload are often referred to as crucial resources in enabling them to care for their children. We assessed the amount of time women spent working outside their homes in order to measure this construct. However, it was also determined that it was important to assess a woman's own perception of how much pressure she felt for time to spend on her daily activities and caring for her children. We asked women how often they felt that they did not have enough time to take care of their homes, their children, and themselves, and also how often they worried about not having enough time to finish their daily work. The information on these four variables was combined by assigning scores of 0 to responses of "never," 1 to a response of "sometimes," and 2 to a response

of “often” to each of the four variables. These scores were then summed to create an overall time stress scale where a low score indicated a low level of stress due to time pressures and a high score reflected a high level of time pressures. The scale thus ranged from 0 to 8.

- 4) Life satisfaction. This was assessed by asking respondents about their satisfaction with various aspects of their lives, including their daily work and the help they received from their spouse and other family members. The individual variables were recoded to assign points to the level of satisfaction that women reported; this was done by assigning a score of 0 to women who reported being dissatisfied with a particular aspect of their lives, 1 if they were ambivalent, and 2 if they were satisfied. The scores on 10 individual variables were then summed to create an overall scale of a respondent’s level of life satisfaction where a low score (minimum of 0) indicated a low level of satisfaction and a high score (a maximum of 20) indicated a high level of satisfaction.

A second life satisfaction variable was created by dropping the variables that pertained to satisfaction with help received from the spouse or mother-in-law - the inclusion of these variables led to missing values for those who were not married or partnered. The life satisfaction scale that did not include these variables had a minimum of 0 and a maximum of 16.

Annex 10.1 presents the variables and the scoring used to develop these scales.

10.2.2 Analysis

A systematic step-wise approach was taken to the analyses, all of which focused on differences between the program communities at the time of the final survey. The following steps were followed:

- 1) The first step was to assess the unadjusted differences between program communities using random effects regression methods that adjusted for the paired design.
- 2) Next, the differences between program communities were examined after adjusting for common confounding factors like respondent’s education level and partner’s education level, depending on the outcome of interest.
- 3) Third, we examined benefits of program participation by separating participants and nonparticipants, and examining the differences between participants and nonparticipants within the two program groups. We also note any relevant differences between nonparticipants in each of the program groups. In examining differences between program participants and nonparticipants within program group, we used two measures of participation: (1) whether a household/mother had ever received program benefits; (2) whether a household/mother was receiving program benefits at the time of the survey. We evaluated the significance of differences between participants and nonparticipants using interaction terms between participation and program group, evaluating separately

the interactions between “ever participation” and program group and between “current participation” and program group.

- 4) Finally, where possible, we draw informal comparisons with the baseline results to evaluate the extent of change between baseline and the final survey.

In general, in evaluating the differential influence of the program approaches on these outcomes of interest, we do not adjust for any variables that could be influenced by the program. Adjusting for such variables would underestimate the impact of the program.

10.3 Results

10.3.1 Impact of the program on household and respondent asset ownership

There were no significant differences between program communities in household or respondent asset ownership at baseline. At the time of the final survey, the preventive group was slightly better off than the recuperative group in terms of household durable good assets and productive assets, but there were no differences between the groups in livestock assets and women’s ownership of assets (Table 10.1).

Although no consistent pattern of differences between participants and nonparticipants within program groups emerges, households from the preventive group who had ever participated in the program had a higher number of household assets compared to ever participants from the recuperative program communities. The same was true for differences between preventive and recuperative groups for nonparticipants - i.e., nonparticipants in the preventive group had higher numbers of household assets than nonparticipants in the recuperative group.

With respect to differences by current participation, ownership of productive assets was higher among current participants compared to nonparticipants in the preventive group. Differences in the recuperative group showed a similar trend but did not reach statistical significance. There were no differences in ownership of durable goods or of livestock assets, or in women’s ownership of assets between current participants and nonparticipants.

In summary,

- ❖ Program communities were not different in household or respondent asset ownership at baseline.
- ❖ Differences between program groups at the final survey were statistically significant, but small, for household durable goods and productive assets ownership. No differences were seen between program groups for ownership of livestock assets or for women’s asset ownership.

Table 10.1 Household and women's asset ownership, by program group, participation, and time of the survey

Household assets	Variable name	Baseline		Final Preventive		Final Recuperative		Final Preventive				Final Recuperative			
		(n = 1,524)		(n = 748)		(n = 752)		Nonparticipant		Participant		Nonparticipant		Participant	
		Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
(1) Ever participated															
Household assets (simple count)	hhasset	10.2	1.8	10.2	1.8	9.8**	1.9	(n = 126)		(n = 622)		(n = 130)		(n = 622)	
Productive assets (simple count)	prasset	2.9	1.4	3.1	1.6	3.0**	1.6	3.3 ^b	1.7	3.1	1.6	2.9	1.6	3.0	1.6
Livestock assets (simple count)	animal	2.4	1.6	2.2	1.6	2.2	1.6	2.2	1.7	2.2	1.6	2.3	1.7	2.2	1.6
Women’s ownership of assets (scale)	q803poss	2.2	1.1	2.1	1.0	2.1	1.1	1.9	1.1	2.2	1.0	2.0	1.1	2.1	1.1
(2) Current participation															
		(results as above)						(n = 423)		(n = 325)		(n = 621)		(n = 131)	
Household assets (simple count)	hhasset							10.1 ^b	1.8	10.2	1.8	9.8	1.9	10.0	2.0
Productive assets (simple count)	prasset							3.0 ^b	1.5	3.3 ^c	1.6	2.9	1.6	3.2 ^c	1.6
Livestock assets (simple count)	animal							2.2	1.6	2.3	1.6	2.3	1.6	2.0	1.6
Women’s ownership of assets (scale)	q803poss							2.1	1.0	2.2	1.0	2.1	1.1	2.0	1.1

** p < 0.05 for difference between program groups (random effects regression).

^a Difference between program groups is significant at p < 0.05 for participants (random effects regression analysis).

^b Difference between program groups is significant at p < 0.05 for nonparticipants (random effects regression analysis).

^c Difference between participants and nonparticipants (within program group) is significant at p < 0.05 (random effects regression analysis).

- ❖ At the final survey, household asset ownership was the same as baseline in the preventive communities but was lower than at baseline in the recuperative communities. There were no differences from baseline in either group for livestock ownership or women's asset ownership.
- ❖ In both groups, there were no differences in any of the assets variables between those who had ever participated in the program and those who had not.
- ❖ In the preventive communities, number of household productive assets owned was higher among households who were *current* recipients of program benefits than those who were not current recipients, which could indicate positive impacts of the program, even though the differences were generally of small magnitude. Conversely, this could suggest that, in fact, the poorest households are unable to participate in the program.

10.3.2 Indirect impact of the program on women's well-being

There were no differences between program communities in any of the measures of women's well-being at baseline, including self-rated overall health, life satisfaction, mental well-being, frequency of stress symptoms, and time related stress. At the final survey, however, respondents in the preventive communities were generally better off than those in the recuperative communities on the women's well-being measures; preventive community respondents reported greater life satisfaction than recuperative community respondents and scored lower on the mental stress symptoms scale ($p < 0.05$) (Table 10.2). Scores on almost all the women's well-being measures were also higher at the time of the final survey for both program groups compared to the baseline. For the mental stress scale, the mean score was unchanged since baseline for the preventive group, but was higher than baseline for the recuperative group, indicating a higher level of stress in that group at the time of the final survey.

There were no consistent patterns of differences between caregivers who ever participated and those who never participated; this was true for both program groups. The regression models that included the participation variables (shown only for self-rated health scores and mental stress in Tables 10.3 and 10.4) show that overall, current participation was associated with higher self-rated health scores and lower mental stress. There was no association between the program participation and well-being for any of the other four well-being measures (not shown). Also, there were no interactions between participation and program group, suggesting that the benefits of current participation did not differ between program groups. Finally, there was also no association between ever having participated and mental well-being measures, suggesting that the benefits of program participation were observed only in the short-term only.

Although the interaction term between program and current participation was not statistically significant in the regression model, the association between participation and well-being appears to be restricted to the preventive group. In the recuperative group, for example, there were no differences between current participants and nonparticipants on any of the well-being measures. However, current participants in the preventive group scored lower on the

Table 10.2 Respondent mental well-being, by program group, participation, and time of survey

Mental well-being	Variable name	Baseline (n = 1,524)		Final Preventive (n = 748)		Final Recuperative (n = 752)		Final Preventive				Final Recuperative			
		Mean	SD	Mean	SD	Mean	SD	Nonparticipant		Participant		Nonparticipant		Participant	
								Mean	SD	Mean	SD	Mean	SD	Mean	SD
(1) Ever participated															
Self-rated health (maximum = 10)	q901	6.1	1.8	6.8	2.2	6.6*	2.1	(n = 126)		(n = 622)		(n = 130)		(n = 622)	
Mental stress (scale) (maximum = 6)	q904ment	3.0	1.6	3.0	1.6	3.3**	1.6	3.2	1.8	2.9	1.6	3.2	1.5	3.3 ^a	1.6
Frequency of stress symptoms (scale) (maximum = 20)	q905hlth	10.1	3.5	9.3	3.9	9.5	3.8	9.7	3.7	9.2	4.0	9.6	3.7	9.5	3.8
Time stress (scale) (maximum = 8)	q905time	3.9	1.9	3.0	2.1	3.0	2.0	3.1	2.2	3.0	2.1	2.7	2.0	3.0	2.1
Life satisfaction (maximum = 20)	q903lsat	11.1	3.8	13.5	3.7	13.0*	3.9	13.0	3.8	13.6	3.7	12.5	4.1	13.1 ^a	3.8
Life satisfaction (without q903g & q903h) (maximum = 16)	q903sat2	n/a	n/a	9.7	3.1	9.3**	3.1	9.4	3.0	9.8	3.1	9.0	3.3	9.4	3.1
(2) Current participation															
		Same as above						(n = 423)		(n = 325)		(n = 621)		(n = 131)	
Self rated health (maximum = 10)	q901							6.6	2.2	7.1 ^b	2.1	6.6	2.2	6.8	1.9
Respondent mental health (scale) (maximum = 6)	q904ment							3.1	1.7	2.8 ^b	1.6	3.3 ^c	1.6	3.2 ^a	1.5
Frequency of stress symptoms (scale) (maximum = 20)	q905hlth							9.5	3.8	9.0	4.1	9.5	3.8	9.4	4.0
Time stress (scale) (maximum = 8)	q905time							3.1	2.1	2.8 ^b	2.1	3.0	2.0	3.0	2.1
Life satisfaction (maximum = 20)	q903lsat							13.3	3.6	13.6	3.8	12.8	4.0	13.7	3.4
Life satisfaction (without q903g & q903h) (maximum = 16)	q903sat2							9.6	3.1	9.9	3.1	9.3	3.1	9.4	3.1

Notes: ** p < 0.05 for difference between program groups; * 0.05 < p < 0.1 (random effects regression).

^a Difference between program groups is significant at p < 0.05 for participants (random effects regression analysis).

^b Difference between participants and nonparticipants (within program group) is significant at p < 0.05 (random effects regression analysis).

^c Difference between program groups is significant at p < 0.05 for nonparticipants (random effects regression analysis).

Table 10.3 Program impact on self-rated health (random effects regression)

	(1)	(2)	(3)	(4)	(5)	(6)
	Bivariate model	Main effects, adjusted	Main effects, ever participated	Interaction model, ever participated	Main effects, currently participating	Interaction model, currently participating
Dependent variable→	Self-rated health	Self-rated health	Self-rated health	Self-rated health	Self-rated health	Self-rated health
Program (reference group=preventive)	-0.18 (0.11)	-0.24 (0.12)	-0.24* (0.12)	0.04 (0.31)	-0.13 (0.13)	-0.08 (0.15)
Respondent's education (years)		0.05 (0.03)	0.05 (0.03)	0.05 (0.03)	0.05 (0.03)	0.05 (0.03)
Partner's education (years)		-0.03 (0.02)	-0.03 (0.02)	-0.03 (0.02)	-0.03 (0.02)	-0.03 (0.02)
Ever participated			0.25 (0.17)	0.42 (0.24)		
Ever participated and Program				-0.33 (0.34)		
Currently participating					0.39** (0.14)	0.47** (0.17)
Currently participating and Program						-0.19 (0.29)
Constant	6.81** (0.12)	6.83** (0.14)	6.62** (0.19)	6.48** (0.24)	6.66** (0.15)	6.62** (0.17)
Observations	1,500	1,250	1,250	1,250	1,250	1,250
Number of pairs	10	10	10	10	10	10

Notes: Standard errors in parentheses. * significant at 5%; ** significant at 1%.

Table 10.4 Program impact on mental stress scale (presence/absence of symptoms) (random effects regression)

	(1)	(2)	(3)	(4)	(5)	(6)
	Bivariate model	Main effects, adjusted	Main effects, ever participated	Interaction model, ever participated	Main effects, currently participating	Interaction model, currently participating
Dependent variable→	Mental stress scale	Mental stress scale	Mental stress scale	Mental stress scale	Mental stress scale	Mental stress scale
Program (reference group=preventive)	0.30** (0.08)	0.31** (0.09)	0.31** (0.09)	0.13 (0.23)	0.24* (0.10)	0.23* (0.11)
Respondent's education (years)		-0.04* (0.02)	-0.04* (0.02)	-0.04* (0.02)	-0.04* (0.02)	-0.04* (0.02)
Partner's education (years)		-0.00 (0.02)	-0.00 (0.02)	-0.00 (0.02)	-0.00 (0.02)	-0.00 (0.02)
Ever participated			-0.02 (0.13)	-0.12 (0.18)		
Currently participating					-0.26* (0.10)	-0.27* (0.13)
Ever participated and Program				0.21 (0.25)		
Currently participating and Program						0.04 (0.22)
Constant	2.95** (0.07)	3.02** (0.10)	3.04** (0.15)	3.12** (0.18)	3.14** (0.11)	3.14** (0.13)
Observations	1,500	1,250	1,250	1,250	1,250	1,250
Number of pairs	10	10	10	10	10	10

Notes: Standard errors in parentheses. * significant at 5%; ** significant at 1%.

mental stress symptoms than current participants in the recuperative group (Table 10.2; $p < 0.05$).

In summary,

- ❖ There were no differences between program communities at baseline on any of the women's well-being measures.
- ❖ At the time of the final survey, respondents in the preventive communities were better off than those in the recuperative communities on four of the women's well-being measures.
- ❖ There were no differences between respondents who had ever been exposed to the program and respondents never exposed to the program. However, current participants in preventive communities had better self-rated health, lower mental stress, and lower time stress than nonparticipants.

10.4 Conclusions

This chapter presented findings that point to the indirect benefit of participation in the program on some aspects of household asset ownership and on women's well-being. We explored these potential benefits based on the strong associations between food insecurity and these outcomes at baseline. Our analysis was grounded in the premise that enrollment in the targeted components of the program (the food assistance, in particular) could alleviate the impact of food insecurity on these outcomes, and thereby reveal indirect benefits of the program.

Our findings suggest that the preventive program communities had slightly higher household asset ownership than the recuperative communities at the end of the evaluation period. There was no change in household asset ownership since baseline in either group, which suggests that food assistance had limited if any effect on asset building.

With regard to maternal health and well-being, four of six well-being measures were better in the preventive group at the time of the final survey. For all but one of the well-being measures, improvements were seen in both program groups since baseline, suggesting a potential benefit of the program inputs on some aspects of women's well-being. Overall, current participation was positively associated with self-rated health and negatively associated with overall mental stress, suggesting a short-term benefit of the program on these outcomes.

Given the conditional nature of the WV MCHN program, it is impossible to tease out whether the benefits of participation seen in the study are linked to the food inputs or the supportive Mothers' Club environments that women participate in. Nevertheless, these results offer an insight into the potentially wide ranging benefits of Title II program inputs in this severely food insecure and impoverished environment.

11. COST-EFFECTIVENESS OF THE PREVENTIVE APPROACH RELATIVE TO THE RECUPERATIVE APPROACH²⁷

11.1 Introduction

In this chapter, we describe and estimate the cost-effectiveness ratio (CER) combining information on the cost structure of the program with the design of the evaluation. Maluccio and Loechl (2006) show that the number of beneficiaries and the time that each one remains in the program drive the differences in costs between the recuperative and preventive program approaches, since the programs are otherwise nearly identical. We exploit this basic feature to calculate the difference in costs across the two program approaches for respondents in the final survey (2005). In doing so, we are able to link cost differentials to the effectiveness calculations presented in Chapter 7.

Given the design of the programs and their multiple potential benefits, particularly improved nutritional status for children that is likely to yield nutritional, health, and economic returns over many years, we caution that it would be incorrect to interpret the CER as a measure of the cost-benefit ratio (CBR) of the program. This is in part because we do not assign a monetary value to the short-term effect of the program on reduced malnutrition. It is also because we neither delineate nor value the potential longer-term benefits due to the improved nutritional status of this group of young Haitians. Finally, in this chapter we consider only a limited number of outcomes, pertaining to the primary program objectives. In other chapters, this report explores some of the other many possible benefits provided by the program, for example, in terms of increased food security of the household, improvements in maternal knowledge, exclusive breastfeeding, and long-term behavioral changes. All of these would need to be included and valued to calculate the CBR.

The evaluation compares the recuperative and preventive program approaches, without the use of a control group. With randomization, this is a strong design, which can answer the main question of the study, i.e., whether the preventive approach is more effective than the traditional recuperative approach of targeting food aid in the context of the MCHN program.

The lack of a control group not receiving either program approach, however, has important implications for the cost-effectiveness analyses. With this design, the analytical approach we must follow is a differential one, typically referred to as a *relative* or *incremental* cost-effectiveness analysis (Jamison et al. 2006; Musgrove and Fox-Rushby 2006). The analysis of the impact of the two program approaches on the main study outcomes - i.e., child nutritional status - explored the effect of the preventive program approach *relative* to the recuperative program approach, i.e., what would happen if one “switched” from the recuperative to the preventive. We must take the same approach to compare the cost-effectiveness of the two program approaches. We examine changes in costs as one “switches” from the recuperative program approach to the preventive, and then associate any additional costs with the gains in effectiveness resulting from that same change.

²⁷ John Maluccio led the writing and analysis for this chapter. Further details are available in Maluccio and Loechl (2006).

11.2 Methodology

11.2.1 The incremental cost-effectiveness between the two program approaches

To measure the incremental cost-effectiveness between the program approaches, we calculate

$$CER = \left(\frac{C_p - C_r}{I_p - I_r} \right),$$

with

- C_p = cost of the preventive program,
- C_r = cost of the recuperative program,
- I_p = impact of the preventive program on a given outcome, and
- I_r = impact of the recuperative program on a given outcome.

As described in Section 11.1, because there is no control group we do not separately observe I_p and I_r , but rather only the difference, $(I_p - I_r)$.

For the numerator, we use comprehensive cost measures that include direct program costs included in the World Vision-Haiti DAP accounting system, other program costs outside the WV-Haiti DAP accounting system, and beneficiary opportunity costs. These are described in the next subsection.

For the denominator, we consider three alternatives, the number of cases prevented of stunting, underweight, and wasting, by changing from the recuperative to the preventive program approach. This approach has the advantage of making the analysis more comparable to existing studies (e.g., Waters et al. 2006). Cases prevented are estimated from the final survey (2005) as the difference in the number of cases of, e.g., of stunting, between the two program approaches.

11.2.2 Measuring the costs

This section describes the three major components of the costs of the program. These include (1) direct program costs; (2) outside program costs; and (3) beneficiary opportunity costs. In this study, we consider direct program costs to be those program costs funded by the monetization of USAID commodities. Outside program costs are the costs of Title II food commodities (provided by USAID) and health-care supplies (provided by MoH-Haiti). Opportunity costs are the costs, real and opportunity costs, incurred by program beneficiaries for using services. In this section we define all the cost components and how we estimate them and in Section 11.3 we calculate them for the program evaluation areas.

Table 11.1 provides an overview of the cost components and the data sources used to estimate them.

Table 11.1 Variables and data sources for incremental cost-effectiveness analysis

Variable	Definition	Data source
Direct program costs	❖ Program costs funded by USAID through wheat monetization and some cash grants	❖ WV-DAP accounting system
Outside program costs	❖ Costs of Title II commodities (from USAID) ❖ Health-care supplies (from MoH, Haiti)	❖ Commodities division monthly distribution summaries, shipping and price records ❖ MCHN division monthly reported beneficiaries x estimated costs of supplies
Beneficiary opportunity costs	❖ Costs to beneficiaries of fulfilling program requirements (includes potentially lost earnings)	❖ Time spent estimated from operations research 2005 (Loechl et al. 2004) & 2002 baseline survey (Menon & Ruel 2003) ❖ Haitian Living Standards Measurement Survey (LSMS) – used to value time
Number of beneficiaries and beneficiary-months	❖ Number of beneficiaries served by each program approach (as beneficiary-months & beneficiaries per month)	❖ Commodities division monthly distribution summaries
Average length in program	❖ Average number of months enrolled in program	❖ Final survey (2005)
Cases of undernutrition prevented	❖ Difference between preventive and recuperative approaches in the numbers of stunted, underweight, and wasted children	❖ Final survey (2005)
Cases of undernutrition prevented	❖ Difference between preventive and recuperative approaches in the numbers of stunted, underweight, and wasted children	❖ Final survey (2005)
Cumulative number of beneficiary months	❖ Total number of beneficiary months of program enrollment (calculated by summing up the total months of participation for all children in each of the program groups)	❖ Final survey (2005)

11.2.2.1 Direct program costs

USAID funds the vast majority of WV-Haiti DAP activities, principally via provision of food commodities which are either monetized or distributed as part of the program. In Haiti, wheat is the food commodity monetized by WV-Haiti to pay for program operations. Typically, the primary source of information for the direct program costs is the program's accounting system. An accounting-based approach to measuring these costs is possible in this study because, although WV-Haiti operations for these two programs do not operate in a completely autonomous fashion, the majority of program-related activities is carried out under the WV-Haiti

Development Activity Program (DAP) and is therefore under its accounting system.²⁸ All direct program costs, funded by wheat monetization and to a lesser extent cash grants, are captured by the WV-Haiti DAP accounting system. We make a number of adjustments to the raw accounting information from the program (further described in Maluccio and Loechl 2006). We adjust all figures by the U.S. inflation rate and report them as 2005 constant U.S. dollars. These costs do not vary on the margin (e.g., when adding an additional beneficiary) and therefore will be treated as *fixed costs* in this study.

11.2.2.2 Outside program costs

Next, we estimate “outside” program costs for WV-Haiti, important program costs that are outside the WV-Haiti DAP accounting system. There are two principal items in this category: Title II commodities provided by USAID that are distributed (i.e., not monetized) and health-care supplies provided by the Ministry of Health in Haiti. Because these costs turn out to be the key *variable costs* underlying the difference between program approaches, we describe them in some detail.

11.2.2.2.1 USAID Title II food commodities

As part of its reporting requirements to USAID, WV-Haiti has an elaborate computerized tracking system designed for the DAP food commodities (known as the commodity tracking system or CTS), which is a system used by WV worldwide. This system tracks commodities from the moment they enter the country until they are distributed to the beneficiaries, documenting movements, amounts distributed, and any losses. Using this system, the Commodities division is able to report numbers of beneficiaries and quantities distributed to each of them. USAID pays to transport the food commodities to Haiti; upon arrival at the port, subsequent expenses for transport and warehousing are covered by WV-Haiti under the DAP accounting system, reflected under the commodities division activities.

The food commodity-related costs not reflected in the WV-Haiti DAP accounting system, then, include the value of the food commodities ultimately transferred to households and the cost of their shipment to Haiti. We use shipping and price records provided by WV-Haiti to calculate the mean value per kilogram, including external shipping costs, for each food commodity for each fiscal year (FY).²⁹ This does not necessarily equal the market price of these items in Haiti if one were to purchase them on the private market there.³⁰ We choose to value food

²⁸ This approach is not always possible. For example, Fiedler (2003), in a cost analysis of a Honduran community-based integrated childcare program that did not have a centralized accounting system, constructs total program costs from the bottom up, estimating the costs required for each “ingredient” activity and then aggregating them. This is a valid approach, also recommended by Adam (2006), and was considered in the design of this study. It allows useful simulations of costs under varying program designs (e.g., excluding certain components) that may more closely approximate marginal costs. An important drawback to the bottom-up approach, however, is that it is difficult to capture *all* of the activities and associated costs borne in the central office of the program. Our view, supported by Waters (2000), is that it would have likely led to an underestimate of the full program costs.

²⁹ These are known as cost, insurance, and freight (CIF) prices as opposed to free on board (FOB) prices, which include only the cost of the items being shipped.

³⁰ It is likely a small secondary market in these food commodities exists, though we have no hard evidence about its depth or the prices for the commodities not commonly available elsewhere, such as Wheat-Soy blend (WSB).

commodities in this way as it is the most relevant in terms of the resources being devoted to the program from a *global* perspective, even though it is possible that the local value of the commodities is more or less than this value. We then use these calculated values to assess the aggregate value of food commodities delivered under the program.³¹ This includes direct rations given to child beneficiaries and their associated family rations, as well as direct rations given to pregnant and lactating mothers and their associated family rations.

This valuation approach differs from the way in which we assess the value of wheat commodities that are monetized and used for direct program costs (described above), which essentially takes the Haiti market valuation. Wheat is a global commodity traded in large volumes, so that local prices are more likely to reflect global prices (plus transport). Nevertheless, monetization results reported in WV-Haiti annual reports suggest that cost recovery does not reach 100%; in FY05, for example, it was estimated at 96%, suggesting we are underestimating slightly the resources required for direct program costs. As made clear in the analysis below, however, the funds used from monetization of wheat fall into the fixed costs categories and therefore do not affect the calculation of the incremental CER, although they would change slightly the overall program costs we report.

11.2.2.2 Health-care supplies

The next important component of outside program costs is the provision of health-care supplies such as vaccines (for both children and women), vitamin A capsules, iron folate supplements, oral rehydration salts, and deworming pills. The costs for delivery (and administration) of these items already are included in the operational costs of the program; it is only the supplies themselves that are outside the WV-Haiti accounting system.

Using MCHN division monthly reports that detail all the services (listed above) provided at Rally Posts (RPs) to program beneficiaries (both children and pregnant and lactating women), as well as to others who attend and receive health-care services but are not eligible for the food commodities, we calculate the total number of persons receiving health-care supplies in each FY. We combine that information with estimates of the costs of each of the components (e.g., the unit cost of each vaccine) to compute the aggregate cost of the in-kind health-care supplies.³² The dominant component (representing over half the health-care supply costs) was the iron folate tablets provided to pregnant women.

11.2.2.3 Beneficiary opportunity costs

While our information is most complete for the direct and outside program costs described above, there are other potential costs, such as beneficiary private costs of participation, that result from the introduction of the program. Failing to pay attention to them may severely underestimate the full program costs.

³¹ Food costs per kg over the four years are (1) WSB: \$0.41–0.46; (2) SFB: \$0.23–0.26; (3) Vegetable oil: \$0.32–0.87; and (4) Lentils: \$0.81–0.96.

³² The price data we use were collected from the UNICEF office in Haiti and pertain to 2004. As with commodity prices, there is a concern about whether these prices reflect true social valuations. This possibility is unlikely to affect substantially the results, however, as it turns out that medical supplies comprise a very small portion of the total value of outside program costs.

It turns out that out-of-pocket private costs for beneficiaries are negligible. Instead, the important beneficiary private costs that stem from the program are opportunity costs incurred by beneficiaries to complete program requirements. Program beneficiaries may incur costs as a result of the program in several ways. The mothers of child beneficiaries must, as a condition to receive the food transfers, attend the Mothers' Clubs (MCs), bring their children to the RPs (although it is permissible for another caregiver to take the child to the Rally Post), and travel to the Food Distribution Points (FDPs) (although it is permissible for any member of the family to carry out this last requirement). Similarly, women beneficiaries (pregnant and lactating women) must attend pre- and postnatal consultations (instead of RPs), as well as MCs and FDPs. If they do not, they cannot participate in the program (and therefore do not receive food transfers) - hence these are necessary and possibly additional, costs that they undertake in order to remain program beneficiaries.

Based on the operations research, Loechl et al. (2004) analyze time spent by mothers of child beneficiaries to fulfill program requirements at the three most important contact points (MCs, RPs, FDPs). We combine that information with information available in the final survey (2005), and estimate that, on average, total time commitments are approximately 12 hours per month per beneficiary.³³ A comparison of whether there were substantial differences in monthly time costs between preventive and recuperative areas indicated there were none, other than that average travel time was about 20 minutes greater in recuperative areas. Mothers who in addition to having a child in the program are themselves beneficiaries (pregnant or lactating with a child < 6 months old) have to fulfill *both* sets of program requirements. They have to attend an additional Mothers' Club and a pre- or postnatal consultation once per month, but only need attend the Food Distribution Point once. The final survey (2005) indicates that this is uncommon, occurring in less than 2% of cases.

About 85% of caregivers in the baseline survey (2002) reported being involved in income-generating activities in the past year (Menon and Ruel 2003). For these women, it is obvious why we should value their time - they may have had to give up remunerative activities in order to attend the program activity. This can occur despite laudable efforts by WV-Haiti to plan events to avoid overlap with important income-generating activities, such as local market days. Even for those women who did not lose earnings or did not report working, however, we must still value their time spent in complying with program requirements.

To value women's time, we begin with the 2001 Haitian Living Standards Measurement Survey (LSMS) and calculate daily earnings for rural women who live and work in Central

³³ The calculations from the operations research were (1) FDPs: average time to and from was 58×2 minutes and average time there was 241 minutes; (2) RPs: average time to and from was 20×2 minutes and average time there was 117 minutes; (3) MCs: average time to and from was 18×2 minutes and average time there was 40 minutes (waiting) and 66 minutes (in session) (Loechl et al. 2004). We did not assess the time spent by pregnant and lactating women at the pre- and postnatal consultations, but we assume that the time commitments are similar to those for Rally Post attendance (average travel time and time spent at the venue). In the final survey (2005), we replicated the travel time questions to get responses for a more representative population and found average travel time to (1) FDPs: 84×2 minutes; (2) RPs: 29×2 minutes; and (3) MCs: 39×2 minutes. Combining the operations research times for non-travel components reported in Loechl et al. (2005) with the travel times from the household survey yields 768 minutes, which we round off to approximately 12 hours. The operations research also revealed it was rare that women paid for transportation, thus no such costs are included.

Plateau.³⁴ In 2001, the median daily wage for women working in rural areas of the Central Plateau was 44 gourdes, which, at 2001 exchange rates, was a little under \$2 a day. As the subset of women who work is not random, we recognize that this valuation likely overstates what many women could and do earn. This is also likely since from the LSMS, we consider only those women working in wage labor. Most women in the program areas are not working in the formal sector and typically informal sector jobs are less remunerative. Finally, to the extent that women can rearrange and substitute their activities across time, it would be possible for many of them not to lose income as a result of having to fulfill the program requirements.

11.3 Data: Costs in the Program Areas

This section presents estimates of the program (11.3.1) and opportunity (11.3.2) costs in the evaluation areas. We use the cost components described above and calculate these costs for the subset of areas in which the program approaches being evaluated were implemented. Further details about the cost calculations are available in Maluccio and Loechl (2006).

11.3.1 Direct and outside program costs in the study areas

The two program approaches compared in the evaluation were similar in most activities and services. The two main differences are

- the number of program beneficiaries and
- the potential length of time each beneficiary remains in the program.

The first element is a function of the number of malnourished children under 5 compared with the number of children 6–24 months old and the utilization rates for those two groups. The second is a difference in design across the two programs. In the preventive program approach, children potentially remain in the program longer, on average, and they and their families receive the same food ration per month as in the recuperative approach. A child entering at 6 months of age is eligible to remain in the program for up to 18 months. In step with this longer eligibility period for the children, mothers under the preventive approach potentially attend the MCs for a longer period of time and children are required to attend the RPs for a longer period. Pregnant and lactating women are treated the same under both approaches. Since the food commodities are the dominant expense and these are very similar for this beneficiary group, this is unlikely to affect the differential costs calculated below. An obvious way to distribute the *variable* costs associated with each approach, then, is to base them on the relative distribution of program beneficiaries per month, as the size of the food rations (per month) are the same across the two program approaches. As part of the study, in addition to its regular tracking of all beneficiaries, WV-Haiti tracked the number of program beneficiaries per month in the study areas.

³⁴ Given the already complex nature of the household surveys for this study, we chose not to add questions about earnings to them knowing we could rely instead on the Haitian LSMS.

11.3.1.1 Differences in beneficiary months

Table 11.2 presents trends for child and pregnant and lactating women beneficiaries for all of Central Plateau as well as separately for the study areas where the two program approaches operate, based on the Commodities division monthly distribution summary reports. Each child or pregnant or lactating woman is counted once for each month he or she is in the program. Thus, the table represents “beneficiary-months” (and not number of children or number of pregnant and lactating women).³⁵ We use extensively in what follows the concept of a beneficiary-month. Since beneficiaries can benefit for multiple months (as described above), the following month a number of the beneficiaries are the same persons, receiving another month of services and therefore representing another beneficiary-month. This measure is the most relevant for the cost analysis. Later, we consider how the results change when we consider instead the raw number of beneficiaries (as in number of different persons) served.

Table 11.2 Number of program beneficiary-months in Central Plateau and study areas

	FY 2002	FY 2003	FY 2004	FY 2005	Total
Children					
Central Plateau	29,375	73,765	86,632	110,371	300,143
Study area: Preventive	508	6,227	11,195	14,000	31,930
(percent of Central Plateau)	(1.7)	(8.4)	(12.9)	(12.7)	(10.1)
(percent of study area)	(52.9)	(56.2)	(69.4)	(66.2)	(64.7)
Study area: Recuperative	453	4,861	4,941	7,147	17,402
(percent of Central Plateau)	(1.5)	(6.6)	(5.7)	(6.5)	(5.8)
(percent of study area)	(47.1)	(43.8)	(30.6)	(33.8)	(35.3)
Study area: Total	961	11,088	16,136	21,147	49,332
(percent of Central Plateau)	(3.3)	(15.0)	(18.6)	(19.2)	(16.4)
Children and pregnant/lactating women					
Central Plateau	48,188	130,816	160,274	195,046	534,324
Study area: Preventive	1,008	8,727	17,497	21,245	48,476
(percent of Central Plateau)	(2.1)	(6.7)	(10.9)	(10.9)	(9.1)
(percent of study area)	(51.4)	(54.2)	(60.9)	(59.6)	(58.8)
Study area: Recuperative	953	7,362	11,242	14,393	33,949
(percent of Central Plateau)	(2.0)	(5.6)	(7.0)	(7.4)	(6.4)
(percent of study area)	(48.6)	(45.8)	(39.1)	(40.4)	(41.2)
Study area: Total	1,961	16,088	28,739	35,638	82,426
(percent of Central Plateau)	(4.1)	(12.3)	(17.9)	(18.3)	(15.4)

Source: WV-Haiti commodities division and authors' calculations.

Notes: Each child or woman is counted as a beneficiary for every month he or she is in the program. Thus, the table represents beneficiary-months (and not number of children or number of pregnant and lactating women). The number of pregnant and lactating women beneficiary-months in the study areas is estimated as described in Maluccio and Loechl (2006).

³⁵ From the final survey (2005) the average duration in the program for children in the preventive approach was 11.7 months and for the recuperative approach, 7.5 months. Combining this information with the number of beneficiary-months presented in Table 11.2, we estimate the total number of beneficiary children over the four years to be 2,729 in preventive areas and 2,320 in recuperative areas.

Over time, the combined study areas accounted for an expanding fraction of the program child beneficiary-months and, to a lesser extent, pregnant and lactating women beneficiary-months, in Central Plateau, reaching nearly one-fifth by FY 2005. This is consistent with the fact that 12 FDPs serve the study areas, nearly one-fifth of the 68 FDPs that serve the Central Plateau region as a whole.

When the program first began in late 2002, the number of child beneficiary-months across the two programs was small and approximately equal, possibly because information about the eligibility for the preventive approach had not yet become widespread. The steady increase in the share of beneficiaries in the study areas is due to increases in both preventive and recuperative areas, but the increases in preventive areas have been much larger, such that in FY 2005, about two-thirds of the child beneficiary-months in the study areas were in preventive areas. The trends seen in these program data agree closely with data from the 2005 household census carried out in the study areas; those data indicate 70% of the current child beneficiaries are living in preventive areas. They also agree with information collected in the final survey (2005), which shows 73% of current child beneficiaries are in preventive areas. The 2005 household census also confirms that the percentages of pregnant and lactating women in each of the areas are similar, and pregnant and lactating women beneficiaries are split evenly between the program approaches, a pattern also seen in Table 11.2.

Lastly, the 2005 household census confirms that the emerging difference in child beneficiaries served is not due to differences in sizes of the populations of the areas, each of which is approximately 20,000 individuals. Instead, it is a function of (1) the difference in the number of 6–24-month-old children participating in the program compared with the number of malnourished children under 5 participating in the program (which, in turn, is a function of the malnutrition rate and the participation rates within each of those groups); and (2) the deliberate design feature of the program that children under the preventive approach potentially remain in the program up to twice as long.

When we examine annual data such as these, the two effects are present since it is possible for a child in the preventive area to show up in 12 consecutive months in the FY but for a malnourished child to show up at most 9 months in that year (although it is possible for a malnourished child to reenter the program later). Even if the same number of beneficiaries were in each program, beneficiary-months in the preventive program could be one-third higher due only to the difference in eligibility periods. Another way to see the extent to which the preventive program has more beneficiaries is to look on a month per month basis. This better isolates the first difference mentioned above, i.e., the comparison between the number of children participating in the 6–24-month age range in preventive areas against the malnourished participants under 5 in the recuperative areas, since a child can appear as a beneficiary only once in any given month. For the study areas, Table 11.3 presents these figures for the final months of FY 2005. While there is fluctuation on a monthly basis, these figures show similar patterns to those in Table 11.2, with only a slightly lower percentage of beneficiaries in preventive areas compared to Table 11.2. This suggests that the annual figures are not inflated greatly by the possibility of participation for 12 versus 9 months in preventive versus recuperative areas. In other words, the higher beneficiary-month figures in Table 11.2 primarily reflect more actual beneficiaries and not just longer periods of participation.

Table 11.3 Number of program child beneficiaries per month in the Central Plateau and study areas (FY 2005)

	April	May	June	July	August
Children					
Study area: Preventive	1,159	1,022	1,163	1,098	1,114
(percent of study area)	(62.9)	(57.4)	(63.7)	(61.9)	(68.3)
Study area: Recuperative	685	760	663	677	516
(percent of study area)	(37.1)	(42.6)	(36.3)	(38.1)	(31.7)
Study area: Total	1,844	1,782	1,826	1,775	1,630

Source: WV-Haiti commodities division and authors' calculations.

Notes: Each child or woman is counted as a beneficiary for every month he or she is in the program.

11.3.1.2 Differences in total program costs

The program beneficiary data show that the study program grew substantially over the three years. Program costs also rose, though not as dramatically, consistent with economies of scale as the program grew. The estimate for full (direct and outside) program costs in the Central Plateau from October 2001 to September 2005 was \$16 million. In FY 2005, the costs were just over \$5 million, fully 40% of which was the value of food commodities distributed to beneficiaries. Ignoring the value of the food commodities distributed would underestimate the program costs substantially. The value of health-care supplies, on the other hand, appears to be less significant in the overall program costs (less than 1%). Another 25% of the full program costs (40% of direct program costs) was for the commodities division of WV-Haiti, in charge of food distribution. The next largest cost centers were the support (administrative) division with 15% and the MCHN division, which carries out the health-care services, at 12% of full (on- and off-budget) program costs (Maluccio and Loechl 2006).

In Maluccio and Loechl (2006), the various program costs for MCHN in Central Plateau were estimated using accounting and other information provided by WV-Haiti. After capturing and valuing as best possible the full range of program costs feeding into the overall program, we isolated the costs that pertain *specifically* to the study areas and, within these areas, the costs that pertain to each approach.³⁶ Only when this is done are the measures of costs and effectiveness on the same basis, allowing assessment of the incremental CER.

The overall program cost figures for the study areas are shown in Table 11.4. Each of the program approaches operate in similarly sized areas that present similar difficulties in terms of access and other infrastructure. They each operate with the same management oversight from WV-Haiti and thus differ little in terms of staff. For example, they have the exact same number of health agents working for the two program approaches, although about five additional low paid *colvols* serve preventive areas. The areas have approximately the same number of RPs and

³⁶ We use the percent of program beneficiary-months in the study areas of all program beneficiary-months in Central Plateau to assign costs to the study areas. Then, we use the relative fraction of beneficiary-months between the two program approaches to allocate the total costs within the study areas (calculated in the previous step) to each of the approaches. We emphasize that this top-down approach, which includes a number of reasonable, but nonetheless ad hoc, assumptions, is imperfect. Nevertheless, it captures the broad patterns of growth and development of the study program over time quite well. Further, it distinguishes between the two approaches based on the principal factor driving differences in their variable costs: the number and length of time that program beneficiaries receive food commodities. For further details, see Maluccio and Loechl (2006).

FDPs - indeed the latter are shared across preventive and recuperative areas. Therefore, we divide all direct program costs that do not vary with the number of beneficiaries (at least in the short term) equally between program areas. These expenses include salaries for all personnel and overhead costs, but importantly do not include the outside program costs - food commodities and health-care supplies. For this reason, the direct program cost rows are identical for the two program approaches in Table 11.4. While in part an assumption, given the similarities across program areas, it is a reasonable one, so that while we do not believe that they are exactly equal, we are confident that they are approximately equal. Table 11.4 also shows the breakdown of these costs by division (but only for the study areas as a whole).

Table 11.4 Direct program and outside program costs in the study areas (\$000)

	FY 2002	FY 2003	FY 2004	FY 2005	Total
Preventive					
Direct program costs	30.7	153.0	272.7	261.0	717.4
Outside program costs					
Food commodity costs	10.6	106.1	195.4	241.4	553.5
Health-care supply costs	0.5	3.7	3.8	1.9	10.0
Total	41.8	262.8	471.9	504.3	1,280.9
Recuperative					
Direct program costs	30.7	153.0	272.7	261.0	717.4
Outside program costs					
Food commodity costs	10.0	89.5	125.5	163.6	388.6
Health-care supply costs	0.5	3.1	2.5	1.3	7.4
Total	41.2	245.6	400.7	425.9	1,113.4
Total Study Area					
Direct program costs	61.4	306.0	545.4	522.0	1,434.8
% of which					
Support division	40%	33%	29%	28%	32%
MCHN division	20%	18%	26%	26%	23%
Commodity distribution division	40%	49%	45%	46%	45%
Outside program costs					
Food commodity costs	20.6	195.7	320.9	405.0	942.2
Health-care supply costs	1.0	6.8	6.3	3.3	17.4
Total	83.0	508.4	872.7	930.3	2,394.3

Source: WV-Haiti Commodities group and authors' calculations.

Notes: Figures are in 2005 constant U.S. dollars except where percent noted.

From the start of the project through September 2005, the full (direct and outside) program costs for the study interventions were \$2.4 million. In the first full year of operations (FY 2003), the full costs across the program approaches were roughly similar, but by FY 2004 the ratio of costs was moving in the direction of the beneficiary-month statistics: food commodity costs in preventive areas in FY 2004 and FY 2005 were 1.5 times those in recuperative areas, reflecting the beneficiary numbers shown in the bottom panel of Table 11.2. Direct program costs form the largest share of the costs, starting at 75% of costs in FY 2002 but declining in importance to just over 55% in FY 2005 as food commodity distribution expanded. Throughout the period, the cost of health-care supplies remains a minor component compared to the other costs. Food commodity costs are more than 98% of the total outside program costs.

11.3.1.3 Differences in program costs per beneficiary-month

At this stage, we are in a position to calculate the average costs per beneficiary-month. Our methodology for computing costs, based on the identical services offered to beneficiaries under both program approaches, means that the direct program costs per beneficiary-month differs across the program approaches, an effect consistent with the economies of scale of operations. The difference in costs across program approaches lie in the *number* of beneficiaries (or more accurately, beneficiary-months), and therefore the (variable) outside program costs per beneficiary. A complication in assessing the direct program costs per beneficiary-month is that while similar, the component parts (and therefore associated costs) of services and goods for child beneficiaries is different from that of pregnant and lactating women beneficiaries. Treating the two as equal, however, we can estimate direct program costs per beneficiary-month by dividing the figures in Table 11.4 by their corresponding beneficiary-month levels (for children and pregnant and lactating women) in Table 11.2. These results are shown in Table 11.5. To calculate the outside program costs (for food commodities and health-care supplies) per beneficiary-month, i.e., those that vary directly with the number of beneficiary-months, we employ the same strategy used to calculate the direct program costs per beneficiary-month, that is, divide the costs by the number of beneficiary-months.

Table 11.5 Direct program and outside program costs per beneficiary-month in the study areas (\$)

	FY 2002	FY 2003	FY 2004	FY 2005	Average
Direct program costs					
Preventive approach	30	18	16	12	15
Recuperative approach	32	21	24	18	21
Outside program costs	11	13	12	11	12

Source: WV-Haiti Commodities group and authors' calculations.

Notes: Figures in 2005 constant U.S. dollars.

Consistent with increasing program efficiency over time and economies of scale, direct program costs per beneficiary-month decline under both program approaches after the first year, and again in FY 2005. This decline occurs as the “fixed” overhead and central office-type expenses are spread over larger numbers of beneficiaries (or more precisely, beneficiary-months), making average costs lower. The decline is steeper in preventive areas, where there were greater increases in beneficiary-months. Outside program costs per beneficiary-month, on the other hand, do not decline dramatically. This is also expected, since the services are unchanging and these components most closely resemble variable costs. In FY 2005, when the program has had time to both grow and mature, outside program costs are \$11 per beneficiary-month (98% of which is for food), and full (direct and outside) program costs are \$23 in preventive areas and \$29 in recuperative areas.

11.3.2 Opportunity costs in the study areas

Valuing the 12 hours of time spent in required program activities at approximately one-day's wage (see Section 11.2.2.3), we approximate beneficiary opportunity costs of the interventions at \$2 per beneficiary per month. We consider this an upper bound for the private costs for the reasons indicated above. For the study area as a whole, then, this translates into

approximately \$170,000 over the course of the evaluation period, or 7% of the total costs reported in Table 11.4. While the estimate of the value of women's time turns out not to be a major cost component relative to the program as a whole, it is important to recognize that this is a cost that varies with beneficiary-months similar to the cost of food commodities.

11.4 Cost-Effectiveness of Preventive versus Recuperative

11.4.1 Incremental cost-effectiveness estimates

We use data from the final survey (2005) to estimate the incremental cost effectiveness of the preventive approach relative to the recuperative approach. The differential effectiveness and costs are estimated as follows:

Difference in program effectiveness: The differential effectiveness measures are also derived from the final survey (2005), which sampled children 12–41 months old. We calculated the simple difference between the preventive and recuperative survey samples in the number of children stunted, underweight, and wasted, thus giving us three measures of differential effectiveness. We suggest that this difference in number of cases of undernutrition between the two approaches is the number of *cases prevented* by the preventive approach relative to the recuperative approach. Table 11.6 presents the number of cases prevented of each of the anthropometric outcomes.

Difference in program costs: What is the difference in costs between the recuperative and the preventive approaches? Above we argued that the driving cost difference is the number of beneficiaries or, more accurately, beneficiary-months and the consequent outside program costs associated with each beneficiary-month. In other words, we treat these as variable costs, varying directly with the number of beneficiary-months. All other costs are treated as fixed and the same across the two program approaches. From the final survey (2005), we are able to calculate the cumulative number of child beneficiary-months in each of the program areas for the sample of children in the cross-section, since for each child we have the number of months he or she was a program beneficiary. For the same sample population of 1,500 children 12–41 months old (approximately 750 in each group) for whom the impact estimates described above have been calculated, they are

- Preventive areas: 6,422 beneficiary-months (for 547 beneficiary children);
- Recuperative areas: 1,589 beneficiary-months (for 212 beneficiary children).

In Tables 11.2 and 11.3, we saw that according to the program data, the preventive program tended to have about two-thirds of the total beneficiary-months on an ongoing basis, due to the combination of participation and average length of participation (see Chapter 6). When we examine the sample of children 12–41 months of age in the final survey (2005), preventive areas comprise 80% of the total number of beneficiary-months. As described above, differences between the groups are due to who is targeted, who participates, and for how long they participate. When we examine a cross-sectional survey containing children 12–41 months of age, however, in the preventive areas we are including a large group over 24 months of age, i.e., those potentially in the program for a full 18 months, and we count all the costs associated

with their participation. Those children in recuperative areas, on the other hand, would only be eligible if they were malnourished at some point, and even then only for 9 months (although with the possibility of re-entry into the program later if they remain or become again malnourished). Only the very youngest (12–15 months) in preventive areas were eligible for less than 9 months. Thus in the cross-section it is unsurprising that we see an even higher percent of beneficiary-months in preventive versus recuperative approaches when we examine the cumulative history of beneficiary-months.

To assess the increase in costs as one goes from the recuperative to the preventive program approach, the direct program costs “drop out” of the equation because they are the same for both programs. Ignoring beneficiary opportunity costs for the moment, we are left only with the outside program costs for food commodities and health-care supplies, the bottom row of Table 11.5. We underscore here that because of the similarity in programs, there appear to be only negligible cost differences between the approaches, apart from those costs that vary with the number of beneficiary-months. Below we reconsider the valuation of these variable costs, particularly food commodity costs.

The additional program costs associated with the preventive program, then, are the number of additional beneficiary-months multiplied by the average outside program costs per beneficiary-month, which we treat as the variable costs, approximately \$56,000.³⁷

Incremental cost-effectiveness

Using the difference in costs of \$56,000 and the differential program effects shown in Table 11.6, we can now calculate the incremental cost-effectiveness per case of the preventive versus recuperative program approaches, shown in the third column of Table 11.6 after rounding off to the nearest 100.

Table 11.6 Incremental cost-effectiveness

	Cases prevented	Incremental CER per case	Incremental CER per case (including private costs)
Stunting	39	\$1,400	\$1,700
Underweight	47	\$1,200	\$1,400
Wasting	28	\$2,000	\$2,400

Source: Authors' calculations.

In the final column of Table 11.6, we present the full costs, incorporating our estimates of the opportunity cost of women's time. While these opportunity costs did not form a large part of the overall program costs (only about 7%), when examining the relative costs of the two programs they take on a more important role since they are variable costs directly linked to the number of beneficiary-months. This is seen in the relatively large differences between the final two columns. These are not costs borne by the program, but do represent investments made by individuals to participate in, and therefore benefit from, the program.

³⁷ $(6,422 - 1,589) \times 11.64 = \$56,256$.

Although distinct from the cost-effectiveness assessment in which what matters is the effect of the program and not how many individuals were served, it also turns out that the number of individual children benefiting from the two programs differs substantially as well. In the final survey (2005), we find 73% of the beneficiaries are in preventive areas. Allocating variable costs across the two approaches yield a per beneficiary cost of \$136 for preventive areas and \$87 for recuperative areas (so that preventive is approximately 50% higher), where the difference is due to differences in the average months (11.7 versus 7.5 months) reported in the final survey (2005). When we include the (fixed) direct program costs, which are lower for preventive areas, the per beneficiary cost remains higher in preventive areas, \$312 versus \$244, although not by as much in percentage terms (approximately 25% higher).

11.4.2 Interpreting the incremental cost-effectiveness estimates

There are a number of important issues to bear in mind when interpreting the incremental cost-effectiveness figures presented in Table 11.6.

(1) Program benefits that go beyond nutrition: Perhaps the most important is that unlike many studies of health-care programs (e.g., Waters et al. 2006), the range of possible benefits from the preventive program approach relative to the recuperative approach is substantial, as documented throughout this report. This is an inherent difficulty of evaluating a complicated program that, while it may have a primary objective of reduced malnutrition, also has a number of secondary objectives. Those same \$1,300 contributing to each prevented case of stunting relative to the recuperative approach not only contributed to the other improved indicators shown in Table 11.6, but also contributed to improved outcomes shown throughout this report, such as food security of the household (due in part to the indirect or family food ration), improvements in maternal knowledge, and exclusive breastfeeding, many of which were for family members outside the target child group. Because it is a program with multiple integrated components, however, it would be inappropriate to exclude costs for specific components, for example, the family rations, whose rationale is to improve the probability that the direct ration goes to the targeted children.

(2) Inability to assign confidence intervals: A second difficulty in interpretation is that unlike in other chapters of this report we are unable to assign confidence intervals (e.g., standard deviations) around our estimates without making assumptions about the statistical distribution of the costs, something we do not know and cannot estimate in a standard framework (in part, what would be needed is repeated cost measures from different samples, for example).

(3) Potential over-valuation of costs: To this point, our analysis has been all-inclusive with respect to costs, incorporating and valuing costs of all types. Our approach is consistent with a broad view examining the program's cost to society and recognizing that all resources have an opportunity cost, that is, if they were not used for the program they would have been used in other, potentially beneficial ways.³⁸ At the same time, however, it is not always clear how to evaluate the costs of all of the inputs into the program, in particular the food commodities and beneficiary opportunity costs. It is possible that either or both of these have been

³⁸ While from the point of view of the program the food is "free," it still has an obvious opportunity cost once the program has received it - i.e., the next best use they could have made of the food. Because we are unable to directly assess that value, we instead use the monetary value at point of purchase, plus cost, insurance, and freight (CIF) as a proxy for it.

overestimated in the above calculations. Nevertheless, even if this were the case and the assessed values were overestimated by some 50%, this would at most reduce the incremental cost-effectiveness figures by one-half, and they would remain on the order of \$500 on a per case basis.

(4) Inability to assign CER to each program approach: The design of the evaluation does not permit an assessment of the CER of each of the program approaches on its own, so it is not possible to determine the average CER for the preventive approach. In the extreme case where we treat the recuperative approach as being completely ineffective, however, then the comparison for this evaluation turns out to have been one that pits the preventive approach against a “control” group. As such, the relative effectiveness measures presented in the first column of Table 11.6 could then be treated as the overall effectiveness. To assess the overall CER, then, one would simply divide the total costs for preventive areas by these effectiveness measures. This would increase the cost per case prevented for each of the outcomes by a factor greater than two.

11.5 Conclusions

The analysis presented here outlines the methodology for calculating costs and the incremental cost-effectiveness ratio for the WV-Haiti preventive versus recuperative programs.

The study areas represent about one-fifth of the overall WV-Haiti DAP intervention area, and therefore account for a similar fraction of the overall costs. We estimate total costs for the study areas, excluding external evaluation costs, to be approximately \$2.4 million, from September 2001 to September 2005. The program grew substantially over the evaluation period, both in terms of beneficiaries and in terms of costs. Clear economies of scale were evident in that the growth in costs was much slower than the growth in beneficiaries. The results also indicate that it is important to consider outside program costs. Ignoring the value of the food commodities distributed would underestimate the program costs by more than one-third in the early years and by almost one-half in FY 2005. The value of health-care supplies and beneficiary opportunity costs, however, appear to be relatively small components of the overall costs. The outside program costs, 98% of which are food commodity costs, represent the dominant variable costs of the program and vary linearly with beneficiary-months so that while economies of scale exist with respect to overall program costs, there are none with respect to these variable costs.

The principal question tested in the project is whether the preventive and recuperative approaches differ in their effectiveness at reducing malnutrition. A concern for assessing the relevance of any such differences is that the two approaches also may differ in terms of their costs. We have shown that even though the total direct costs and the variable cost per beneficiary-month are the same for both programs, the preventive approach as undertaken in this study is more costly than the recuperative approach, both on a per beneficiary-month basis and on a per beneficiary basis. This is due to a combination of design differences, the existing malnutrition rate, and different participation rates across the approaches. Importantly, the difference in program beneficiary-months across the program approaches is due to (1) the difference in the number of 6–24-month-old children participating in the program compared with the number of malnourished children under 5 participating in the program (which, in turn, is a

function of the malnutrition rate); (2) the deliberate design feature of the program that children under the preventive approach remain in the program longer, potentially twice as long; and (3) higher program participation rates among eligible children in the preventive approach compared with eligible children in the recuperative approach. If once they participated, children remained in the program for the full potential period, only in a setting where the malnutrition rate was 33% and participation rates were identical would there be no cost difference due to *who* is targeted (assuming a uniform distribution of children 0–5 years old). The malnutrition rate would have to be 66% for the design differences to entirely cancel out. Since average length of participation is not twice as long in preventive versus control, but participation rates are higher for the preventive group, malnutrition would still have to be higher than 33% for there to be no cost difference in the program as implemented. In the study program areas, however, malnutrition rates were substantially lower than this and participation rates were higher under the preventive approach, leading to substantial cost differences across the programs.

The evaluation compares the recuperative and preventive approaches, without the use of a control group. This design, without a control group, has important implications for how we carry out the cost-effectiveness analyses. What is possible under this design is a *relative* or *incremental* cost-effectiveness analysis. The impact chapter explored the effect of the preventive approach relative to the recuperative approach, in other words, what would happen when we switch from recuperative to preventive. In assessing cost-effectiveness, we took the same approach. We calculated increases in costs as one “switches” from the recuperative approach to the preventive approach, and then contrasted those costs with the gains in effectiveness seen for that same change.

We presented our estimates of incremental CERs for three indicators: prevented cases of stunting, underweight, and wasting. In this context, the evidence suggests that the gains in improved nutrition for the sample population cost on the order of \$1,000 or more per case prevented. That said, we caution the reader in interpreting these estimates for several reasons. The principal concern is that such CER estimates assess effectiveness for only a single outcome at a time, which is inappropriate for a program with multiple objectives - and multiple improved outcomes are demonstrated elsewhere in this report. The second reason we caution overemphasis on these estimates is that the value of food commodities is central to our calculations and there are reasons to question the conservative (in the sense of valuing it highly) approach we took in assessing those costs.

What is clear, however, is that the additional cost of the preventive over the recuperative approach, and therefore the incremental cost-effectiveness, depends crucially on program design, program participation, and the existing rate of malnutrition.

12. KEY FINDINGS, AND PROGRAM AND RESEARCH IMPLICATIONS

This chapter presents the main highlights of the evaluation and discusses the program and research implications of the findings. First, we summarize and discuss the results of the main question addressed by the study, i.e., is the preventive approach more effective than the recuperative approach in reducing childhood undernutrition? Second, we discuss the differences in cost and cost-effectiveness between the two program approaches. We then discuss the potential pathways of impact, placing the results within the program theory framework described in Chapter 2. Finally, we discuss the policy, program, and research implications of the findings of the study.

12.1 Relative Impact and Cost-Effectiveness

12.1.1 Is the preventive approach more effective than the recuperative approach at reducing childhood undernutrition?

Main impact – probability results: The key finding of the study is that in communities randomly allocated to receive a *preventive* approach of Title II-MCHN program, the prevalence of stunting, underweight, and wasting was 4, 6, and 4 percentage points lower after 3 years of operation than in communities exposed to the *recuperative* program approach. Mean anthropometric indicators (HAZ, WAZ, and WHZ) were also statistically significantly higher in preventive compared to recuperative communities, with differences of +0.14 for HAZ, +0.24 for both WAZ and WHZ. The magnitude of differences in favor of the preventive group for mean anthropometric indicators is comparable to other effectiveness trials aimed at reducing undernutrition through improved complementary feeding (Caulfield, Huffman, and Piwoz 1999); and to the average impact of USAID Title II MCHN programs documented by Swindale and collaborators (2004) (i.e., approximately 2 percentage points reduction in underweight prevalence per year). Although the studies included in these reviews, which used before/after or post-intervention designs with a control group are not directly comparable to our study design (which compared two food-assisted MCHN program approaches), they are indicative of a range of effect that may be expected from this type of intervention. If we assume that our recuperative approach had some impact on reducing undernutrition (as suggested by the review of USAID food-assisted MCHN programs), then the larger impact of the preventive approach must be viewed as additional to that of the recuperative approach.

To our knowledge, our study is the first to compare two different approaches of targeting Title II-MCHN programs using a randomized, community-level, longitudinal design. This probability design allows us to conclude (with only 5% probability that the results are due to chance) that the differences in undernutrition observed at the end of the study between the preventive and recuperative communities are due to differences between the two program approaches.

Plausibility of impact: The study also gathered additional data to confirm the plausibility of the results. First, we showed that children who were exposed at the optimal time (i.e., from 6 months of age) and for the entire period between 6 and 23 months of age (i.e., children 24-35 months at final survey) benefited more from the intervention than children exposed at a later age

or for a shorter duration. Second, our analysis of younger siblings (i.e., infants 0-11 months of age) of index children shows that differences between the preventive and recuperative groups are stronger and more significant for older children - a result that is plausible, given that the two programs offered the same preventive services to mothers during pregnancy and the first 6 months of lactation.

Finally, when compared to baseline, children's nutritional status appears to have deteriorated among the recuperative group, especially with regards to the prevalence of underweight and wasting. The prevalence of underweight went up by 2.8 percentage points in the recuperative communities, while it went down by 2.8 percentage points in the preventive communities. Wasting went up by 4 percentage points in the recuperative areas while it decreased slightly in the preventive areas. The prevalence of stunting, on the other hand, decreased by 3.5 percentage points since baseline in the preventive communities, while it went up very slightly in the recuperative communities. These results suggest that the preventive approach mitigated the deleterious effects on childhood malnutrition of the economic and political crisis that occurred in Haiti during the study period.

Evidence of deterioration in nutritional status in the area of the study is provided by new results from the 2005 Demographic and Health Survey (EMMUS IV 2007). The comparison of changes in malnutrition rates between 2000 and 2005 in the Central Plateau (the region where our study took place) show that stunting increased by 5 percentage points, while underweight prevalence almost doubled (from 17% in 2000 to 32% in 2005) and wasting more than tripled (from 2.2% in 2000 to 7.6% in 2005). These prevalences are not directly comparable with the prevalences reported in our study because the DHS used the NCHS/CDC/WHO reference standards (WHO 1983), whereas our study used the recently released WHO standards (WHO 2006). Nonetheless, the findings documented by the DHS point to a severe deterioration in children's nutritional status in our study area, especially in the two weight-based indicators. Thus overall it appears that both the preventive and recuperative programs may have helped mitigate the impact of the economic and political crisis on childhood undernutrition in Haiti, but that the preventive model was more effective in doing so.

12.1.2 What is the relative cost-effectiveness of the preventive and recuperative approaches?

We estimated program costs by calculating direct program costs and outside program costs. Direct costs include the costs of central expenses, overhead, salaries, and other fixed costs, while outside costs include the costs of the food and health-care supplies.

We find that the *total* direct costs for the two program approaches are the same (\$717,400), but the total variable/outside costs of the preventive approach, as designed and implemented in the evaluation area, are much higher than the costs of the recuperative approach (\$563,500 versus \$396,000), mainly because of the higher number of beneficiaries in the preventive approach (48,476 versus 33,949, including children and women).

When examining costs *per beneficiary month*, rather than total costs, we find that the direct program costs per beneficiary-month are higher in the recuperative than in the preventive approach (\$21 USD versus \$15 USD). The outside program costs (which include the costs of the

food commodities and health-care supplies) are the same in both groups – \$12 per beneficiary month.

There are three main reasons for the difference:

(1) *There is a larger number of eligible children in preventive compared to recuperative areas:* this is due to the fact that the prevalence of underweight children among the under-5s - the criterion used to screen children into the program in the recuperative areas - is approximately 25%, or 1 in 4 children. In preventive areas, all children 6-24 months (or approximately 33% [1 in 3] children less than 59 months) are eligible. This results in larger numbers of children eligible for the program in preventive than in recuperative program areas.

(2) By design, *the duration of eligibility is longer in the preventive compared to the recuperative approach:* it was decided at the outset of the study that children in the recuperative approach would stay in the program for 9 months as per the original program design; children in the preventive approach, however, would be eligible to remain in the program for the whole period between 6 and 24 months, i.e., for up to 18 months - double the duration of the recuperative approach.

(3) *The participation rates are higher in the preventive compared to the recuperative areas:* we showed in Chapter 6 that the preventive approach seemed to elicit higher participation rates than the recuperative approach, thereby further increasing the gap between the number of beneficiary-months in the two approaches.

When looking at the relative cost-effectiveness of the preventive compared to the recuperative approach, the results suggest that it cost well over \$1,200 for each additional case of stunting, underweight, or wasting prevented by the preventive approach, relative to the recuperative approach. (Unfortunately, without a control group, we cannot derive individual program cost-effectiveness estimates for comparative purposes.)

The findings for the relative cost-effectiveness should be interpreted with caution for several reasons. First, such cost-effectiveness ratio estimates assess effectiveness for only a single outcome - in this case, undernutrition. This leads to an underestimate of overall impacts of a program such as the MCHN program evaluated here, which has multiple objectives, and which has been shown to have impact on a wide range of outcomes (as documented in Chapters 8, 9, and 10). Second, the value of food is central to our calculations and there are reasons to question the high cost at which it was assessed in these calculations. A lower valuation of the food costs would yield *considerably* lower incremental cost-effectiveness ratio estimates.

As noted above, the additional expense of the preventive over the recuperative approach, and therefore the incremental cost-effectiveness, depends crucially on program design (i.e., duration of participation), program uptake, and the existing rates of malnutrition. Some modifications, especially in duration of participation, can be made to reduce the cost of the preventive approach, as long as there is evidence to indicate that shorter durations of participation might be as effective in reducing undernutrition. Also, future computation of cost-benefit ratios can help develop a fuller understanding of the longer-term benefits of investment in a preventive approach.

In summary, the nutritional impact of the preventive approach relative to the recuperative approach is substantial. The direct program costs of the preventive approach costs are lower on a per-month basis because of the economies of scale of the preventive approach. The outside program costs (including the food costs) of the two approaches are exactly the same on a per beneficiary-month basis. The total costs of the preventive approach are higher in this setting and are attributable to design, program utilization/coverage, and the existing rates of malnutrition in this setting. It is not possible, therefore, to extrapolate the relative cost-effectiveness information to other settings from this estimation without taking into account these factors.

12.2 Limitations of the Study

A limitation of the study is the lack of a control group that received no intervention. Inclusion of a control group was not feasible for cost and logistical reasons. A control group, however, would have permitted a full assessment of the impact of each program approach in and of itself, rather than an estimation of relative impact of the preventive versus recuperative approaches. The lack of a control group also has implications for the cost-effectiveness estimations, which can only be done on a relative or incremental basis, given this study design. For the impact on nutritional status, the lack of a control group is not as crucial as it is for the cost-effectiveness estimation, because information on trends in undernutrition in Haiti during the study period can be obtained from other data sources, as discussed above.

A second limitation is that this study used a community-longitudinal design, rather than a child-level longitudinal study design. The use of a community-longitudinal design based on cross-sectional surveys was an appropriate parsimonious design to examine the impact of the two program approaches on community-wide undernutrition prevalence. However, this design does not permit an analysis of the impact of the preventive and recuperative approaches on the timing of growth faltering or recovery from malnutrition among individual children, nor does it permit us to examine the impact on individual children of differences in the timing of the interventions. An additional child-level longitudinal substudy over the period necessary to obtain adequate data to examine these issues would have been prohibitively costly.

The study also had low statistical power for subgroup analyses by child age and other characteristics. Although differences in favor of the preventive approach are seen across the age range from 12 to 41 months, some of the sub-age group differences are not statistically different, due to lack of power for subgroup analyses, although they are of meaningful magnitude.

An additional limitation that prevents certain analyses and interpretation is the confounding of child age with participation. Our sample of children in the age range of 12-41 months includes a substantial proportion of children who are still currently enrolled in the program and who have not fully benefited from the program inputs (i.e., all children 12-24 months of age in the preventive approach). Similarly, many of the children in the recuperative program are also still enrolled in the program at the time of the survey. An estimation of full nutritional benefit is only possible among older children, for whom we have limited statistical power due to limited sample size. As shown in subsequent sections, however, this issue is not a problem for ascertaining the impact of program participation on other outcomes such as household food security, maternal health, and nutrition knowledge. We recommend that a follow-up survey be conducted in the study communities to include only children who have

graduated out of both the preventive and the recuperative program as this will allow an examination of the full benefit of the two programs.

Finally, it is critical to acknowledge the role of the economic and political crisis in Haiti and its potential influence on the evaluation results. Although the crisis had little impact on the actual program operations and services delivered, it had large impacts on fuel and food prices in Haiti over the time period of this study. These price impacts are expected to have led to household-level crises that could have dampened the impact of the programs on such outcomes as food security, feeding practices, and nutritional outcomes. The deterioration in nutritional status over time (in the recuperative group), the lack of long-term improvements in food security in spite of large food assistance inputs, and the lower consumption of expensive foods like animal source foods at the final compared to the baseline survey, are all indicative of the deleterious impact of the economic crisis on poor rural households such as the ones included in our sample.

12.3 Pathways of Impact


As discussed in Chapter 2, the evaluation in Haiti paid particular attention to measuring the pathways of influence as defined by the “program theory” articulated previously. In this chapter, we use the program theory to frame and discuss the results of the evaluation. Figure 12.1 revisits the “flow” of programmatic inputs from WV-Haiti to the child. Chapters 5 through 11 presented and discussed the results that come together in this framework to inform our understanding of how the preventive and recuperative programs implemented by WV-Haiti had an impact on child outcomes, maternal behavior, household food security, and other outcomes. Here, we attempt to bring these results together with some of the planned design differences between the program approaches to discuss their implications for interpreting the differential nutritional impact of the two program approaches, as well as the implication of the results for programs and for further research.

Thus, the purpose of this section is to describe (1) differences in program design, (2) differences in program delivery and program access and uptake; and (3) differences in intermediary program utilization-related outcomes (household care context, caregiver resources, and care practices) by program approach and by program participation. In describing these three aspects, we aim to provide insights into mechanisms by which different aspects of the program pathway were affected by the program and may be responsible for the differential nutritional impact of the two program approaches. A summary of the key results for these outcomes are presented in Annex 12.1.

12.3.1 Factors pertaining to program design

Some key facets of the program design were likely responsible for the differences seen between the two program approaches. These include (1) the timing, duration and continuity of supplementation; and (2) age-appropriateness and continuity of BCC. These are discussed briefly below.

Figure 12.1 Program impact pathways



(1) *Timing, duration, and continuity of supplementation*: Research on the efficacy of supplementation indicates that impact is enhanced with earlier supplementation. It is apparent from the participation patterns and timing of exposure to the program in Chapter 6 that in fact, many children in the preventive approach were indeed exposed to the program inputs early in

infancy, and even prenatally. Although a large proportion of children in the recuperative approach were also exposed prenatally, they were not exposed in early infancy and at the time of greatest potential to benefit from the supplementation. Furthermore, the total duration of exposure to the supplementation was higher in the preventive group than in the recuperative.

(2) *Age-appropriateness and continuity of BCC*: The timing, continuity, and age appropriateness of the BCC strategy could also have contributed to differences in program outcomes. Specifically, by virtue of being (potentially) enrolled in the program from pregnancy until the child was 2 years of age, mothers in the preventive group were exposed to relevant knowledge at the most appropriate time in their child's developmental trajectory, as well as for a longer duration. While women in the recuperative group were also exposed to the BCC strategy from pregnancy until the end of 6 months of lactation, there was no continuity of the BCC after 6 months of age.

12.3.2 Factors pertaining to program delivery

12.3.2.1 Program management

The study analyzed the field-based program management in some detail, focusing on the work context of the field health staff, i.e., the health promoters and assistant health promoters. The operations research conducted in 2004 (Menon et al. 2005) demonstrated that the WV health promoters and assistant health promoters were highly motivated in their jobs, and perceived that they received adequate supervision. There were no differences in staff perceptions and attitudes between the two program approaches. Overall, the positive perceptions of staff reflect the close attention paid by the higher management at WV-Haiti to staff work context and organizational issues influencing staff motivation. In addition, the strong support from the WV-Haiti management team to this evaluation, as well as their flexibility and receptivity in relation to the program design and improvement inputs provided by the IFPRI-Cornell team via the formative and operations research, is indicative of an organization that pays attention to quality of program implementation and management.

12.3.2.2 Program implementation

Chapters 4 and 5 discussed the use of formative research to design the interventions, as well as the use of operations research activities to evaluate and improve implementation of the preventive and recuperative programs. Although the programs were operating under extremely difficult political circumstances, evidence presented in Chapter 5 suggests that both program approaches were implemented at high quality. More important, there were no major differences between the program approaches in the quality of their implementation. For the impact evaluation, this finding is important, and leads to greater evidence that differences in nutritional impact of the two approaches can be attributed to the concept and design of the preventive versus the recuperative approaches rather than implementation differences.

The engagement between the IFPRI-Cornell team and the WV-Haiti staff was strengthened by the program design process described in Chapter 4 and also by the operations research process described in Chapter 5. Both of these factors led to a productive collaboration between the evaluation group and the implementation team, without one supplanting the roles of

the other. For instance, although the IFPRI-Cornell group was actively engaged in designing and assessing the operations of the two approaches, there was no involvement by the evaluation team in the day-to-day aspects of program implementation that were solely WV-Haiti's responsibility.

12.3.2.3 Program operations and inputs

At this level, we consider factors pertaining to the delivery of the Maternal and Child Health and Nutrition (MCHN) package of services to the program beneficiaries. These include the food assistance, the health and nutrition behavior change and communication strategy, and the other preventive health and nutrition MCHN program services. As highlighted in Chapter 5, these program inputs were delivered mostly as intended. More important, the operations research showed that WV staff at all levels were highly motivated in their jobs, in addition to being well-trained in some of the core skills necessary for field health staff.

The operations research conducted in 2003 (Loechl et al. 2004) showed that food assistance was generally targeted appropriately and that for the most part, the program beneficiaries received the intended amounts. Although there were some concerns about waiting times and crowding of food distribution point, the factors leading to these problems were infrastructural and largely outside the control of the WV-Haiti team. At the same time, some other concerns relating to hygiene and food handling, as well as communications between the commodities team and the health team, were amenable to change, and were addressed by WV-Haiti management.

In 2003, just two months after the BCC strategy was introduced, the operations research showed that the delivery of the BCC intervention was particularly strong in some dimensions while still lacking in others, with no differences between program approaches. Specifically, the facilitation skills of the health staff were found to be excellent, as was their coverage of the technical content of the MCs. However, the organization of MCs according to the intended age subgroups in the preventive program was not as envisioned or planned. Following discussion of these results with WV-Haiti staff, additional training and motivation of staff was undertaken and supervision was intensified. In 2004, the second round of operations research showed that the MCs organization had improved substantially, while the high quality of education and facilitation skills of the health staff was maintained.

The preventive services delivered at the RP by WV were assessed in 2003 and in 2004. In 2003, a variety of organizational issues related to the RPs were identified, as were some technical issues pertaining to the large group health and nutrition education and the growth monitoring and promotion (GMP) activities at the RP. These issues were discussed with the WV health team at the consultative workshop in 2004, following which remedial measures were implemented to reduce waiting times and crowding at the RPs, as well as to improve the large group education sessions and the GMP. An assessment of these issues in 2004 revealed substantial improvements in both these areas. Some other aspects of the services provided at the RPs, e.g., availability of vaccines and vitamin A, were less amenable to change because they were dependent on supply and logistical issues outside of WV's control. There were no differences between program approaches in any of these aspects.

Since all the program areas covered by this evaluation received services from WV-Haiti, any variability in program participation between the two program approaches is related to differential uptake/utilization of the program services by the communities served by these programs.

In sum, the engagement with WV through the formative research and operations research processes demonstrated a strong programmatic commitment to delivering high quality services with wide coverage in spite of the economic, political, and social constraints that existed in Haiti at the time of the study. These aspects were exactly the same between program approaches.

12.3.3 Factors pertaining to program access and utilization

Even well-designed, well-delivered programs with wide coverage can fail to have the expected impact if the intended users do not access the program, or if they do not use program services as intended. This is particularly important for programs that include a BCC component, where the main product delivered by the program is information and knowledge. The translation of knowledge into appropriate actions is then dependent on user acceptance of the information, and the ability of the user to implement the actions.

In the case of the preventive and recuperative programs compared in this evaluation, both programs provided information (via the MCs) as well as some direct resources (i.e., the food assistance) that could enable the translation of that information into appropriate child feeding and care practices. In addition, the peer group approach used by the MCs provided social support to mothers and engagement with others in their neighborhoods that shared the same life circumstances. In providing these programmatic inputs, therefore, the WV program had the potential to substantially impact the household care context for those households that were exposed to the program. The expectation that follows is that this would in turn lead to changes in care practices such as infant feeding practices, and that this ultimately would result in improved child nutritional outcomes. In this section we briefly summarize the impact of the WV program on the care context, focusing on differences between the two program approaches and differences by participation. We separate the care context into household and maternal contexts/resources and also discuss the differential impact of the two programs on care practices. Before doing this, however, we first discuss the differential uptake of the program by households in the preventive and recuperative programs.

12.3.3.1 Program access and uptake

Program enrollment

As shown in Chapter 6, patterns of enrollment in the targeted food assistance and BCC component of the overall MCHN program mirrored the specific targeting mechanisms used in the two program approaches. For instance, enrollment and duration of participation were the same between the two program approaches during pregnancy and the first six months of lactation, while they differed thereafter, as expected by design. In preventive program communities, participation is high between 6 and 24 months of age, also as designed, while in the recuperative communities, participation is highest among 12-23-month-old children who are more likely to be underweight and thus eligible for the program.

Program uptake

At the time of the final survey, 75% of children in preventive areas who were in the targeted age range (6-24 months) were enrolled in the program; by contrast, in recuperative areas, only 29% of currently underweight children were enrolled in the program. When looking at the percentage of current participants in the final survey who meet eligibility criteria (i.e., accuracy of targeting), the results also show that the preventive approach is doing better, with 93% of participants being in the targeted age range, compared to 57% of currently enrolled children in the recuperative group being underweight. Thus, even though targeting for the recuperative group appears to be reasonably good, uptake of underweight children is still low.

This apparently lower uptake in the recuperative group could be due to several factors such as the following: currently underweight children may have just become underweight and not have yet been identified by the program; underweight children who participated in the program previously might not have recovered; poor measurements at the RP may have resulted in misclassification³⁹; or children identified by the program as eligible may not have participated. It is also possible that mothers in recuperative areas were less likely to take their child to the RP for screening (and other services) because they did not know their child's nutritional status - while mothers in preventive areas knew whether or not their child was eligible based on age.

Clearly, the “underweight” measure at the time of the survey is a less accurate proxy than age to reflect program eligibility. For instance, currently underweight children may have just become underweight and not have yet been identified by the program; poor measurements at the RP may have resulted in misclassification; or underweight children may have participated in the program previously and not have recovered. In spite of these limitations, the results do suggest a markedly lower uptake of eligible beneficiaries in the recuperative compared to the preventive program.

Thus, the preventive approach seems to elicit greater uptake by targeted households - and possibly also less misclassification by program staff - than the recuperative approach. This is not entirely surprising, given the much greater clarity and transparency of the “age” eligibility criterion, compared to the criterion requiring child anthropometric measurements. It is likely that mothers who are not certain that their child will be eligible for the targeted food assistance - because they do not know the exact child's nutritional status - are less motivated to take the child to the RP than a mother who knows her child is eligible because he is in the right age bracket.

Use of RPs—The entry point to the program

The enrollment, eligibility, and uptake patterns described above are a reflection of programmatic practices, but they are also driven by differences in how communities use the primary entry point for the programs, i.e., the Rally Posts. All the communities in this evaluation were covered by the RPs and differences in participation between approaches are thus attributable to differential uptake patterns of the entry point into the program, i.e., the RP.

³⁹ Note that the operations research in 2003 (Loechl et al. 2004) did reveal some misclassification of children, but this was not large. Therefore, we do not believe that misclassification is a major factor explaining the lower coverage of potentially eligible children in the recuperative areas.

Although overall RP participation was not different between program groups, we found that RP participation was age-dependent and that age-based patterns of RP use differed between the preventive and recuperative approaches. Although there were no differences in RP participation for infants in their first year, there were marked differences for older children, with fewer children 24 months and older attending the RPs in preventive compared to recuperative communities. These age differences in participation, which were also observed in our 2004 operations research (Menon et al. 2005), support the hypothesis proposed above of clearer community and caregiver understanding of the different eligibility mechanisms used in the two program approaches. In preventive areas, mothers know that after 24 months of age, unless their child is severely malnourished ($WAZ < -3$ Z-scores), s/he is no longer eligible for program food commodities, and thus, they stop attending RPs. In recuperative areas, mothers are probably less clear about when exactly their child is sufficiently undernourished to enter the program, but it seems that at least they understand the importance of taking the child to the RP in the first year for immunization and other preventive services.

Characteristics of participant and nonparticipant households and caregivers

Given that household characteristics can significantly impact program participation patterns, we carefully examined the characteristics of households in the preventive and recuperative groups, as well as the characteristics of participant and nonparticipant households within each of the program groups. We found no significant differences in household or caregiver characteristics between program groups at the time of the final survey, although caregivers in the preventive communities were slightly more likely to be working away from home. There were also few differences in household and caregiver characteristics between participant and nonparticipant households in each of the program groups, with the exception of small differences in household size and caregiver resources such as communication, asset ownership, and marital status. Because these could reflect a cluster of support-related resources that made it possible for households to participate in the program, more extensive analysis of these aspects will be pursued in the future to better characterize understand the determinants of program participation. Overall, however, it appears that program participants and nonparticipants were not different.

12.3.3.2 Program participation: Impact on care resources and care practices

The program inputs provided by the WV MCHN program were intended to improve the household and caregiver context within which childcare occurs, and consequently the care practices themselves. In this section we document how the program impacted on these aspects, by comparing differences between the two program approaches at final survey as well as differences between participants and nonparticipants within the program approach. Understanding differences in the program pathway between the preventive and recuperative approaches is useful to help explain which intermediary mechanisms led to a greater impact of the preventive approach on child nutritional outcomes. Understanding differences between program participants and nonparticipants helps clarify the overall impact of WV-Haiti's program on targeted communities.

Impact on household context and household resources

The two indicators of household context and resources used in our survey were household food insecurity and asset ownership. At final survey, households in preventive communities had significantly lower food insecurity than households in recuperative areas, but differences were small. More meaningful differences were observed between current participation and nonparticipants in both program groups, suggesting a positive short-term impact of food assistance on household food security. Compared to baseline, overall food insecurity did not improve over the three-year duration of the study, and continued to be severe in the survey area. There was thus no evidence of a long-term effect of the program on food security; the same was true for asset ownership, which had not improved since baseline, and was not different between program communities or between participants and nonparticipants. Thus, overall, the program seems to have had a positive impact on food security in the short term, suggesting that the slightly greater impact on households in preventive compared to recuperative communities may be due to their longer eligibility to receive food assistance. There was, however, no evidence that the program had a long-term impact on food insecurity or asset ownership, which is most likely due to the severe economic constraints faced by poor households in Haiti at the time of the study. It is possible that more sustained impacts of such a program could be obtained in less constrained times.

Impact on maternal context and resources

Maternal resources that are thought to be important determinants of childcare practices and child nutrition include knowledge, education, physical and mental well-being, asset ownership, social support, and decisionmaking power (Engle, Menon, and Haddad 1999). In this section we briefly summarize our findings related to the caregiver resources that could have been affected by the program, namely knowledge and physical and mental well-being. We did not find any differences between program groups on other maternal resources such as education, social support, autonomy in decisionmaking either at baseline or at the final survey.

Impact on maternal knowledge: Maternal knowledge of several topics related to infant and young child feeding and general health and nutrition topics was higher among the preventive compared to the recuperative group at final survey, but differences were generally small. More meaningful differences were found between caregivers who had ever participated in the program, compared to those who had never participated. Maternal knowledge also significantly improved from baseline to final survey in both program groups, highlighting an overall positive impact on maternal knowledge of WV's MCHN program in the study area, despite large differences in uptake between program groups.

Physical and mental well-being: Our measures of physical well-being were based on body mass index and a visual analogue scale of self-rated health. The mental well-being measures assessed mental stress, time-related stress, and the frequency of stress-related symptoms. There were no differences between program communities at baseline on any of these measures. At baseline, we also found that food insecurity was strongly associated with all measures of women's well-being, particularly mental well-being (Menon et al. 2004). This finding was indicative of the wide-ranging consequences of food insecurity for Haitian households.

At the time of the final survey, respondents in the preventive communities were better off than those in the recuperative communities on four of the women's well-being measures. There were no differences between respondents who had ever been exposed to the program and those who had not. However, current participants in preventive communities had better self-rated health, lower mental stress, and lower time stress than nonparticipants. This could be due as suggested by our baseline analyses, to short-term less severe food insecurity. This potential mediating pathway between program participation and women's well-being through increased food security will be examined in future analyses of the data.

In summary, the impact of participation in the WV program is seen both on maternal health and nutrition knowledge, as well as on women's mental well-being. The mechanisms through which these intermediate programmatic outcomes were affected, however, appear to differ, with knowledge being associated with any exposure to the program, but mental well-being being associated only with current participation. In both cases, however, it is likely that differences between program approaches are driven by differences in participation rates and duration of participation between the two program approaches.

Impact on care practices

The following three subsections discuss the impact of the preventive and recuperative approaches on care practices. In reviewing these results, it is useful to be aware that these data were all gathered through survey questionnaires that rely on recall-based methods. Program participants could be more likely to report good practices because of their increased awareness and knowledge, even if the practices were not adhered to completely. This is a well-known issue with recall methods. At the same time, it should be noted that multiple questions focused on the same type of practice have yielded similar responses in this study, and improvements are not seen for all practices, thus providing reassurance that the recall-based data are not systematically biased.

Awareness, trial, and adoption of recommended practices

The final survey included an assessment of the awareness, trial, and adoption of seven key practices recommended by the program. The practices ranged from the use of expressed breast milk when mothers left the house to feeding a convalescing child an extra meal a day after a bout of illness. For most key practices, respondents in preventive program areas were more likely to report awareness, trial, and adoption than were respondents in recuperative areas. In most cases, however, differences between program areas were of relatively small magnitude. However, differences between those exposed to the program and those not exposed (in each area) were much larger than differences between program areas. Reported adoption rates for different practices among recuperative group participants ranged from 15% to 48%, while it ranged from 19% to 57% among participants in the preventive group. Among nonparticipants (in both groups), reported adoption ranged from 2% to 31%.

Feeding practices and care

We assessed the impact of the program on care practices by focusing mainly on breastfeeding and complementary feeding practices, preventive and curative care-seeking

practices, and markers of hygiene practices. There were no differences in any of these practices at baseline.

Breastfeeding: There were no differences between program groups in early feeding practices or in the timing of introduction of liquids and complementary foods. This was expected since the two program approaches offered exactly the same services until the child reached 6 months of age. Large improvements in these practices were seen since baseline, however, and equally large differences were observed between participants and nonparticipants for both program approaches. Breastfeeding duration was the same in both program groups at baseline and at final survey. Haitian women in our sample breastfed until the child was between 18 and 24 months of age, and the program was able to sustain this positive practice. More important, the maintenance of this pattern of breastfeeding since baseline is evidence that the provision of donated food commodities as complementary foods did not diminish the duration of breastfeeding.

Complementary feeding practices, including consumption of WSB: We assessed many dimensions of complementary feeding such as frequency of feeding, dietary diversity, use of baby bottles, assistance during feeding, feeding during and after illness, and use of vitamin supplements such as vitamin A. The appropriateness of reported practices was assessed by comparing practices to age-specific recommendations. We also examined consumption of WSB, the main fortified commodity targeted to the beneficiary children.

There were few differences between program groups in complementary feeding practices at final survey, with the exception of diet quality and the consumption of animal source foods, which were slightly higher in the preventive group. The summary indicator of infant and young child feeding, which combined breastfeeding, frequency of feeding, and dietary diversity information, also showed a statistically significant, but small difference in favor of the preventive, compared to the recuperative group. Compared to baseline, consumption of animal source foods was lower at final survey, probably a reflection of the economic crisis and related price increases in Haiti over the study period.

Although the proportion of children currently in the program and consuming recipes made with WSB was not different between the program groups, there is a much larger *number of* program beneficiaries receiving WSB and other fortified foods in the preventive group, particularly among the younger children. It can thus be assumed that the fortified commodities contribute more to caloric and nutrient intake among children in the preventive approach than in the recuperative approach.

Several practices had markedly improved since baseline; these included the use of baby bottles (which was halved since baseline), vitamin A supplementation, and feeding during and after diarrhea.

As with maternal knowledge, differences in many of the feeding practices were larger between participants and nonparticipants than between program groups. The practices for which substantial differences were seen between participants and nonparticipants were meal frequency, use of baby bottles, vitamin A supplement consumption, and consumption of fortified donated commodities (WSB) (the latter being expected). The pattern of differences between participants

and nonparticipants and the reported reasons for non-trial and adoption of practices suggests that overall, practices that required few material resources to try and adopt were more likely to have improved. For instance, feeding the child animal source foods and nutrient-rich vegetables were less likely to have improved than avoidance of baby bottles.

Preventive and curative care-seeking and hygiene

Immunization rates were not different between groups at baseline, but they were extremely low, with only 11% of children fully immunized. At the final survey, immunization rates were not different between the program groups, and had increased substantially since baseline, although they were still excessively low at approximately 30%. The reasons for this seemed to be largely due to poor supply as opposed to low demand, as suggested by our operations research results (Menon et al. 2005).

Care-seeking for fever, cough, fast breathing, and diarrhea were not different between program groups at baseline or at the final survey and were not different between participants and nonparticipants. However, care-seeking rates were lower at the final survey when compared to the baseline survey. Since care provision in the program areas was better at the time of the final survey compared to the baseline, it is likely that lower curative care-seeking at the final survey reflected decreased severity of illness and/or better home management of illness. For instance, use of ORS and SSS were higher since baseline, suggesting improved home care for diarrhea as a result of the program.

There were no differences in markers of hygiene practices between program groups either at baseline or final surveys, but there was a slight decrease was seen in hygiene scores since baseline. This again may be a reflection of the deteriorating socioeconomic conditions in the study communities following the economic crisis. There were also no meaningful differences in markers of hygiene practices between participants and nonparticipants.

12.3.4 Conclusions on pathways of impact

Our findings suggest that the pathways of impact, which led to better child nutritional outcomes among preventive communities, operated mainly through differences in design between the two programs, as well as differences in participation and uptake patterns and the overall childcare context:

- ❖ *Design aspects:* The key design differences in the two programs that likely contributed to difference in outcomes are the timing and duration of participation in the program, as well as the continuity of the BCC strategy for the preventive group. Children in the preventive program were more likely to be exposed to program inputs at an age when they are known to be more likely to respond to supplementation. They also received benefits for a longer duration, which is also known to enhance the impact of supplementation.
- ❖ *Program uptake and participation:* An additional benefit of the preventive approach is that it clearly elicited greater participation and thus reached a significantly larger number of mothers and children. Empowering women with improved knowledge, awareness, and care practices will not only have a positive

impact on infants and young children today; it will also have an impact on these children's younger siblings and possibly on whole communities in the longer run.

- ❖ *Childcare context, care practices, and consumption of fortified commodities:* Participants in preventive communities had greater short-term food security and caregivers scored better on mental well-being scales. They also scored higher on several complementary feeding practices and preventive health-care-seeking practices. In addition, greater consumption of fortified commodities among the preventive group also likely contributed to differences in impact. Overall, the number of children who received the fortified commodities during the period when they were most likely to benefit from nutritional supplementation is larger in the preventive compared to the recuperative group. This is due to high program uptake, good program targeting, as well as use of the fortified commodities for the target child. The children also received the foods for longer in the preventive approach.

Our results also showed that the two program approaches were operating equally well and that none of the program implementation and staff-related factors differed between the two approaches. This allows us to conclude with certainty that the greater nutritional impact observed in preventive communities was truly due to a more effective program approach.

12.4 Implications for Programs, Policies, and Future Research

12.4.1 Implications for programs and policies

A preventive approach to addressing childhood undernutrition is more effective than a curative approach. The results of this three-year evaluation provide evidence that targeting of a food-assisted MCHN program to children under-two years of age using a preventive approach is more effective at reducing community-wide undernutrition than a recuperative approach. The direct implications of these results are that in order to improve effectiveness, food-assisted MCHN programs should target all children under the age of two years, as opposed to malnourished children under-five, and continue to target pregnant and lactating women. Severely malnourished children should continue to be screened and receive appropriate care.

Focusing on the under-twos is both feasible and successful in a programmatic context. There is renewed attention around the need to target the critical under-two age group, but few examples exist of feasible, successful, and effective programs. This evaluation provides a successful example of how targeting food assistance and behavior change communication to this age group can reduce the prevalence of undernutrition. Perhaps more important, it provides an example of *how* such programs can be developed, strengthened, and monitored under real programmatic conditions. Our work thus provides programmatic evidence supporting a focus of nutrition interventions on the under-twos.

In addition, our findings suggest that targeting is more efficient with the preventive approach since it relies on age rather than nutritional status. In addition, the preventive approach seems to elicit greater participation among those eligible for benefits. This could be because the programmatic targeting mechanisms are more transparent to community members and parents.

A well-designed and well-implemented behavior change strategy can improve infant feeding practices regardless of whether a preventive or recuperative approach is used for targeting. The challenge of developing and implementing locally relevant, programmatically feasible strategies for improving feeding practices among children under-two is a major hurdle to programmatic success. Our work provides an example of an approach and a specific set of tools that were used to develop and implement an effective behavior change and communication strategy integrated within a food-assisted MCHN program.

Cost per beneficiary-month is the same for the two programs but total costs are higher for the preventive approach as designed in this study. The implications of these findings are that programs should carefully review their program design, geographic priority areas, and targeting mechanisms based on their resources and target number of beneficiary-months. Programs can also attempt to balance or reduce the costs of the preventive approach by changing the age range for targeting, the duration for which children are enrolled in the program, and/or even the amount of the food assistance provided. The impact of making any of these changes to the preventive approach, however, should be rigorously tested through evaluation research to ensure that these modifications do not result in losses in effectiveness and nutritional impact of the program.

Investing in formative and operations research is important for program success. This evaluation provides strong evidence that investing in formative research can help design BCC programs that are grounded in the sociocultural context and locally relevant, and therefore, are effective. Additionally, using formative research to explore potential opportunities for integrating the BCC strategy within an existing program and identifying the critical capacities that need to be strengthened allows the development of a solid and well-grounded BCC strategy. The study also provides evidence that operations research is a critical component of program evaluation and that it provides critical insights regarding the quality of implementation and service delivery. In addition, it is a good investment for programs because it helps solve operational bottlenecks and ensure smooth sailing. Several elements of operations research procedures and methods used by the evaluation team were adopted by WV-Haiti on a routine basis to examine and correct implementation problems in the non-study areas.

An important element of the formative and operations research done as part of this evaluation was the packaging and communication of the results for use by the program implementation team at WV-Haiti. In both cases, the results of the research were summarized in matrices that brought the results together in a format that allowed for clear communication and engaged discussions of potential programmatic solutions. In addition, the results were presented and discussed with program staff at all levels in the WV-Haiti hierarchy, including field staff. This allowed for engagement with the formative and operations research process and outcomes, and likely led to greater commitment to improving the program at all levels.

How generalizable and replicable are the results of the Haiti evaluation? An important question for programs is whether or not the results of the Haiti study are generalizable and replicable. The results are *generalizable*, given the remarkably similar patterns of child growth globally, as well as the similar determinants of poor child growth worldwide, i.e., poor infant feeding and care and repeated illnesses. They are also generalizable, given what is known about the global impact of programs to address nutrition by improving infant feeding and care, and reducing illness.

For *replicability*, we believe that there are four important conditions: (1) good program design based on sound formative research; (2) effective implementation and service delivery monitored with regular operations research; (3) good incentive structures and high staff motivation monitored and fostered by effective supervision; and (4) similar or higher levels of undernutrition than in Haiti (e.g., at least 25%-30% underweight/stunting). Although a similar preventive program could be effective in a population with lower levels of undernutrition than in Haiti, the cost per case of undernutrition prevented would be higher. Most poor countries in Africa and Asia, and even some countries of Latin America, currently have higher prevalences of undernutrition than Haiti.

12.4.2 Research implications

Evaluation of the preventive approach in other settings, preferably using a control group. As discussed previously, it was not possible to include a control group in this evaluation. However, it would be useful if future evaluations of the preventive approach could include a control group, so that the magnitude of the absolute effect of the preventive approach could be assessed. In addition, evaluations in other contexts should pay adequate attention to program theory as well as the design and implementation of the preventive approach. This will help generate knowledge and consensus on best practices for developing and implementing programmatic approaches for delivering nutrition services to this critical age group.

Separating contributions of food assistance and BCC components. Given the integrated nature of the MCHN program implemented by WV-Haiti in Haiti, it was not possible to separate out the contributions of the food component from that of the BCC component. Since the food component contributes the majority of the cost of the program approach, it would be useful to conduct evaluations that can help separate the contributions of these two components. At the same time, it should be recognized that in some extremely impoverished settings, such as in Haiti, it would likely be futile to expect that BCC alone could lead to substantial impacts on childhood undernutrition. As demonstrated by our research on the effectiveness of the BCC strategy, even the provision of additional resources such as food assistance did not fully enable the translation of all program recommendations into practice.

Age range and duration for targeting the preventive program. A major reason for the relatively high cost of the preventive approach in Haiti was the large age range across which children were recruited into the program (6-18 months) and the duration that children remained in the program for (up to 18 months). In this setting, it is unclear whether a shorter age range for targeting or a shorter duration of enrollment would have yielded the same impact. However, this is something that should be investigated both in similar settings as Haiti as well as in other contexts. It should be noted that the age range and the duration of enrollment were designed based on the best available efficacy evidence at the time this study was designed. This can and should be re-evaluated as more efficacy and programmatic evidence becomes available, since reducing the age range and/or the duration of food assistance in the preventive program could lead to considerable cost savings. We also recommend a follow-up of children after they are 60 months old in both program approaches. This would allow an examination of the full impact of both approaches.

12.5 Lessons for Program Evaluations

Using program theory to develop evaluations. The approach to this evaluation was based on program theory that considered the full pathway of expected impacts. Developing the evaluation activities based on this framework was useful not only for the measurement of the different program inputs and outcomes but also to help identify bottlenecks that could influence the program's impact. Based on these experiences, we highly recommend a systematic, program-theory-based approach to conducting evaluations. At the same time, it is important that researchers recognize the limitations that are posed by the program theory that they adopt. In our case, we used the UNICEF conceptual framework for nutrition and the extensive work on care resources and care practices as guidance for developing the program theory. Although the UNICEF framework is comprehensive, the use of the food-health-care lens of the UNICEF framework leads to a different approach to data collection and interpretation of results than the application of other frameworks that are, say, more focused on poverty, livelihoods, gender, intrahousehold allocation, or other lenses.

Addressing probability, plausibility, and adequacy. The approach used for this evaluation brought together different levels of evidence for the impact of the preventive approach when compared to the recuperative approach. The community-randomized design provided probability evidence, while the careful attention to measuring the full programmatic pathways provided evidence for the plausibility of impact. The reduction in stunting seen in the preventive group between baseline and the final survey exceeded the impact proposed by WV-Haiti in their four-year DAP; this provided evidence for the adequacy of the impact. We believe it is important that program evaluations carefully assess the level of evidence they require, and this should be based on the intended purpose of the evaluation (Habicht, Victora, and Vaughan 1999).

Intervention design. We paid careful attention to the design of the preventive approach in particular because it was important to ensure that the approach was truly preventive in nature, both from a food package perspective as well as from the BCC strategy perspective. There was also no example of a truly preventive Title-II food assisted MCHN program documented in the literature that we could draw on (unlike the widely used recuperative approach, for instance). This attention to design is important to consider in all evaluations, because evaluating a package that is not designed to be true to the concept behind it is inefficient and does not allow appropriate interpretation of the results of an evaluation.

Documenting implementation quality and program utilization. Attention to the quality of implementation of the intervention package is also critical in any impact evaluation. As with the design, a poorly implemented intervention is much less likely to yield the expected impact than a well implemented intervention. Thus, attention to design and to implementation quality are both essential to draw conclusions about the effectiveness of interventions. This is often neglected in impact evaluations and leads to erroneous conclusions about effectiveness.

In addition to assessing implementation quality, it is also important to assess both program participation and uptake of program inputs by eligible beneficiaries. As discussed previously, an intervention's effectiveness can be reduced by quality of design, quality of implementation, and also by the level of uptake. We examined each of these determinants of

effectiveness carefully in this evaluation and it is clear that assessing different elements of program uptake can provide insights into patterns of access to services (participation) as well as actual household uptake of services among beneficiaries. It also provides insights into the constraints that prevent adequate and appropriate use of the intervention as intended.

Age of intervention versus age of expected impact. Finally, we suggest that it is extremely important that evaluations focused on child growth outcomes pay attention to the age at which an intervention to improve growth is delivered versus the age at which impact is expected. All outcome assessments should be conducted within the age group that is most likely to show impact. This is particularly important, given the new policy direction to target the under two age group with nutritional interventions to improve undernutrition outcomes globally. While it is appropriate to target this age group for programmatic inputs, it is not entirely appropriate to measure outcomes among the younger children in this age range because they have not yet had a chance to fully benefit from the intervention. Ideally, the age at which impact on child anthropometric outcomes should be assessed will be based on a clear understanding of growth patterns in each context and expectations for timing of the impact of an intervention. It is certainly appropriate to include some children in age groups that are less likely to show impact (as we have done) because this enhances the plausibility of impact, but it is inappropriate to expect full impact among children who are still too young to manifest the full impact of a nutritional intervention.

12.6 Conclusions

This evaluation shows that a preventive approach to a Title-II MCHN program is more effective than the traditional, recuperative approach at reducing the prevalence of childhood undernutrition among children aged 12-41 months. Both program approaches cost exactly the same to implement on a beneficiary-per-month basis but the preventive approach was more expensive because of differences in design and participation rates between the two approaches. Moreover, the relatively low levels of undernutrition in this population (compared to other equally poor countries) results in a relatively larger number of children under-two, compared to underweight children under 5 years, and thus more beneficiary-months in the preventive approach. Both programs also had a significant impact on improving maternal knowledge and feeding practices compared to baseline, showing that a well-designed and well-implemented BCC strategy integrated within a Title-II MCHN program can be highly effective. These benefits were obtained with two carefully designed and implemented program approaches operating under particularly difficult field conditions in rural Haiti.

Based on these results, we recommend that Title II food-assisted MCHN programs focus their resources on children under the age of two years, using a preventive focus that also targets pregnant and lactating women. We also recommend that formative research be used to develop locally appropriate BCC strategies to help improve infant and young child-feeding practices among the target population for a preventively-focused food-assisted MCHN program. Together, a well-designed and well-implemented preventively focused BCC strategy and a food assistance package can achieve significant reductions in undernutrition.

REFERENCES

- Adam, T. *Sources of Variability in Costing Methods: Implications for the Transferability of Cost-Effectiveness Results*. Rotterdam: University Medical Center, 2006.
- Allen, L. H. "Nutritional Influences in Linear Growth." *European Journal of Clinical Nutrition* 48 (Suppl. 1) (1994): S75-S89.
- Beaton, G. "Which Age Group Should Be Targeted for Supplementary Feeding?" In *Nutritional Issues in Food Aid*. ACC/SCN Symposium Report Nutrition Policy Discussion Paper No. 12, pp. 37-54. Geneva: United Nations Administrative Committee on Coordination/Subcommittee on Nutrition (ACC/SCN), 1993.
- Beaton, G. H., and H. Ghassemi. "Supplementary Feeding Programs for Young Children in Developing Countries." *American Journal of Clinical Nutrition* 35 (Supplement 4) (1982): 863-916.
- Bhandari, N., S. Mazumder, R. Bahl, J. Martines, R. E. Black, and M. K. Bhan. "Infant Feeding Study Group. An Educational Intervention to Promote Appropriate Complementary Feeding Practices and Physical Growth in Infants and Young Children in Rural Haryana, India." *Journal of Nutrition* 134 (9) (2004): 2342-48.
- Bhandari, N., S. Mazumder, R. Bahl, J. Martines, R. E. Black, and M. K. Bhan. "Infant Feeding Study Group. Use of Multiple Opportunities for Improving Feeding Practices in Under-Twos within Child Health Programmes." *Health Policy and Planning* 20 (5) (2005): 328-36.
- Bilinsky, P., and A. Swindale. *Months of Inadequate Household Food Provisioning (MIHFP) for Measurement of Household Food Access: Indicator Guide*. Washington, D.C.: Food and Nutrition Technical Assistance Project, Academy for Educational Development, 2005.
- Brownie, C., J.-P. Habicht, and B. Cogill. "Comparing Indicators of Health and Nutritional Status." *American Journal of Epidemiology* 124 (6) (1986): 1031-44.
- Caulfield, L. E., S. L. Huffman, and E. G. Piwoz. "Interventions to Improve the Intake of Complementary Foods by Infants 6-12 Months of Age in Developing Countries: Impact on Growth and Prevalence of Malnutrition and Potential Contribution to Child Survival." *Food and Nutrition Bulletin* 20 (2) (1999): 183-200.
- Cogill, B. *Anthropometric Indicators Measurement Guide*. Washington, D.C.: Food and Nutrition Technical Assistance Project, Academy for Educational Development, 2003.
- Cohen, J. *Statistical Power Analysis for the Behavioral Sciences*. 2nd Edition. Hillsdale, N.J.: Lawrence Erlbaum Associates, 1988.

- Dewey, K. G., and K. H. Brown. "Update on Technical Issues Concerning Complementary Feeding of Young Children in Developing Countries and Implications for Intervention Programs." *Food and Nutrition Bulletin* 24(1) (2003): 5-28. "Erratum" in *Food and Nutrition Bulletin* 24 (2) (2003): 239.
- Dewey, K. G., R. J. Cohen, and N. C. Rollins. "Informal Working Group on Feeding of Nonbreastfed Children. WHO Technical Background Paper: Feeding of Non-Breastfed Children from 6 to 24 Months of Age in Developing Countries." *Food and Nutrition Bulletin* 25 (4) (2004): 377-402.
- EMMUS-III (Enquête Mortalité Morbidité et Utilisation des Services) 2000/2001. Ministère de la Santé Publique et de la Population (MSPP), Institut Haitien de l'Enfance, ORC Macro, Calverton, Md., 2001.
- EMMUS-IV. (Enquête Mortalité Morbidité et Utilisation des Services) 2005/2006. Ministère de la Santé Publique et de la Population (MSPP), Institut Haitien de l'Enfance, ORC Macro, Calverton, Md., 2007.
- Engle, P. L., P. Menon, and L. Haddad. "Care and Nutrition: Concepts and Measurement." *World Development* 27 (8) (1999): 1309-37.
- Fiedler, J. L. *A Cost Analysis of the Honduras Community-Based Integrated Child Care Program*. Health, Nutrition, and Population Discussion Paper. Washington, D.C.: World Bank, 2003.
- Foreman, E. K. *Survey Sampling Principles*. New York: Marcel Dekker, Inc., 1991.
- Habicht, J. P., C. G. Victora, and J. P. Vaughan. "Evaluation Designs for Adequacy, Plausibility and Probability of Public Health Programme Performance and Impact." *International Journal of Epidemiology* 28 (1999): 10-18.
- International Crisis Group. "Haiti's Transition: Hanging in the Balance." Latin America/Caribbean Briefing N°7, 8 February 2005
- Jamison, D. T., J. G. Breman, A. R. Measham, G. Alleyne, M. Claeson, D. B. Evans, P. Jha, A. Mills, and P. Musgrove. *Priorities in Health*. Washington, D.C.: Oxford University Press and the World Bank, 2006.
- Katz, J. "Sample-Size Implications for Population-Based Cluster Surveys of Nutritional Status." *American Journal of Clinical Nutrition* 61 (1995): 155-160.
- Loeche, C., P. Menon, M. T. Ruel, and G. Pelto. "Process Used to Design an Integrated Health and Nutrition Program to Prevent Child Malnutrition in Rural Haiti." Submitted to the Food and Nutrition Technical Assistance (FANTA) Project, Academy for Educational Development, Washington, D.C., July, 2003a.

- Loechl, C., P. Menon, G. Pelto, and M. Ruel. "Behavior Change Communication to Improve Infant and Young Child Feeding Practices in Rural Haiti: Training and Communication Materials." Submitted to the Food and Nutrition Technical Assistance (FANTA) Project, Academy for Educational Development, Washington, D.C., January 31, 2003b.
- Loechl, C, M. T. Ruel, G. Pelto, and P. Menon. "An Operations Evaluation of WV-Haiti's Integrated Health and Nutrition Program in Central Plateau, Haiti." Submitted to the Food and Nutrition Technical Assistance (FANTA) Project, Academy for Educational Development, Washington, D.C., March 2004.
- Lutter, C. K., J. O. Mora, J.-P. Habicht, K. M. Rasmussen, D. S. Robson, and M. G. Herrera. "Age-Specific Responsiveness of Weight and Length to Nutritional Supplementation." *American Journal of Clinical Nutrition* 51 (3) (1990): 359-64.
- Mackintosh, U. A., D. R. Marsh, and D. G. Schroeder. "Sustained Positive Deviant Child Care Practices and Their Effects on Child Growth in Viet Nam." *Food and Nutrition Bulletin* 23 (4) (2002): 18-27.
- Maluccio, J., and C. Loechl. "Preventive versus Recuperative Targeting of Food Aid: Accounting for the Costs." Mid-term Cost Report. Submitted to the Food and Nutrition Technical Assistance (FANTA) Project, Academy for Educational Development, Washington, D.C., December 2004.
- Maluccio, J. A., and C. Loechl. "Preventive versus Recuperative Targeting of Food Aid: Accounting for the Costs." Final cost report submitted to the Food and Nutrition Technical Assistance Project, Academy for Educational Development, Washington, D.C., 2006.
- Martorell, R. ed. "The Effects of Improved Nutrition in Early Childhood: The INCAP Follow-Up Study." *Journal of Nutrition* 125 (4S) (1995): 1027S-1138S.
- Menon, P., and M. Ruel. "Childcare, Nutrition, and Health in the Central Plateau of Haiti: The Role of Community, Household, and Caregiver Resources." A report of the IFPRI-Cornell WV-Haiti Baseline Survey, Haiti, 2002, submitted to the Food and Nutrition Technical Assistance Project, Academy for Educational Development, Washington, D.C., 2003.
- Menon, P., M. T. Ruel, G. Pelto, and J.-P. Habicht. "Review of Health and Nutrition Education Messages and Delivery System Currently Used in Haiti, and Recommendations for Further Research." Report submitted to Food and Nutrition Technical Assistance (FANTA), Academy for Educational Development, Washington, D.C., 31 December 2001.
- Menon, P., M. T. Ruel, M. Arimond, and E. A. Frongillo. "Hungry and Stressed: Household Food Insecurity Is Associated with Caregiver Physical and Mental Well-Being in Rural Haiti." Experimental Biology meeting abstracts [accessed at <http://select.biosis.org/faseb>]. *FASEB Journal* 18; Abstract # 848. Presented at Experimental Biology, 2004, Washington, D.C., 2004.

- Menon, P., C. Loechl, G. Pelto, M. Ruel, and E. Metellus. "Development of a Behavior Change Communications Program to Prevent Malnutrition in the Central Plateau of Haiti: Results and Challenges from a Formative Research Study." Report submitted to the Food and Nutrition Technical Assistance (FANTA) Project, Academy for Educational Development (AED), Washington, D.C., September 2002b.
- Menon, P., M. T. Ruel, G. Pelto, Y. Francois, E. Metellus, and A. Ferrus. "A Qualitative Study of the Patterns of Infant Feeding and Care in the Hinche Area of Plateau Central, Haiti." Report submitted to Food and Nutrition Technical Assistance (FANTA), Academy for Educational Development, Washington, D.C., April 8, 2002a.
- Menon, P., M. Arimond, M. T. Ruel, G. Pelto, J.-P. Habicht, and C. Loechl. "Using Operations Research to Monitor Program Implementation and Inform Impact Evaluation. Results of a Second Round of Operations Research of the WV-Haiti Maternal and Child Health Program in Haiti." Submitted to the Food and Technical Assistance (FANTA) Project, Academy for Educational Development, Washington, D.C., March 2005.
- Mukuria, A. G., M. T. Kothari, and N. Abderrahim. *Infant and Young Child Feeding Updates*. Calverton, Md.: ORC Macro, 2006.
- Murray, D. M. *Design and Analysis of Group-Randomized Trials*. New York: Oxford University Press, 1998.
- Musgrove, P., and J. Fox-Rushby. "Cost-Effectiveness Analysis for Priority Setting." In *Disease Control Priorities in Developing Countries*, edited by D. T. Jamison, J. G. Breman, A. R. Measham, G. Alleyne, M. Claeson, D. B. Evans, P. Jha, A. Mills, and P. Musgrove. Washington, D.C.: Oxford University Press and the World Bank, 2006.
- Pachon, H., D. G. Schroeder, D. R. Marsh, K. A. Dearden, T. T. Ha, and T. T. Lang. "Effect of an Integrated Child Nutrition Intervention on the Complementary Food Intake of Young Children in Rural North Viet Nam." *Food and Nutrition Bulletin* 23 (4) (2002): 62-9.
- PAHO/WHO (Pan American Health Organization/World Health Organization). *Guiding Principles for Complementary Feeding of the Breastfed Child*. Pan American Health Organization/World Health Organization, Washington, D.C., 2003.
- Penny, M. E., H. M. Creed-Kanashiro, R. C. Robert, M. R. Narro, L. E. Caulfield, and R. E. Black. "Effectiveness of an Educational Intervention Delivered Through the Health Services to Improve Nutrition in Young Children: A Cluster-Randomized Controlled Trial." *Lancet* 365 (9474) (2005): 1863-72.
- Prochaska, J. O., and C. C. DiClemente. "States and Processes of Self-Change of Smoking: Toward an Integrative Model of Change." *Journal of Consulting Clinical Psychology* 51 (3) (1983): 390-95.
- Rivera, J. A., and J.-P. Habicht. "The Recovery of Guatemalan Children with Mild-to-Moderate Wasting: Factors Enhancing the Impact of Supplementary Feeding." *American Journal of Public Health* 86 (10) (1996): 1430-34.

- Rivera, J. A., and J.-P. Habicht. "Effect of Supplementary Feeding on the Prevention of Mild-to-Moderate Wasting in Conditions of Endemic Malnutrition in Guatemala." *Bulletin of the World Health Organization* 80 (12) (2002): 926-32.
- Robert, R. C., J. Gittelsohn, H. M. Creed-Kanashiro, M. E. Penny, L. E. Caulfield, M. Rocio Narro, and R. Black. "Process Evaluation Determines the Pathway of Success for a Health Center-Delivered, Nutrition Education Intervention for Infants in Trujillo, Peru." *Journal of Nutrition* 136 (2006): 634-641.
- Rossi, P. H., M. W. Lipsey, and H. E. Freeman. *Evaluation: A Systematic Approach*, 7th edition. Thousand Oaks, Calif.: Sage, 2004.
- Roy, S. K., G. J. Fuchs, Z. Mahmud, G. Ara, S. Islam, S. Shafique, S. S. Akter, and B. Chakraborty. "Intensive Nutrition Education with or without Supplementary Feeding Improves the Nutritional Status of Moderately-Malnourished Children in Bangladesh." *Journal of Health, Population, and Nutrition* 23 (2006): 320-30.
- Ruel, M. T. "The Natural History of Growth Failure: Importance of Intrauterine and Postnatal Periods." In *Nutrition and Growth*, edited by R. Martorell and F. Haschke, pages 123-158. Nestlé Nutrition Workshop Series, Pediatric Program, Vol. 47. Philadelphia, Pa.: Nestec Ltd., Vevey/Lippincott Williams & Wilkins, 2001.
- Ruel, M. T., and M. Arimond. "Spot-Check Observations for Assessing Hygiene Practices: Review of Experience and Implications for Programs." *Journal of Health, Population, and Nutrition* 20 (2002): 65-76.
- Ruel, M. T., P. Menon, C. Loechl, and G. Peltó. "Fortified Donated Cereal Blends Improve the Nutrient Density of Traditional Complementary Foods, But Are Insufficient to Fill the Iron and Zinc Gaps in 6-8 Month Old Infants in Haiti." *Food and Nutrition Bulletin* 25 (4) (2004): 361-376.
- Schroeder, D. G., R. Martorell, J. A. Rivera, M. T. Ruel, and J.-P. Habicht. "Age Differences in the Impact of Nutritional Supplementation on Growth." *Journal of Nutrition* 125 (1995): 1051S-1059S.
- Schroeder, D. G., H. Pachon, K. A. Dearden, C. B. Kwon, T. T. Ha, T. T. Lang, and D. R. Marsh. "An Integrated Child Nutrition Intervention Improved Growth of Younger, More Malnourished Children in Northern Viet Nam." *Food and Nutrition Bulletin* 23 (4) (2002): 53-61.
- Shrimpton, R., C. G. Victora, M. de Onis, R. C. Lima, M. Blössner, and G. Clugston. "Worldwide Timing of Growth Faltering: Implications for Nutritional Interventions." *Pediatrics* 107 (2001): E75.
- Sletten, P., and W. Egset. *Poverty in Haiti* (2004). FAFO Paper 31. Oslo, Norway: FAFO, 2004. (<http://www.faf.no>).

- Swindale, A., M. Deitchler, B. Cogill, and T. Marchione. *The Impact of Title II Maternal and Child Health and Nutrition Programs on the Nutritional Status of Children*. Washington, D.C.: Food and Nutrition Technical Assistance Project, Academy for Educational Development, 2004.
- UNDP (United Nations Development Programme). *Human Development Report 2005*. New York: United Nations Development Program, 2005.
- UNESCO (United Nations Educational, Scientific, and Cultural Organization) Institute of Statistics. *World Education Indicators*. New York: UNESCO, 2002.
- Waters, H. R. “A Review of the Literature with Applications for Conducting Cost-Effectiveness Studies and for Advocacy.” Support for Analysis and Research in Africa (SARA) Project. Bureau for Africa, Office of Sustainable Development, Washington, D.C., 2000.
- Waters, H. R., M. E. Penny, H. M. Creed-Kanashiro, R. C. Robert, R. Narro, J. Willis, L. E. Caulfield, and R. E. Black. “The Cost-Effectiveness of a Child Nutrition Education Programme in Peru.” *Health Policy and Planning* 21 (4) (2006): 257–64.
- WHO (World Health Organization). *Measuring Change in Nutritional Status: Guidelines for Assessing the Nutritional Impact of Supplementary Feeding Programmes for Vulnerable Groups*. Geneva: World Health Organization, 1983.
- WHO (World Health Organization). *Child Growth Standards: length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: Methods and Development*. Geneva: World Health Organization, 2006.
- WHO (World Health Organization). *Vaccines and Biologicals. WHO Vaccine-Preventable Diseases: Monitoring System. 2002 Global Summary*. Geneva: World Health Organization, 2002.
- WHO (World Health Organization). *Guiding Principles on Feeding Non-Breastfed Children 6-24 Months of Age*. Geneva: World Health Organization, 2005.
- World Bank. “Inequality in Latin America and the Caribbean – Breaking into history?” World Bank Latin America and Caribbean Studies, World Bank, Washington, D.C., 2003.
- World Bank. *Repositioning Nutrition as Central to Development*. World Bank: Washington, D.C., 2006.
- WV-Haiti, Inc. Haiti. “FY 2002-FY 2006 Development Activity Program Proposal.” April 2001.

ANNEXES


ANNEX 2.1 LIST OF HOUSEHOLD SURVEY MODULES

	Baseline Survey	Final Survey
1.	Household composition and socioeconomic status	Household composition and socioeconomic status
2.	Education, civil status and employment of the respondent mother ^a and her spouse	Education, civil status and employment of the respondent mother and her spouse
3.	Maternal prenatal and delivery care and childcare and feeding practices ^b	Maternal prenatal and delivery care and childcare and feeding practices
4.	Child health and immunization	Child health and immunization
5.	Child, maternal, and household cleanliness	Child, maternal, and household cleanliness
6.	Respondent mother's empowerment and decisionmaking	Respondent mother's empowerment and decisionmaking
7.	Household food security	Household food security
8.	Knowledge and attitudes about child feeding and care	Knowledge and attitudes about child feeding and care
9.	Respondent mother's mental and physical health	Respondent mother's mental and physical health
10.	Not assessed at baseline	Participation in the food assisted MCHN program for the index child, the child's younger sibling (if any), other sibling in the 12-41 month age range (if any), and the child's mother (if currently pregnant).
11.	Not assessed at baseline	Attendance at Mothers' Clubs for the index child, the child's younger sibling (if any), or any other sibling in the 12-41 month age range (if any)
12.	Not assessed at baseline	Use of Rally Post services for the index child and younger sibling
13.	Not assessed at baseline	Awareness, trial and adoption of key practices recommended by the program
14.	Not assessed at baseline	Knowledge of infant and young child topics taught at the Mothers Clubs.


^a The terms "respondent mother" and "caregiver" are used interchangeably throughout this document because all main caregivers of the index child were the respondent mothers.

^b Some of the questions on child feeding, appetite and feeding during illness, and child cleanliness, were also asked about the younger child in the household in cases where there was one.

ANNEX 3.1 ORGANIZATIONAL STRUCTURE OF THE HEALTH COMPONENT OF THE PROGRAM



ANNEX 3.2 ORGANIZATIONAL STRUCTURE OF THE COMMODITIES COMPONENT OF THE PROGRAM



ANNEX 4.1 INFANT AND CHILD FEEDING PRACTICES IN HAITI COMPARED TO BEST PRACTICES, AND CONSTRAINTS AND OPPORTUNITIES FOR BEHAVIOR CHANGE IN CENTRAL PLATEAU

(reproduced from Menon et al. 2002b)

Goals	Practices to promote	Practices in Haiti	Facilitating conditions for behavior change	Issues that may affect capacity for behavior change
A. INFANT FEEDING FROM 0-6 MONTHS OF AGE				
Exclusive Breastfeeding (BF)				
<ul style="list-style-type: none"> ➤ Ensure exclusive BF ➤ Prevent bacterial contamination 	<ul style="list-style-type: none"> ➤ Early initiation of exclusive BF (EBF) ➤ Feeding of colostrum ➤ BF on demand ➤ Avoidance of pre- and post-lacteal feeds ➤ Using expressed breast milk if needed ➤ Avoidance of baby-bottles 	<p><i>Positive:</i></p> <ul style="list-style-type: none"> ➤ BF widely practiced ➤ Reported to be mostly on demand <p><i>Non optimal:</i></p> <ul style="list-style-type: none"> ➤ Pre-lacteals and post-lacteal liquids and gruels widely used ➤ Complementary liquids and foods introduced at a <u>very young age</u> ➤ Widespread use of baby bottles <p><i>Not enough information:</i></p> <ul style="list-style-type: none"> ➤ Timing of initiation of BF^a ➤ Colostrum use^a 	<ul style="list-style-type: none"> ❖ Experienced, successful positive deviant mothers (who EBF) exist in communities ❖ Positive deviant mothers had received information from health agents, media, health center staff ❖ EBF moms report it is cheaper to EBF and child is healthier ❖ No objection to use of expressed breast milk – some mothers do it; but training needed 	<ul style="list-style-type: none"> ➤ Water-based liquids and teas given to treat colic (<i>gaz</i>) ➤ Gruels given because mothers need to leave home for work or other activities ➤ Mothers' time and employment constraints ➤ Mothers are concerned about feeling too weak and depleted if they EBF ➤ Concept of <i>let cho</i> (prevents mothers from breastfeeding, but seems to be only in the short term) ➤ Use of expressed breast milk is rare, milk expression unknown in some areas
B. FEEDING PRACTICES FOR INFANTS AND YOUNG CHILDREN 6-24 MONTHS OF AGE				
Continued breastfeeding				
<ul style="list-style-type: none"> ➤ Ensure sustained, frequent, on demand BF up to 24 months of age and beyond 	<ul style="list-style-type: none"> ➤ Continue to BF frequently and on demand ➤ Using expressed breast milk if needed ➤ Avoidance of baby-bottles 	<p><i>Positive:</i></p> <ul style="list-style-type: none"> ➤ Mothers traditionally continue to BF up to 24 months of age <p><i>Non-optimal:</i></p> <ul style="list-style-type: none"> ➤ Widespread use of baby bottles <p><i>Not enough information:</i></p> <ul style="list-style-type: none"> ➤ Mothers may not always BF on demand because of need to leave home for work or other tasks ➤ Children whose mothers are frequently absent may not receive sufficient nutrients from breast milk 	<ul style="list-style-type: none"> ❖ No objection to expression of breast milk, but training needed 	<p>No need for behavior change, but continue promotion of continued BF up 24 months and beyond</p> <p><i>Potential constraints to frequent, on demand BF:</i></p> <ul style="list-style-type: none"> ➤ Mothers do need to leave home to work and/or go to markets ➤ Milk expression rarely practiced, unknown in some areas

Annex 4.1 (continued)

Goals	Practices to promote	Practices in Haiti	Facilitating conditions for behavior change	Issues that may affect capacity for behavior change
Complementary Foods				
➤ Provide foods to complement breast milk and to allow adequate intake of energy and micronutrients	<ul style="list-style-type: none"> ➤ Feed child special energy- and nutrient-dense foods of appropriate texture and consistency for age ➤ From 6 months on, gradually increase amounts and quantity of foods as child gets older ➤ Increase number of times child is fed CF as he/she gets older (at least 2-3 times/d for 6-8 mo old; 3-4 times/d for 9-24 mo old) ➤ Feed a variety of foods (gradually increase variety with age); animal foods should be eaten daily, or as often as possible 	<p><i>Non optimal:</i></p> <ul style="list-style-type: none"> ➤ Complementary foods (CF) of low energy and very low nutrient-density ➤ Variety of foods seems low; animal foods consumed infrequently and in small amounts; low intake of vitamin A fruits and vegetables ➤ Frequency of feeding is low (2-3 times/d) and does not seem to increase with age ➤ Evening meal not fed to young children ➤ No “special” complementary food for child; gruels are consumed by all family members 	<ul style="list-style-type: none"> ❖ No cultural barriers to feeding young children animal foods ❖ Mothers know that eggs, liver are good for child ❖ Mothers usually feed child when they are present ❖ Mothers leave prepared food for child when they have to leave ❖ Good recognition of importance of fluid replacement during diarrhea 	<ul style="list-style-type: none"> ➤ Lack of availability and access to food, especially animal foods and micronutrient-rich fruits and vegetables ➤ Overall poverty, lack of economic resources ➤ Poor access to water, sanitation, health services ➤ Time constraints of caregivers to prepare “special foods” ➤ Belief that evening meal causes indigestion ➤ Lack of recognition of importance of high feeding frequency for young children ➤ Belief that children are ready for family foods and family meal patterns by 12 months of age ➤ Some cultural barriers to feeding young children specific fruits/vegetables
Feeding during diarrhea				
➤ Continue to BF and feed CF to child during diarrhea; ensure fluid replacement	<ul style="list-style-type: none"> ➤ Increase fluid intake during illness, including more frequent breastfeeding, and encourage the child to eat soft, varied, appetizing, favorite foods. ➤ After illness, give food more often than usual and encourage the child to eat more. 	<p><i>Positive practices:</i></p> <ul style="list-style-type: none"> ➤ Mothers continue to BF and give liquids when child has diarrhea <p><i>Non-optimal practices</i></p> <ul style="list-style-type: none"> ➤ Mothers reduce feeding of CF during diarrhea 	<ul style="list-style-type: none"> ❖ Knowledge about fluid replacement can be used to introduce issues of encouraging consumption of food during and after an episode of illness and providing special foods for recuperation 	

Annex 4.1 (continued)

Goals	Practices to promote	Practices in Haiti	Facilitating conditions for behavior change	Issues that may affect capacity for behavior change
Responsive feeding				
➤ Practice responsive feeding, applying the principles of psychosocial care	<ul style="list-style-type: none"> ➤ Feeding with a balance between giving assistance and encouraging self feeding, as appropriate to the child's level of development ➤ Feeding with positive verbal encouragement, without verbal or physical coercion. ➤ Feeding with age-appropriate, as well as culturally appropriate, eating utensils ➤ Feeding in a protected and comfortable environment ➤ Feeding in response to early hunger cues ➤ Feeding by an individual with whom the child has a positive emotional relationship and who is aware of and sensitive to the child's individual characteristics, including his or her changing physical and emotional states 	<p><i>Positive practices</i></p> <ul style="list-style-type: none"> ➤ Child is usually fed in a separate bowl or plate rather than common plate ➤ Fathers seem involved in childcare and feeding <p><i>Insufficient information at this point on:</i> Psychosocial care (needs to be assessed through observations, which was beyond the scope of the present study)</p>	<ul style="list-style-type: none"> ❖ Some aspects of responsive feeding are already practiced (e.g., feeding in a separate bowl, involving fathers in care) 	

^a Data on these practices were obtained from the baseline survey carried out in 2002 but were not available at the time of the formative research study.

ANNEX 4.2 IDENTIFICATION OF PROGRAMMATIC OPTIONS TO ADDRESS THE CONSTRAINTS TO INFANT FEEDING, AND TO SUPPORT FACILITATING FACTORS

Constraints/facilitators to appropriate infant feeding:	Program options within current structure and delivery system: (BCC, food donations and other interventions delivered by the WV-Haiti Programme at Rally Posts, food delivery points, and Mothers' Clubs)	Program options that will require NEW program structure or complementary interventions:
A. INFANT FEEDING FROM 0-6 MONTHS OF AGE		
Exclusive breastfeeding		
<i>Factors that constrain exclusive breastfeeding</i>		
➤ Water-based liquids and teas given to treat colic (<i>gaz</i>)	➤ BCC program: ensuring women are given enough knowledge and confidence about dealing with infant colic	
➤ Gruels given because mothers need to leave home for work (economic, reasons) or other activities (mothers have multiple household responsibilities; time constraints)	➤ BCC program: training mothers in the use of expressed breast milk ➤ Food aid component (could aid in delaying extremely early resumption of work outside home)	➤ Microcredit programs targeting women and increasing their potential involvement in income-generating activities at (or close to) home ➤ Long-term poverty reduction strategies needed ➤ Public transportation and road improvement projects ➤ Childcare support
➤ Mothers are concerned about feeling too weak and depleted if they EBF	➤ BCC program: can address need to replace fluids frequently when EBF ➤ Food aid component can possibly alleviate weakness due to poor quality diet and lack of food	➤ Fathers' Clubs: ensure that fathers are sensitized to the need for EBF women to be supported. ➤ Agriculture production activities to increase availability/access to food
➤ Concept of <i>let cho</i> (prevents mothers from breastfeeding, but seems to be only in the short term)	➤ BCC program	
➤ Use of expressed breast milk is rare, milk expression unknown in some areas	➤ BCC program: ensuring adequate training in the use and appropriate storage of expressed breast milk	
<i>Factors that facilitate exclusive breastfeeding</i>		
➤ Experienced, successful positive deviant mothers (who EBF) exist in communities	➤ BCC program: use mothers' clubs as support groups to encourage EBF	
➤ Positive deviant mothers had received information from health agents, media, health center staff	➤ BCC program: ensure that mothers receive the same information from different sources	
➤ EBF moms report it is cheaper to EBF and child is healthier	➤ BCC program: use benefits of EBF on household medical expenses as a motivator	

Constraints/facilitators to appropriate infant feeding:	Program options within current structure and delivery system: (BCC, food donations and other interventions delivered by the WV-Haiti Programme at Rally Posts, food delivery points, and Mothers' Clubs)	Program options that will require NEW program structure or complementary interventions:
B. FEEDING PRACTICES FOR INFANT AND YOUNG CHILDREN 6-24 MONTHS OF AGE		
Continued breastfeeding		
<i>Factors that constrain continued breastfeeding</i>		
➤ Mothers need to leave home to go to work and/or markets	➤ BCC program: promote and encourage continued and sustained breastfeeding up to 24 months of age	➤ Microcredit programs targeting women and increasing their potential involvement in income-generating activities at (or close to) home ➤ Long-term poverty reduction strategies needed ➤ Public transportation and road improvement projects ➤ Childcare support
➤ Expression of breast milk rarely practiced	➤ BCC program: ensuring adequate training in the use and appropriate storage of expressed breast milk	
<i>Factors that facilitate continued breastfeeding</i>		
➤ Mothers traditionally breastfeed up to 18 or 24 months	➤ BCC program: Promote and encourage continued and sustained breastfeeding up to 24 months of age	
Complementary foods		
<i>Factors that constrain feeding of optimal complementary foods</i>		
➤ Time constraints of caregivers to prepare "special foods"	➤ BCC Program: Promote easy to prepare, time efficient recipes and ideas for nutritious CF	➤ Need for public transport and road projects that can ensure that women spend more time commuting to place of work. ➤ Engage other family members (grandmothers, fathers, etc.) in BCC program to provide more support to mothers
➤ Belief that evening meal causes indigestion	➤ BCC program: encourage feeding of at least gruels at night, rather than juices or teas	
➤ Lack of recognition of importance of high feeding frequency for young children	➤ BCC Program: ensure mothers are sensitized to higher feeding frequency needs of infants and young children	

Constraints/facilitators to appropriate infant feeding:	Program options within current structure and delivery system: (BCC, food donations and other interventions delivered by the WV-Haiti Programme at Rally Posts, food delivery points, and Mothers' Clubs)	Program options that will require NEW program structure or complementary interventions:
➤ Belief that children are ready for family foods and family meal patterns by 12 months of age	➤ BCC program: Need to ensure that children 12-23 are given adequate attention and appropriate foods (use brain development as a motivator)	
➤ Some cultural barriers to feeding young children specific fruits/vegetables	➤ BCC program: encourage trials of small amounts of these foods	
➤ Lack of access to micronutrient rich foods, especially animal foods and micronutrient-rich fruits and vegetables	➤ BCC program: encourage use of small amounts of meat, liver, or eggs for children ➤ Encourage consumption of goat milk (especially among goat owners)	➤ Livestock projects to increase access to animal source foods ➤ Livestock care projects to improve health of animals and milk production ➤ Home garden promotion; solar drying of fruits/vegetables ➤ Microcredit programs to facilitate income generation through livestock rearing ➤ Market interventions to encourage sale of small pieces of meat and liver
➤ Overall poverty, lack of economic resources	➤ Food aid component can help somewhat	➤ Overall community development projects and poverty reduction interventions
➤ Poor access to water, sanitation, health services		➤ Community development projects for improving water, sanitation, etc.
<i>Factors that facilitate feeding of optimal complementary foods</i>		
➤ No cultural barriers to feeding young children animal foods; and mothers are aware that eggs and liver are good for young child	➤ BCC program: encourage and support feeding of animal foods to young children	➤ Same as above (3 rows up): Livestock projects and market interventions to increase availability and access to animal source foods
➤ Mothers leave prepared food for child when they have to leave	➤ BCC program: encourage preparation of enriched recipes rather than traditional low nutrient density gruels/juices	

Constraints/facilitators to appropriate infant feeding:	Program options within current structure and delivery system: (BCC, food donations and other interventions delivered by the WV-Haiti Programme at Rally Posts, food delivery points, and Mothers' Clubs)	Program options that will require NEW program structure or complementary interventions:
Feeding during diarrhea		
<i>Factors that constrain optimal feeding during and after diarrhea</i>		
➤ Feeding of CF during illness is decreased	➤ BCC program: encourage caregivers to continue attempts to feed children during illness ➤ BCC program: stress the need for extra food and the use of enriched recipes when children are recovering from illness	
<i>Factors that facilitate optimal feeding during and after diarrhea</i>		
➤ Good recognition of importance of fluid replacement during diarrhea	➤ BCC program: encourage caregivers to sustain fluid replacement with ORS and other safe fluids when child has diarrhea	➤ Community development projects for improving water quality and sanitation ➤ Ensuring availability of and access to ORS
Responsive Feeding		
<i>Factors that could constrain responsive feeding</i>		
➤ Mothers' time and workload constraints	➤ BCC program: encourage mothers to entrust adult members and inform them about responsive feeding as well	
<i>Factors that facilitate responsive feeding</i>		
➤ Fathers seem involved in childcare and feeding		➤ Ensure that fathers are engaged in BCC program as well through Fathers' Clubs and sensitized to responsive feeding practices
➤ Mothers usually feed child when they are present	➤ BCC program: encourage responsive feeding and encourage mothers to entrust adult members to feed child when possible	

ANNEX 4.3 BCC STRATEGY MATRIX FOR A BCC PROGRAM TO PREVENT MALNUTRITION AMONG CHILDREN BETWEEN 0-24 MONTHS

Behaviors to promote	Who will messages be targeted to?	When will messages be delivered?	Where will communication be delivered ?	How will communication be delivered ?	What is needed to help with communication ?
0-5 Months: Exclusive breastfeeding 1. Initiate breastfeeding immediately after the child is born 2. Give the child colostrum (and avoid <i>lòk</i>) 3. Breastfeed exclusively (avoid other liquids and foods) 4. Breastfeed frequently, on demand 5. Use expressed breast milk as needed (avoid other liquids and foods) 6. Use a cup and spoon to feed the infant expressed breast milk (avoid baby bottles) 7. Increase the frequency of breastfeeding when the infant is sick	Pregnant mothers Midwives Grandmothers Health professionals in health centers (pre-natal care) Lactating mothers Fathers Grandmothers	Pregnancy, before delivery At delivery First 1-2 months of lactation	➤ Pre- and postnatal consultations (can also be group education) ➤ Food distribution ➤ Mothers' Clubs ➤ Fathers' Clubs ➤ Markets	➤ Individual counseling at pre-natal and postnatal consultation ➤ Group education of pregnant women at health centers (if they invite all pregnant women to attend on a particular day for the pre-natal control) ➤ Discussion and problem solving related to breastfeeding at Mother's Clubs ➤ Discussions on support needs with fathers at Father's Clubs ➤ Home visits	➤ <i>Training of pre-natal and postnatal counseling staff</i> (health staff as well as midwives) in communication methods and content of practices to encourage ➤ <i>Training of health agents and colvols</i> in group discussion and problem solving methods ➤ <i>Provision of resource materials for communication</i> (counseling cards, flip charts, other visual material) ➤ <i>Make attendance at pre- and postnatal consultation and mothers' clubs mandatory</i> in order to receive food
6-8 Months: Complementary feeding and continued breastfeeding 1. Continue to breastfeed on demand and use expressed breast milk as necessary 2. Gradually introduce enriched porridges, gruels and special foods (enriched using beans, eggs, fish, breast milk, milk, pumpkin, etc.) 3. Feed the infant enriched foods 2-3 times per day	Lactating mothers Fathers Grandmothers Lactating mothers Fathers	First 2-3 months of lactation During 3-9 months of	➤ Rally Posts ➤ Food distribution centers ➤ Mothers' Clubs ➤ Father's Clubs	➤ Group education at Rally Posts ➤ Recipe demonstrations at Food Distribution Points ➤ Recipe trials at Mothers' Clubs ➤ Group discussions	➤ <i>Training of agents de santé and colvols</i> in group discussion and problem solving methods ➤ <i>Training of health agents and colvols in recipe trials/ demonstrations</i> to be

Behaviors to promote	Who will messages be targeted to?	When will messages be delivered?	Where will communication be delivered ?	How will communication be delivered ?	What is needed to help with communication ?
4. Increase the quantity of enriched foods as the child grows older 5. Feed nutritious snacks (like <i>cham cham</i> , fruits, peanut butter) 1-2 times per day 6. Use a cup and spoon to feed the infant expressed breast milk and other liquids (avoid baby bottles) 7. Feed infants directly, and feed slowly and patiently. 8. Encourage children to eat, but do not force them; if children refuse many foods, experiment with different food combinations, tastes, textures and positive methods of encouragement 9. Minimize distractions during meals 10. Talk to children during feeding, with eye to eye contact 11. Increase frequency of breastfeeding and liquids when the infant is ill 12. Feed the child his or her favorite foods when ill 13. Increase the frequency of feeding and feed more enriched foods when the infant is convalescing after an illness	Grandmothers	lactation		and problem solving at Mother's Clubs ➤ Group discussions with fathers at Father's Clubs? ➤ Home visits	implemented at food distribution points and mothers' clubs ➤ <i>Provision of resource materials for communication</i> (counseling cards, flip charts, etc.) ➤ <i>Make attendance at postnatal consultation and mothers' clubs mandatory</i> in order to receive food
9-23 Months : Complementary feeding and continued breastfeeding 1. Continue to breastfeed on demand and use expressed breast milk as necessary 2. Continue to feed enriched porridges, gruels, and special foods (enriched using beans, eggs, fish, breast milk, milk, pumpkin, etc.) 3. Increase the variety of foods fed to the infant by adding other family foods to the child's diet	Lactating mothers Fathers Grandmothers	9 months onwards	➤ Food distribution centers ➤ Rally Posts ➤ Mothers' Clubs ➤ Father's Clubs	➤ Group education at Rally Posts ➤ Recipe demonstrations at Food Distribution Points ➤ Recipe trials at Mothers' Clubs ➤ Group discussions and problem solving	➤ <i>Training of health agents and colvols</i> in group discussion and problem solving methods ➤ <i>Training of health agents and colvols in recipe trials/ demonstrations</i> to be implemented at food

Behaviors to promote	Who will messages be targeted to?	When will messages be delivered?	Where will communication be delivered ?	How will communication be delivered ?	What is needed to help with communication ?
4. Feed the infant enriched porridges/gruels or special foods 3-4 times per day 5. Increase the quantity of food as the child grows older 6. Feed nutritious snacks (like cham cham, fruits, peanut butter) 1-2 times per day 7. Use a cup and spoon to feed the infant expressed breast milk and other liquids (avoid baby bottles) 8. 9-11 months: Feed infants directly, and feed slowly and patiently 9. 12-23 months: Assist and supervise feeding to ensure adequate intake, and feed slowly and patiently 10. Encourage children to eat, but do not force them; if children refuse many foods, experiment with different food combinations, tastes, textures and positive methods of encouragement 11. Minimize distractions during meals 12. Talk to children during feeding, with eye to eye contact 13. Increase frequency of breastfeeding and liquids when the infant is ill 14. Feed the child his or her favorite foods when ill 15. Increase the frequency of feeding and feed more enriched foods when the infant is convalescing after an illness				at Mother's Clubs ➤ Group discussions with fathers at Father's Clubs? ➤ Group education at Community Health sessions ➤ Home visits	distribution points ➤ <i>Provision of resource materials for communication</i> (counseling cards, flip charts, etc.) ➤ <i>Make attendance at post-natal consultation, Rally Posts and Mothers' Clubs mandatory</i> in order to receive food

Behaviors to promote	<i>Who</i> will messages be targeted to?	<i>When</i> will messages be delivered?	<i>Where</i> will communication be delivered ?	<i>How</i> will communication be delivered ?	<i>What</i> is needed to help with communication ?
All the above behaviors + overall attention and focus on the under-two child	Entire community		<ul style="list-style-type: none"> ➤ Community meetings ➤ Radio ➤ Posters ➤ Other 	<ul style="list-style-type: none"> ➤ Sensitization to various practices at the community meetings ➤ Radio messages ➤ Community posters 	<ul style="list-style-type: none"> ➤ Utilization of community leaders and pastors for sensitization in churches and at community meetings? ➤ Design of effective posters ➤ Identification of key venues to display posters ➤ Design of effective radio messages/ stories/ songs/dialogues ➤ Identification of key radio stations and key times during the day to broadcast messages

ANNEX 4.4 SCHEDULES OF LEARNING SESSIONS AND TOPICS AT MOTHERS' CLUBS

A. Schedule of learning sessions at Mothers' Clubs (for pregnant and lactating women)

Month of Pregnancy	Mothers' Clubs for pregnant women
5	Other topics: Diet for pregnant women
6	Other topics: Dangerous signs during pregnancy
7	Other topics: Preparation of child delivery
8	<u>Session 1</u> Importance of breastfeeding (initiation of breastfeeding, exclusive breastfeeding, continue breastfeeding until 2 years of age or beyond) Discouragement of bottle use Comparison of recommendations with local beliefs and practices
9	<u>Session 2</u> Initiation of breastfeeding, importance of colostrum Exclusive breastfeeding until 6 months Position and attachment of the baby during feeding, frequency of breastfeeding
Child age (months)	Mothers' Clubs for lactating women
1	<u>Session 3</u> Sharing experience with exclusive breastfeeding Review of exclusive breastfeeding Review of position and attachment of the child during feeding, frequency of breastfeeding and care of nipples and breasts Expression of breast milk Drinking water while breastfeeding
2	<u>Session 4</u> Sharing experiences related to drinking water while breastfeeding and expression of breast milk Sharing experience related to exclusive breastfeeding Discussing constraints/problems related to exclusive breastfeeding and offering solutions
3	<u>Session 5</u> Exclusive breastfeeding and Lactational Amenorrhea Method (LAM)
4	<u>Session 6</u> Introduction of complementary foods when children are about 6 months old Importance of continued breastfeeding until 2 years of age or beyond
5	<u>Session 7</u> Overview on child development and feeding chart (for children 6-12 months of age: food consistency, participating in feeding, frequency, quantity of food) Learning how to eat Important information about the first food (in addition to breast milk) given to children Preparation of the next session: preparing nutritious foods
6	<u>Session 8</u> Preparing nutritious foods/cooking session Tasting and discussion Repetition of learning how to eat <u>Session 7</u> Preparation of the next session: preparing nutritious foods

B. Schedule of learning sessions at Mothers' Clubs (for mothers of 6-24 month old children)

Month in the program	Mothers of 6 to 24 months old children
1	<u>Session 8</u> Preparing nutritious foods/cooking session Tasting and discussion Repetition of learning how to eat
2	<u>Session 9</u> Repetition on child development and feeding chart (for children 0 to 12 months of age) Helping children to eat Feeding during and after illness Preparation of the next session: variety of food
3	<u>Session 10</u> Sharing experience with one new feeding practice (related to helping children to eat) Variety of food Sharing experience with preparing nutritious foods at home
4	<u>Session 11</u> Hygiene in food preparation, handling and storage – Diarrhea prevention Feeding during and after illness
5	<u>Session 12</u> Child development and feeding chart (for children 12-23 months of age) Discussing food variety issues (special complementary foods, fruits and vegetables, vitamin A-rich foods, animal foods, evening meal) Preparing a creative way to communicate one feeding recommendation of the child development and feeding chart
6	<u>Session 13 (P)</u> Causes of malnutrition Different types of malnutrition Recuperation of moderately malnourished children
7	Other topics: Diarrhea
8	Other topics: Immunization
9	Other topics: Hygiene
10	Other topics: Use of Moringa Oleifera
11	Other topics: HIV/AIDS
12	Other topics: Family Planning
13	Other topics: Home gardening
14	<u>Session 12</u> Child development and feeding chart (for children 12-23 months of age) Discussing food variety issues (special complementary foods, fruits and vegetables, vitamin A-rich foods, animal foods, evening meal) Preparing a creative way to communicate one feeding recommendation of the child development and feeding chart
15	Other topics: HIV/AIDS
16	Other topics: Family Planning
17	Other topics, placement in schedule to be determined
18	Other topics, placement in schedule to be determined

C. Schedule of learning sessions at Mothers' Clubs for mothers of malnourished children 6-to-59 months of age (recuperative program group)

Month in the program	Mothers of malnourished children
1	<u>Session 13</u> Causes of malnutrition Different types of malnutrition Recuperation of moderately malnourished children Introduction to the child development and feeding chart Preparation for the next session: preparing nutritious foods
2	<u>Session 8</u> Preparing nutritious foods/cooking session Tasting and discussion Repetition of learning how to eat
3	<u>Session 9</u> Repetition on child development and feeding chart (for children 0 to 11 months of age) Helping children to eat Feeding during and after illness Preparation of the next session: variety of food
4	<u>Session 10</u> Sharing experience with one new feeding practice (related to helping children to eat) Variety of food Sharing experience with preparing nutritious foods at home
5	<u>Session 12</u> Child development and feeding chart (for children 12 months to 5 years of age) Discussing food variety issues (special complementary foods, fruits and vegetables, vitamin A-rich foods, animal foods, evening meal) Preparing a creative way to communicate one feeding recommendation of the child development and feeding chart
6	<u>Session 1</u> Importance of breastfeeding (initiation of breastfeeding, exclusive breastfeeding, continue breastfeeding until 2 years of age or beyond) Discouraging bottle use Comparison of recommendations with local beliefs and practices Other topics: Immunization
7	<u>Session 11</u> Hygiene in food preparation, handling and storage—Diarrhea prevention Feeding during and after illness Other topics: Diarrhea
8	Other topics: HIV/AIDS
9	Other topics: Family Planning

ANNEX 5.1 SUMMARY MATRIX FROM CONSULTATIVE WORKSHOP, FEBRUARY 2003

Summary of discussions held in Haiti with WV-Haiti staff on the findings from the first round of operations research, and follow-up actions identified

A - RALLY POSTS (RPs)

Aspects/activities	Constraint(s) identified	Potential corrective action(s)	Feasibility of implementing corrective action(s)	Potential impact of improving this aspect on effectiveness and impact of program
Organization	Too crowded, too many beneficiaries; ratio of participants/staff is too high	<p>Increase number of RPs</p> <p>Increase staff (AS/<i>colvols</i>)</p> <p>Continue to work on a case-by-case basis to split the RPs that are really too big and can be split</p> <p>Train <i>colvols</i> better so that they can help the health agents more</p> <p>Improve venue of the RPs by providing basic furniture (chairs, benches, tables, etc.)</p>	<p>Not possible at this point; health agents already have too many RPs/month</p> <p>Not feasible in short-term because of program criteria and lack of resources</p> <p>Feasible; already being done; should be continued</p> <p>This is already being done; <i>colvols</i> have been included in all recent training; salary has also been increased and they are now referred to as "health promoter agents"</p> <p>This will be done through mobilization of community resources</p>	<p>- Will reduce staff and participants time burden and frustrations</p> <p>- May increase participation at RPs, which, in turn, may increase impact of program</p> <p>- May increase quality of services at RPs</p> <p>Will increase assistance that health agents receive and reduce their time burden</p> <p>Could make the RPs more efficient and more pleasant for staff and participants</p>

Aspects/activities	Constraint(s) identified	Potential corrective action(s)	Feasibility of implementing corrective action(s)	Potential impact of improving this aspect on effectiveness and impact of program
	Registration is a major bottleneck; sequence of activities varies a lot between RPs and does not usually follow the implementation plan	The following sequence was recommended for the future: 1) give a number to each participant 2) provide general education session 3) weigh child, plot weight on growth chart in health card 4) do general registration (including register weight in registry) 5) evaluate nutritional status; 6) inform and advise mother (see below in GMP) 7) give required immunization, deworming, Vitamin A, and ORS	Feasible; the team has discussed several possible sequences and concluded that this one was probably the best choice; supervisors will assist health agents/ <i>colvols</i> in implementing and testing this sequence	See above
Specific activities (organization and quality)				
1. Education	Still mostly one session, first thing in the morning and therefore many participants miss the session	Give at least two education sessions and if needed, up to three	Feasible; was done in some of the RPs observed; the supervisors will be key in motivating the health staff to comply with this recommendation	Greater coverage of participating mothers; this may increase maternal knowledge and reinforce certain topics discussed at the MCs, thereby strengthening the BCC strategy
	Quality of education is non-optimal (messages tend to be vague, little use of visual materials, limited interaction between staff and participants) Monthly education topics recommended by the program are rarely used	May be difficult to improve, given crowded, busy, noisy environment; may be best to focus on simple topics and messages of general interest (e.g., ORS, immunization, family planning, etc.) Ensure that the calendar of topics is distributed to all supervisors; also ensure that supervisors work closely with health staff during their monthly meetings to reinforce adherence to monthly schedule of topics.	Feasible to focus on simple topics, ensuring that information is accurate and messages are clear and simple; supervisors will have to monitor this. Feasible; requires greater communication between health staff and their supervisors and monitoring of implementation of the calendar of topics	Following the 12-month calendar of education topic will allow mothers to receive education on the full set of topics considered important and of general interest to mothers by the program. This can increase their general knowledge of health, nutrition and hygiene.

Aspects/activities	Constraint(s) identified	Potential corrective action(s)	Feasibility of implementing corrective action(s)	Potential impact of improving this aspect on effectiveness and impact of program
2. Vitamin A/deworming/ORS	Many participants do not receive the services and this seems to be due largely to a lack of supply of the products	This problem is due to lack of availability from the main providers (Ministry of Health, UNICEF, etc.)	This problem is outside of the program's control, but efforts should be pursued to ensure a constant supply	These preventive health services are very important for child's health, growth, development and survival. Therefore it is important to achieve high coverage.
3. Growth monitoring and promotion	Children are weighed, their weight is recorded, and their nutritional status assessed, but there is little use of the information to communicate with the mother and to give her tailored individual advice regarding her child's growth and progress	Retrain personnel to ensure that each mother receives the following information: 1) Child's weight 2) Child's nutritional status 3) Child's progress (gained or lost weight) 4) Brief message of encouragement (if child is growing well) and recommendations for follow-up with other program activities (e.g., MCs) if child is not growing well. If child is M3, make sure s/he is scheduled for a home visit.	Feasible; will require that supervisors motivate staff and monitor that these actions are implemented	The information provided to mothers about their child's nutritional status and growth can help raise awareness among mothers and motivate them to engage more actively in the BCC activities and to adopt the recommended practices.
	The quality of the weighing and plotting varies; errors result in the misclassification of some children into the different Gomez categories, and thus in errors in screening beneficiary children (true only for the recuperative group, which uses children's nutritional status as a screening criteria)	Retrain health staff to improve the quality of the weighing (making sure that needle of the hanging scale is still before recording the weight will help). Retrain staff to improve their skills in plotting the child's weight in the growth chart.	Very feasible; can be achieved through good training and supervision	Improving the weighing and plotting of the weights in the growth chart will reduce misclassification errors, thereby improving the cost-effectiveness of the program.
4. Immunization	Lack of supplies (similar to the problems of vitamin A, ORS, and deworming tablets)	See above (vitamin A, ORS, and deworming)	See above (vitamin A, ORS, and deworming)	See above (vitamin A, ORS, and deworming)

B - MOTHERS' CLUBS (MCs)

Aspects/activities	Constraint(s) identified	Potential corrective action(s)	Feasibility of implementing corrective action(s)	Potential impact of improving this aspect on effectiveness and impact of program
Organization	Problems with reorganization of MCs for different subgroups of mothers/children (especially in preventive group)	Ensure that supervisors understand the rationale for the reorganization of the clubs, and continue to reinforce importance of implementing this reorganization as planned. Slightly modify current system, which is to form two groups with 6-23 month old (in preventive group): 6-8 and 9-23. Replace with the following two groups: 6-11 and 12-23.	Very feasible, but requires close supervision to ensure that groups are formed as recommended by the program and that mothers attend the groups they are assigned to.	The importance of the timely delivery of the education for the effectiveness of the BCC cannot be overemphasized. A much greater impact on behavior change is expected if the intervention reaches the beneficiaries at their best learning moment. Achievement of behavior change may, in turn, reduce childhood malnutrition.
Quality of the education and BCC intervention	Very good quality overall; marked improvements since development of BCC strategy and training of staff; performance varies, however, and continued improvements can still be achieved	Strengthen supervision and ongoing training of staff; strengthen the communication of concepts and technical content of the sessions, as well as the utilization of the communication material and the adult education techniques. Continue to use the supervision checklists to provide feedback to the staff and to monitor progress.	Very feasible; is already being done; important to continue to strengthen the supervision and feedback provided to the staff. Supervisors also need to monitor that enthusiasm and motivation of the staff, and interest of beneficiaries are maintained over time.	The greater the quality of the BCC in MCs, the greater the potential impact of the program on improved child feeding and care and consequently on children's nutritional status, health, and development.

C - FOOD DISTRIBUTION POINTS (FDPs)

Aspects/activities	Constraint(s) identified	Potential corrective action(s)	Feasibility of implementing corrective action(s)	Potential impact of improving this aspect on effectiveness and impact of program
Organization	Very large number of beneficiaries per distribution point and high beneficiary/staff ratios	Some organizational aspects have already been addressed by the program and some are in the process of being resolved; the staff makes every effort to improve the logistics of the food distribution process, including improving local storage facilities.	Difficult to solve completely, because most logistical problems are inherent to the rough environmental conditions of the area.	Improving the logistics of food distribution could reduce the staff and beneficiary time burden and increase satisfaction.
	Important delays in arrival of food and staff at FDP			
	Very high time burden for beneficiaries (including travel time and time spent at the FDP) and staff			
Quality	Amount of food received by beneficiaries is not always the amount allocated by the program	A system is already in place to verify that the quantity of food received by the beneficiaries corresponds to the amount allocated.	Feasible; system is already in place. Need to ensure that check is done systematically on a sample of beneficiaries as they exit the FDP.	Could increase beneficiary satisfaction and reduce food losses.
	Communication difficulties have been reported between commodity and health staff and commodity staff and beneficiaries	These problems are addressed on an ongoing basis, as they occur, by the supervisors or regional coordinators.	Very possible, already being addressed.	Could improve staff satisfaction with their work and beneficiary satisfaction with the program.
	Certain problems of hygiene in the manipulation of the food during the distribution process have been identified	Suggestions made to provide gloves to food handlers and small compensation to motivate them to adopt more hygienic practices. Use of plastic sheets and cleaning of measurement tools have already been adopted. Possibility of acquiring large containers with a tap for the distribution of oil was also recommended.	Feasible, but requires financial means to acquire the materials proposed.	Would reduce the risk of contamination of the food.

ANNEX 5.2 MOTHERS' CLUBS OBSERVATION CHECKLIST ITEMS

For each domain, the items listed are the positive behaviors included on the checklist.

Technical content

- Information conveyed is correct and complete (according to session-specific observation form).
- Give accurate responses to most or all questions from participants.
- When incorrect information is expressed by participants, bring focus back to promoted behavior.
- Health promoter/assistant acknowledges when a question is beyond his/her technical knowledge.

Session management and organization

- Complete all session steps.
- Complete all steps in order.
- Do not read aloud instructions meant for staff only.
- Do read Crèole session information for group with no difficulty.
- Complete session within +/- 15 minutes of the planned length.
- Have all materials ready and organized before session.

Teaching and facilitation skills

- Use small groups as suggested in session guide.
- Define the question/topic to be discussed.
- Help arrange participants to face each other.
- Circulate around the room to help.
- Ask for group reports.
- Use open-ended questions as in session guide.
- Use open-ended questions to probe and encourage at other times.
- Speak loudly and clearly.
- Show most or all visuals planned for the session.
- Ensure that all participants can see the visuals.
- Use other teaching/facilitation techniques per session guide.

Attitudes displayed (with observer to explicitly note how attitude was demonstrated)

- Provide praise/affirmation to participants.
- Demonstrate respect for participants.
- Help women feel at ease with participating.
- Attempt to create good dialogue and limit lecture-style presentation.

Atmosphere at the learning session

- No distracting side-conversations during session
- No disruptive children, who continue to stay in the room
- No other distractions (trucks driving by, etc.)
- Laughter during the session
- Use animations with the group.
- Participants do not appear bored.

ANNEX 5.3 QUESTIONS FOR STAFF RELATED TO MOTIVATION

For each of these statements, could you please indicate HOW OFTEN you, personally, have felt this way – **ALWAYS, OFTEN, SOMETIMES, RARELY or NEVER.**

1. Do you feel like you enjoy your work as a health agent/*colvol*?
2. Do you feel that the program will improve people's lives?
3. Do you feel confident that you are prepared for your responsibilities in the program?
4. Do you feel that Mothers' Clubs are not useful to poor women because they don't have the money to change their behavior?
5. Do you feel like you are responsible for so many activities that it's not possible to do them all well in the time you have?
6. Do you feel like you are very connected with other health agents/*colvols* in the program?
7. Do you feel well supported by the other health agents/*colvols* you work with?
8. Do you feel that if you need help running your Rally Post, you can always get help from the other health agents/*colvols* whom you know. (*ask only for health agents*)?
9. How often have you had thoughts of leaving this job?

This time, for each of these statements, could you please indicate HOW STRONGLY you, personally, AGREE with the statement? **Do you strongly agree, agree, agree somewhat, disagree, or strongly disagree.**

10. I am contributing to improving the conditions of the communities I am working in.
11. Mothers' Clubs help changing behaviors even of poor women in a positive direction.
12. I receive adequate training to meet my current responsibilities.
13. I feel like I receive refresher training as often as I need it.
14. I am very much involved personally in my job.
15. I have a lot of pressure in this job. It really seems like the workload keeps increasing.
16. I find my job to be motivating and I like to do it.
17. I feel that the program management values the work that we do in the communities.
18. I am satisfied with my salary when I compare this job to other similar jobs.
19. I am satisfied with the amount I am paid for the work I do.
20. I feel secure that I will not lose my job in the near future.
21. The program beneficiaries value our efforts to improve their lives.
22. The program beneficiaries care much more for the food than for our efforts to improve their practices.
23. The other health promoters/health promoter assistants respect the work that I do.
24. Would you want to leave this job if it were possible/if you had other alternatives?

ANNEX 5.4 QUESTIONS FOR STAFF RELATED TO SUPERVISION

For each of the statements, please think about your direct supervisor and indicate HOW OFTEN you have experienced each of the following behaviors with your supervisor. Tell me if you experience this **always, often, sometimes, rarely, or never.**

1. My supervisor keeps me informed about the follow-up of my concerns/worries.
2. My supervisor informs me about upcoming trainings/meetings, etc., in a timely fashion.
3. My supervisor respects my fixed monthly activities when planning other meetings.
4. My supervisor consults with me before making changes to the activities that I am involved in, e.g., the re-organization of the MCs.
5. My supervisor takes my concerns into account when planning activities that involve me.
6. When I make a mistake on the job, my supervisor scolds me.
7. My supervisor praises me when I do something really well.
8. I feel that my supervisor takes my concerns up to the higher management level.
9. My supervisor helps me to organize my time and activities in an efficient manner.
10. My supervisor uses times when I make mistakes or don't perform well as opportunities to help me improve my skills.
11. When my supervisor visits me at the Mothers' Club, she uses the checklist to observe how I conduct the session.
12. When my supervisor visits me at the Mothers' Club, she uses the checklist to provide me with feedback on the session.
13. When my supervisor visits me at the Rally Post, she inspects the registers that I use.
14. When my supervisor visits me at the Rally Post, she provides me with feedback on my work there.

For each of the statements, please think about your direct supervisor and indicate HOW STRONGLY each of the statements captures the behavior or attitude of your supervisor. Please tell me if you **strongly agree, agree, agree somewhat, disagree, or strongly disagree.**

15. I feel well informed by my supervisor about changes/modifications to the program activities that I am involved in.
16. I feel that my supervisor devotes enough time and effort for the preparation of BCC activities each month.
17. My supervisor respects my fixed monthly activities when planning other meetings.
18. My supervisor ensures that I have enough of the supplies that I need to do my daily work.
19. When I disagree with my supervisor I feel safe to express my opinion.
20. Having a supervisor visit me at the MC/RP motivates me to do a better job there.

21. The way the supervisor provides feedback on my performance at the MC inspires me to be my best.
22. My supervisor's commitment to her job encourages me to do my best.
23. My supervisor takes into account/considers my suggestions to improve things.
24. My supervisor works with me to identify solutions to program activity related problems.
25. I feel that my supervisor is sympathetic to my problems/cares about my problems.
26. I constantly learn new things about maternal and child health from my supervisor (technical information).
27. My supervisor gives me enough guidance and structure to help me do my job.
28. My supervisor uses times when I make mistakes or don't perform well as opportunities to help me improve my skills.
29. I feel that my supervisor helps me plan not just my job but also my future.

ANNEX 8.1 SCORING OF QUESTIONS IN INFANT AND YOUNG CHILD FEEDING KNOWLEDGE TEST

	Measured knowledge item	Variable name	Scoring
A. Appropriate age of introduction of new foods			
1.	Water/other liquids - water - other liquids		<p>For each food group (liquids, semisolids, staples, vegetables, eggs, meats): Introduction of any food in the food group at 6-8 mo=+1 None of foods introduced in the appropriate period of 6-8 mo=0 (i.e., introduction was either too early or too late).</p> <p><u>Food group variables created:</u> <i>Knowliqd, knowsemi, knowstap, knowveg, knowegg, knowmeat.</i></p> <p>Values for each food group variable were then summed up to create an overall scale.</p>
2.	Semi-solid foods - bread soup - gruels - bean sauce		
3.	Staple foods - rice - millet - cornmeal		
4.	Vegetables - vegetables added to the food - vegetables cooked on their own		
5.	Eggs - egg yolk - whole egg		
6.	Meats - chicken - fish - meat		
	Total possible score	Intrknow	Minimum=0 Maximum=6
B. Appropriate feeding frequency score			
1.	No. of meals/day for a 6-8 mo old child		0 to 1 meal/day = 0 2 and higher =1
2.	No. of snacks/day for a 6-8 mo old child		0 snacks/day = 0 1 and higher =1
3.	No. of meals/day for a 9-11 mo old child		0 to 2 meals/day = 0 3 and higher =1
4.	No. of snacks/day for a 9-11 mo old child		0 snacks/day = 0 1 and higher =1
5.	No. of meals/day for a 12-24 mo old child		0 to 2 meals/day = 0

	Measured knowledge item	Variable name	Scoring
			3 and higher =1
6.	No. of snacks/day for a 12-24 mo old child		0 snacks/day = 0 1 and higher =1
	Total possible score	apprfreq	Minimum=0 Maximum=6
B. Overall feeding knowledge scale, weighted			
1.	Knowledge of appropriate introduction of new foods, weighted		<i>Intrknow</i> /3 – brings contribution of the knowledge on appropriate introduction to 2 points in total (range 0 to +2)
2.	Knowledge scale on appropriate feeding frequency, weighted		<i>Apprfreq</i> /3 – brings contribution of the knowledge on appropriate feeding frequency to 2 points in total (range 0 to +2)
3.	Age until when a child should be breastfed		0-5 mo=-2 6 to 11 mo=0 12 to 17 mo=0.5 18 thru 23=1 24 and beyond=2
	Total possible score	Feedknow	Minimum=-2 Maximum=+6

ANNEX 8.2 SCORING OF QUESTIONS IN GENERAL NUTRITION AND HEALTH KNOWLEDGE TEST (RESPONSES THAT WERE CONSIDERED APPROPRIATE)

Knowledge question	Variable name	Responses considered acceptable Scoring = 1 for ANY acceptable response (exceptions are noted below)
1. Until what age (months) do you think you should breast feed your child ?	<i>Q604</i>	≥ 24 months
2. How many times per day should a child 6-8 months old eat?	<i>Q605</i> <i>Q606</i>	A. Meals: 2 or more _ per day B. Snacks: _1 or more per day
3. How many times per day should a child 9-11 months old eat?	<i>Q607</i> <i>Q608</i>	A. Meals: 2 or more _ per day B. Snacks: _1 or more per day
4. How many times per day should a child older than 12 months eat?	<i>Q609</i> <i>Q610</i>	A. Meals: _3 or more per day B. Snacks: _2 or more _ per day
5. How long after birth should a baby start breastfeeding?	<i>Q611</i>	Immediately Less than 1 hour after
6. What should a mother do with the “first milk” or colostrum?	<i>Q612</i>	Give it to her baby by breastfeeding soon after birth
7. If a mother thinks her baby is not getting enough breast milk, what should she do?	<i>Q613</i>	Breastfeed more often/more frequently Mother needs to drink more water (assigned only 0.5 point)
8. Do you think that infants under 6 months of age should be given water in addition to breast milk if the weather is very hot?	<i>Q614</i>	No
9. If a mother has a young baby (less than 6 months) and needs to be away from her baby and the baby gets hungry, what should the baby be fed?	<i>Q615</i>	Mother’s expressed breast milk
10. Can a mother who has a young baby and who is not well fed produce enough breast milk?	<i>Q616</i>	Yes
11. What are the things you can do to increase milk production?	<i>Q617</i>	Breastfeed more often/more frequently Mother needs to drink more water (assigned only 0.5 point)

Knowledge question	Variable name	Responses considered acceptable Scoring = 1 for ANY acceptable response (exceptions are noted below)
12. What are special foods mothers could make for her children to complement breast milk? (Focus was on evaluating knowledge about special foods developed through recipe trials)	Q618	Enriched gruel with breast milk Enriched gruel with other kinds of milk Enriched gruel with egg Enriched gruel with beans and g/nuts Mashed plantain with pumpkin and dried fish sauce
13. Will feeding a child enriched gruel in the evening cause indigestion?	Q619	No
14. At meal times, how much food should a child 6-9 months old be offered/served each time you feed him/her? (Using a local measure)	Q620	¼ of a Gode 7 ½ of a Gode 7 ¾ of a Gode 7 One Gode 7 More than one Gode 7
15. Do you think that a 1-year old child should eat only the same foods as the rest of the family?	Q621	No
16. At meal times, how much food should a child older than 12 months be offered/served each time you feed him/her?	Q622	½ of a Gode 7 ¾ of a Gode 7 One Gode 7 More than one Gode 7
17. Do you think children need an extra meal per day after they have been sick?	Q623	Yes
18. For how long do children need an extra meal per day after they have been sick?	Q624	2 weeks More than 2 weeks
19. What should you do when your child has diarrhea? (ORS scored separately from dietary changes)	Q625	Give ORS/home-prepared solution
20. What should you do when your child has diarrhea? (Scoring of dietary changes)	Q625	Feed more food than usual Give more liquids than usual Continue breastfeeding Breastfeed more often

Knowledge question	Variable name	Responses considered acceptable Scoring = 1 for ANY acceptable response (exceptions are noted below)
21. What can you do to protect a child against polio?	<i>Q626</i>	Vaccinate your child
22. What are ways to protect yourself against AIDS?	<i>Q627</i>	Abstain from sex Stay with one partner Don't have many partners Stay away from contaminated blood Avoid used needles Don't use another person's razor Always use a condom
23. When should you wash your hands?	<i>Q628</i>	Before eating After using the toilet Before feeding the child After cleaning a child who has defecated
24. What are the ways to protect a child from getting worms?	<i>Q629</i>	Wash hands Cut nails Children should wear pants Wash fruits and vegetables Children should wear sandals Give them treated water
25. What are the ways to make drinking water safe?	<i>Q630</i>	Boil water Treat with chlorine
26. What are some of the things to encourage young children to eat their food?	<i>Q631</i>	Feed slowly and patiently Talk to the child Reduce distractions Feed other foods Change flavor of the food
27. Why do you think children get malnourished?	<i>Q632</i>	Don't eat enough food/poor appetite Don't eat frequently Child is ill (diarrhea, infection, etc.) Child is weaned abruptly Child is not fed with affection (0.5 point) Unbalanced meals Insufficient quantity of food

Knowledge question	Variable name	Responses considered acceptable Scoring = 1 for ANY acceptable response (exceptions are noted below)
28. What are some of the ways in which we can help malnourished children recuperate?	<i>Q633</i>	Increase the amount fed at each meal Feed the child more frequently Feed tasty foods, foods the child likes Encourage the child to eat If sick, take to clinic/health agent/etc If marasmus/kwashiorkor, take to clinic/doctor Give them balanced meals Give them enough food

ANNEX 8.3A GUIDING PRINCIPLES FOR COMPLEMENTARY FEEDING OF THE BREASTFED CHILD

Excerpted from PAHO/WHO (2003).

1. Duration of exclusive breastfeeding & age of introduction of complementary foods

Practice exclusive breastfeeding from birth to 6 months of age, and introduce complementary foods at 6 months of age while continuing to breastfeed.

2. Maintenance of breastfeeding

Continue frequent, on-demand breastfeeding until 2 years of age or beyond.

3. Responsive feeding

Practice responsive feeding, applying the principles of psycho-social care. Specifically: a) feed infants directly and assist older children when they feed themselves, being sensitive to their hunger and satiety cues; b) feed slowly and patiently, and encourage children to eat, but do not force them; c) if children refuse many foods, experiment with different food combinations, tastes, textures and methods of encouragement; e) minimize distractions during meals if the child loses interest easily; f) remember that feeding times are periods of learning and love - talk to children during feeding, with eye to eye contact.

4. Safe preparation and storage of complementary foods

Practice good hygiene and proper food handling by a) washing caregivers' and children's hands before food preparation and eating, b) storing foods safely and serving foods immediately after preparation, c) using clean utensils to prepare and serve food, d) using clean cups and bowls when feeding children, and e) avoiding the use of feeding bottles, which are difficult to keep clean.

5. Amount of complementary food

Start at six months of age with small amounts of food and increase the quantity as the child gets older, while maintaining frequent breastfeeding. The energy needs from complementary foods for infants with average breastmilk intake in developing countries are approximately 200 kcal per day at 6-8 months of age, 300 kcal per day at 9-11 months of age, and 550 kcal per day at 12-23 months of age. In industrialized countries these estimates differ somewhat (130, 310 and 580 kcal/d at 6-8, 9-11 and 12-23 months, respectively) because of differences in average breastmilk intake.

6. Food consistency

Gradually increase food consistency and variety as the infant gets older, adapting to the infant's requirements and abilities. Infants can eat pureed, mashed and semi-solid foods beginning at six months. By 8 months most infants can also eat "finger foods" (snacks that can be eaten by children alone). By 12 months, most children can eat the same types of foods as consumed by the rest of the family (keeping in mind the need for nutrient-dense foods, as explained in #8 below). Avoid foods that may cause choking (i.e., items that have a shape and/or consistency that may cause them to become lodged in the trachea, such as nuts, grapes, raw carrots).

7. Meal frequency and energy density

Increase the number of times that the child is fed complementary foods as he/she gets older. The appropriate number of feedings depends on the energy density of the local foods and the usual amounts consumed at each feeding. For the average healthy breastfed infant, meals of complementary foods should be provided 2-3 times per day at 6-8 months of age and 3-4 times per day at 9-11 and 12-24 months of age, with additional nutritious snacks (such as a piece of fruit or bread or chapatti with nut paste) offered 1-2 times per day, as desired. Snacks are defined as foods eaten between meals-usually self-fed, convenient and easy to prepare. If energy density or amount of food per meal is low, or the child is no longer breastfed, more frequent meals may be required.

8. Nutrient content of complementary foods

Feed a variety of foods to ensure that nutrient needs are met. Meat, poultry, fish or eggs should be eaten daily, or as often as possible. Vegetarian diets cannot meet nutrient needs at this age unless nutrient supplements or fortified products are used (see #9 below).

Vitamin A-rich fruits and vegetables should be eaten daily. Provide diets with adequate fat content. Avoid giving drinks with low nutrient value, such as tea, coffee and sugary drinks such as soda. Limit the amount of juice offered so as to avoid displacing more nutrient-rich foods.

9. Use of vitamin-mineral supplements or fortified products for infant and mother

Use fortified complementary foods or vitamin-mineral supplements for the infant, as needed. In some populations, breastfeeding mothers may also need vitamin-mineral supplements or fortified products, both for their own health and to ensure normal concentrations of certain nutrients (particularly vitamins) in their breastmilk. [Such products may also be beneficial for pre-pregnant and pregnant women].

10. Feeding during and after illness

Increase fluid intake during illness, including more frequent breastfeeding, and encourage the child to eat soft, varied, appetizing, favorite foods. After illness, give food more often than usual and encourage the child to eat more.

ANNEX 8.3B GUIDING PRINCIPLES FOR FEEDING NON-BREASTFED CHILDREN 6-24 MONTHS OF AGE

Excerpted from WHO (2005).

1. **Amount of food needed.** Ensure that energy needs are met. These needs are approximately 600 kcal per day at 6-8 months of age, 700 kcal per day at 9-11 months of age, and 900 kcal per day at 12-23 months of age.
2. **Food consistency.** Gradually increase food consistency and variety as the infant gets older, adapting to the infant's requirements and abilities. Infants can eat pureed, mashed and semi-solid foods beginning at six months. By eight months most infants can also eat "finger foods" (snacks that can be eaten by children alone). By 12 months, most children can eat the same types of foods as consumed by the rest of the family (keeping in mind the need for nutrient-dense foods, as explained in #4 below). Avoid foods in a form that may cause choking (i.e., items that have a shape and/or consistency that may cause them to become lodged in the trachea, such as whole nuts, whole grapes or raw carrots, whole or in pieces).
3. **Meal frequency and energy density.** For the average healthy infant, meals should be provided 4-5 times per day, with additional nutritious snacks (such as pieces of fruit or bread or chapatti with nut paste) offered 1-2 times per day, as desired. The appropriate number of feedings depends on the energy density of the local foods and the usual amounts consumed at each feeding. If energy density or amount of food per meal is low, more frequent meals may be required.
4. **Nutrient content of foods.** Feed a variety of foods to ensure that nutrient needs are met.
 - a. Meat, poultry, fish or eggs should be eaten daily, or as often as possible, because they are rich sources of many key nutrients such as iron and zinc. Milk products are rich sources of calcium and several other nutrients. Diets that do not contain animal source foods (meat, poultry, fish or eggs, plus milk products) cannot meet all nutrient needs at this age unless fortified products or nutrient supplements are used.
 - b. If adequate amounts of other animal source foods are consumed regularly, the amount of milk needed is 200-400 ml/d; otherwise, the amount of milk needed is 300-500 ml/d. Acceptable milk sources include full cream animal milk (cow, goat, buffalo, sheep, camel), Ultra High Temperature (UHT) milk, reconstituted evaporated (but not condensed) milk, fermented milk or yogurt, and expressed breast milk (heat treated if the mother is HIV positive).
 - c. If milk and other animal source foods are not eaten in adequate amounts, both grains and legumes should be consumed daily, if possible within the same meal, to ensure adequate protein quality.
 - d. Dairy products are the richest sources of calcium. If dairy products are not consumed in adequate amounts, other foods that contain relatively large amounts of calcium,

such as small fish that include the bones (dried or fresh, with the bones crushed or otherwise processed so that they are safe to eat) and lime treated maize tortillas, can fill the gap. Other foods such as soybeans, cabbage, carrots, squash, papaya, dark green leafy vegetables, guava and pumpkin are useful additional sources of calcium.

- e. The daily diet should include Vitamin A rich foods (e.g. dark coloured fruits and vegetables; red palm oil; vitamin A fortified oil or foods); vitamin C rich foods (e.g. many fruits, vegetables and potatoes) consumed with meals to enhance iron absorption; and foods rich in the B vitamins including riboflavin (e.g. liver, egg, dairy products, green leafy vegetables, soybeans), vitamin B6 (e.g. meat, poultry, fish, banana, green leafy vegetables, potato and other tubers, peanuts) and folate (e.g. legumes, green leafy vegetables, orange juice).
 - f. Provide diets with adequate fat content. If animal source foods are not consumed regularly, 10-20 g of added fats or oils are needed unless a fat rich food is given (such as foods or pastes made from groundnuts, other nuts and seeds). If animal source foods are consumed, up to 5 g of additional fats or oils may be needed.
 - g. Avoid giving drinks with low nutrient value, such as tea, coffee and sugary soft drinks. Limit the amount of juice offered, to avoid displacing more nutrient rich foods.
5. **Use of vitamin-mineral supplements or fortified products.** As needed, use fortified foods or vitamin mineral supplements (preferably mixed with or fed with food) that contain iron (8-10 mg/d at 6-12 months, 5-7 mg/d at 12-24 months). If adequate amounts of animal source foods are not consumed, these fortified foods or supplements should also contain other micronutrients, particularly zinc, calcium and vitamin B12. In countries where vitamin A deficiency is prevalent or where the under-five mortality rate is over 50 per 1000, it is recommended that children 6-24 months old receive a high dose vitamin A supplement (100,000 IU once for infants 6-12 months old and 200,000 IU bi-annually for young children 12-23 months old).
6. **Fluid needs.** Non-breastfed infants and young children need at least 400-600 ml/d of extra fluids (in addition to the 200-700 ml/d of water that is estimated to come from milk and other foods) in a temperate climate, and 800-1200 ml/d in a hot climate. Plain, clean (boiled, if necessary) water should be offered several times per day to ensure that the infant's thirst is satisfied.
7. **Safe preparation and storage of foods.** Practise good hygiene and proper food handling by a) washing caregivers' and children's hands with soap (or a rubbing agent such as ash) before food preparation and eating, b) storing foods safely and serving foods immediately after preparation, c) using clean utensils to prepare and serve food, d) using clean cups and bowls when feeding children, and e) avoiding the use of feeding bottles, which are difficult to keep clean.

8. **Responsive feeding.** Practise responsive feeding, applying the principles of psycho social care. Specifically: a) feed infants directly and assist older children when they feed themselves, being sensitive to their hunger and satiety cues; b) feed slowly and patiently, and encourage children to eat, but do not force them; c) if children refuse many foods, experiment with different food combinations, tastes, textures and methods of encouragement; e) minimize distractions during meals if the child loses interest easily; f) remember that feeding times are periods of learning and love talk to children during feeding, with eye to eye contact.
9. **Feeding during and after illness.** Increase fluid intake during illness and encourage the child to eat soft, varied, appetizing, favourite foods. After illness, give food more often than usual and encourage the child to eat more.

ANNEX 8.4 AWARENESS, TRIAL, AND ADOPTION OF KEY RECOMMENDED PRACTICES, BY PROGRAM GROUP AND PARTICIPATION (EVER)

Recommended practices	Awareness				Trial				Adoption			
	Ever participated		Never participated		Ever participated		Never participated		Ever participated		Never participated	
	Prev. N=622	Recup. N=626	Prev. N=126	Recup. N=126	Prev. N=622	Recup. N=626	Prev. N=126	Recup. N=126	Prev. N=622	Recup. N=626	Prev. N=126	Recup. N=126
Leaving expressed breast milk when going out	97.6 ^a	98.2 ^a	73.8	81.0	65.1 ^{a b}	59.3 ^a	22.2	24.6	47.0 ^a	41.9 ^a	16.7	18.3
Adding breast milk to gruel	74.4 ^{a b}	67.9 ^a	17.5	26.2	31.7 ^{a b}	24.0 ^a	3.2	4.0	19.0 ^a	15.3 ^a	2.4	2.4
Adding an egg to child's gruel	96.3 ^{a b}	89.1*	38.1	45.2	72.5 ^b	53.7 ^a	76.2	70.6	47.1 ^{a b}	34.4 ^a	11.9	9.5
Preparing gruel with beans & nuts	88.9 ^{a b}	78.9 ^a	32.5	27.0	45.5 ^{a b}	35.9 ^a	7.9	4.0	25.9 ^{a b}	18.5 ^a	6.4	2.4
Feeding enriched gruel in evening	95.3 ^{a b}	90.4 ^a	24.6 ^c	40.5	76.7 ^{a b}	67.1 ^a	14.3	15.9	56.6 ^{a b}	48.1 ^a	13.5	11.9
Preparing mashed plantain with pumpkin	97.6 ^{a b}	94.9*	54.8	55.6	79.7 ^{a b}	67.3 ^a	36.5	29.4	54.5 ^{a b}	46.2 ^a	31.0 ^c	17.5
Feeding an extra meal after illness	96.0 ^{a b}	92.5	50.0	46.8	90.7 ^{a b}	84.0	77.8	81.4				

* Failed to converge with random effects regression analysis—differences were highly significant ($p < 0.001$) using the logit model.

^a Difference between participants and nonparticipants (within program group) is significant at $p < 0.05$ (random effects regression analysis).

^b Difference between program groups is significant at $p < 0.05$ for participants (random effects regression analysis).

^c Difference between program groups is significant at $p < 0.05$ for nonparticipants using random effects regression models.

ANNEX 8.5 PERCENT OF INFANTS FED ACCORDING TO RECOMMENDATIONS DURING EARLY POSTNATAL PERIOD, BY INTERVENTION AREA AND BY PARTICIPATION

	Variable name	Preventive			Recuperative			All children
		Ever exposed	Not exposed	All	Exposed	Not exposed	All	
		Percent						
Index children (12-41 mo)		(n = 622)	(n = 126)	(n = 748)	(n = 626)	(n = 126)	(n = 752)	(n = 1,500)
Breastfed within 1 hour ^a	<i>bf1hr</i>	72.9 ^d	28.6 ^d	64.6	71.2 ^d	39.3 ^d	65.8	65.2
Only breast milk on first day	<i>onlybf</i>	95.3 ^d	63.5 ^{d, e}	90.0	93.1 ^d	77.0 ^{d, e}	90.4	90.2
Gave colostrum	<i>q325n</i>	90.8 ^d	52.4 ^{d, e}	84.4*	90.9 ^d	75.4 ^{d, e}	88.3*	86.3
Younger sibling (0-11 mo)		(n = 152)	(n = 7) ^b	(n = 159)	(n = 171)	(n = 14) ^b	(n = 185)	(n = 344)
Breastfed within 1 hour ^a	<i>bf1hr</i>	85.4	--	84.1	84.9 ^d	-- ^d	81.9	83.0
Only breast milk on first day	<i>onlybf</i>	98.7	--	98.1 ^c	94.2	--	93.0	95.3 ^c
Gave colostrum	<i>q325n</i>	95.4	--	95.0	93.6	--	94.1	94.5

* Significant at $p \leq 0.05$. P-value for comparison of preventive and recuperative areas; results from xtlogit for dichotomous variables.

^a The sample for this variable includes 264 missing cases (the original coding is 88) most likely due to recall problem (to be confirmed with field staff).

^b The result is not presented due to small sample size.

^c P-value is not available because the maximization procedure failed to converge to a solution.

^d Difference between participants and nonparticipants (within program group) is significant at $p < 0.05$ (random effects regression analysis).

^e Difference between program groups is significant at $p < 0.05$ for nonparticipants (random effects regression analysis).

ANNEX 8.6 AGE OF INTRODUCTION OF SELECTED LIQUIDS AND FOODS, BY INTERVENTION AREA AND BY PARTICIPATION

	Prev + exp	Prev + not exp	All prev	Recp + exp	Recp + not exp	All recp	All children
	Percent						
Index children (12-41 mo)	(n = 622)	(n = 126)	(n = 748)	(n = 626)	(n = 126)	(n = 752)	(n = 1,500)
<i>At age of child started giving other liquids (variable name: othliq)</i>							
0-5.9 mo	11.9	65.1	20.9	16.0	46.8	21.1	21.0
6-8.9 mo ^a	87.6 ^{b, c}	34.9 ^{b, d}	78.7	84.0 ^{b, c}	53.2 ^{b, d}	78.9	78.8
9-11.9 mo	0.0	0.0	0.0	0.0	0.0	0.0	0.0
12 mo or older	0.5	0.0	0.4	0.0	0.0	0.0	0.2
Not yet	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<i>At age of child started giving semi-solid foods (variable name: semisol)</i>							
0-5.9 mo	8.8	52.4	16.2	12.1	42.9	17.3	16.7
6-8.9 mo ^a	90.8 ^b	46.8 ^b	83.4	87.5 ^b	57.1 ^b	82.4	82.9
9-11.9 mo	0.2	0.0	0.1	0.3	0.0	0.3	0.2
12 mo or older	0.2	0.8	0.3	0.0	0.0	0.0	0.1
Not yet	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<i>At age of child started giving solid (variable name: solid)</i>							
0-5.9 mo	3.5	25.4	7.2	5.1	25.4	8.5	7.9
6-8.9 mo ^a	71.4	50.8	67.9	71.1	59.5	69.1	68.5
9-11.9 mo	15.4	16.7	15.6	14.9	8.7	13.8	14.7
12 mo or older	9.6	7.1	9.2	8.9	6.3	8.5	8.9
Not yet	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<i>At age of child started giving meats (variable name: meat)</i>							
0-5.9 mo	3.1	21.4	6.1	4.5	19.8	7.0	6.6
6-8.9 mo ^a	70.1	45.2	65.9	65.0	55.6	63.4	64.7
9-11.9 mo	13.3	15.9	13.8	14.5	11.1	14.0	13.9
12 mo or older	13.5	17.5	14.2	16.0	13.5	15.6	14.9
Not yet	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<i>At age of child started giving eggs (variable name: eggs)</i>							
0-5.9 mo	4.3	20.6	7.1	4.9	19.1	7.3	7.2
6-8.9 mo ^a	78.3 ^b	49.2 ^{b, d}	73.4	72.5	60.3 ^d	70.4	71.9
9-11.9 mo	8.4	10.3	8.7	12.0	6.3	11.1	9.9
12 mo or older	9.0	19.8	10.8	10.6	14.3	11.2	11.0
Not yet	0.0	0.0	0.0	0.0	0.0	0.0	0.0

Note: Shaded area indicates n in cell size <= 20.

^a We tested feeding pattern between exclusively breastfed versus given more than just breast milk (or not yet breastfed). P-value for comparison of preventive and recuperative areas; results are from xtlogit for dichotomous variables.

^b Difference between participants and nonparticipants (within program group) is significant at p < 0.05 (random effects regression analysis).

^c Difference between program groups is significant at p < 0.05 for participants (random effects regression analysis).

^d Difference between program groups is significant at p < 0.05 for nonparticipants (random effects regression analysis).

ANNEX 8.7 PERCENT OF CHILDREN WHO WERE FED AT LEAST THE MINIMUM RECOMMENDED NUMBER OF TIMES,^a AND MEAN NUMBER OF MEALS AND SNACKS, BY AGE GROUP, CURRENT PROGRAM PARTICIPATION, AND BREASTFEEDING STATUS

	Variable name	Non-breast			Breastfed			All children		
		Not current	Current participation	All	Not current	Current participation	All	Not current	Current participation	All
Index children		(n = 843)	(n = 141)	(n = 984)	(n = 270)	(n = 246)	(n = 516)	(n = 1,113)	(n = 387)	(n = 1,500)
Age group 12-23 mo										
- % with minimum number of meals	<i>mealfrq2</i>	8.3 ^b	22.8 ^b	14.4	54.8	62.7 ^b	58.9	39.9	52.8 ^b	46.2
- mean number of meals	<i>numsol</i>	2.7 ^b	3.0 ^b		2.6 ^b			2.6 ^b		
- mean number of snacks	<i>numsnack</i>	2.2	2.5	2.4	2.0	2.2	2.1	2.1 ^b	2.3 ^b	2.2
Age group 24 mo and older										
- % with minimum number of meals ^c	<i>mealfrq2</i>	-	- 2.8	-	- 2.8	- 2.7	-	- 2.8	- 2.7	-
- mean number of meals	<i>numsol</i>	2.7	2.6	2.7	2.7	- ^d		2.7	2.6	2.7
- mean number of snacks	<i>numsnack</i>	2.2	2.5	2.2	2.3	- ^d	2.3	2.2	2.5	2.2
Younger siblings		(n = 0)	(n = 0)	(n = 0)	(n = 80)	(n = 35)	(n = 115)	(n = 80)	(n = 35)	(n = 115)
Age group 6-11 mo										
- % with minimum number of meals	<i>mealfrq2</i>	-	-	-	61.3	68.6 ^{2.7}	63.6	61.3	68.6 ^b	63.6
- mean number of meals	<i>numsol</i>	-	-	-	2.2 ^b			2.2 ^b		
- mean number of snacks	<i>numsnack</i>	-	-	-	1.6	2.1	1.8	1.6	2.1	1.8

^a Give recommendations by age/breastfeeding status and cite PAHO/WHO 2003 and WHO 2005.

^b Difference between participants and nonparticipants is significant at $p < 0.05$ (random effects regression analysis). 2.3 2.7 2.3

^c The result is not applicable as the minimum recommended number of meals only covers only children 6-23 months.

^d The result is not reported due to small sample size.

ANNEX 8.8 PERCENT OF CHILDREN WHO CONSUMED SELECTED FOOD GROUPS IN THE PREVIOUS 24 HOURS, BY INTERVENTION AREA AND PARTICIPATION

Consumption of selected food groups	Variable name	Prev + exp	Prev + not exp	All prev	Recp + exp	Recp + not exp	All recp	All children
					(percent)			
Index children (12-41 mo)		(n = 622)	(n = 126)	(n = 748)	(n = 626)	(n = 126)	(n = 752)	(n = 1,500)
All cereals	<i>grains</i>	98.1	97.6	98.0	97.4	95.2	97.1	97.5
Roots, tubers, starchy vegetables	<i>rootstub</i>	34.9	24.6	33.2	37.4	35.7	37.1	35.2
Legumes	<i>legumes</i>	87.1	88.1	87.3	84.0	84.9	84.2	85.7
Vitamin A-rich vegetables	<i>vafrveg</i>	72.0 ^a	73.0	72.2*	67.9 ^a	64.3	67.3*	69.7
All other fruits and vegetables	<i>othveg</i>	38.0 ^a	40.5	38.4***	31.0 ^a	27.0	30.3***	34.3
Milk and formula	<i>dairy</i>	28.2 ^a	22.2	27.2**	21.7 ^a	18.3	21.1**	24.1
Meat, fish, and egg	<i>meategg</i>	64.1 ^a	54.8	62.5***	54.5 ^a	49.2	53.6***	58.1
Nuts	<i>nuts</i>	23.8	19.0	23.0	24.9	19.8	24.1	23.5

* Significant at $p \leq 0.05$. P-value for comparison of preventive and recuperative areas; results from xtlogit for dichotomous variables.

** Significant at $p \leq 0.01$. P-value for comparison of preventive and recuperative areas; results from xtlogit for dichotomous variables.

*** Significant at $p \leq 0.001$. P-value for comparison of preventive and recuperative areas; results from xtlogit for dichotomous variables.

^a Difference between program groups is significant at $p < 0.05$ for participants (random effects regression analysis).

ANNEX 8.9 PERCENT OF CHILDREN WHO CONSUMED SELECTED ANIMAL SOURCE FOODS IN THE PREVIOUS 24 HOURS, BY INTERVENTION AREA AND PARTICIPATION

Consumption of selected animal products	Variable name	Prev + exp	Prev + not exp	All prev	Recp + exp	Recp + not exp	All recp	All children
					(percent)			
Index children (12-41 mo)		(n = 622)	(n = 126)	(n = 748)	(n = 626)	(n = 126)	(n = 752)	(n = 1,500)
Eggs	<i>egg</i>	22.7 ^{a, b}	11.1 ^a	20.7*	18.2 ^{a, b}	6.3 ^a	16.2*	18.5
Chicken	<i>chicken</i>	7.6	8.7	7.8	5.6	5.6	5.6	6.7
Fish and seafood	<i>fish</i>	36.3 ^b	26.2	34.7	30.2 ^b	30.2	30.2	32.4
Beef and pork	<i>beefpork</i>	28.5 ^b	28.6	28.4***	20.9 ^b	21.4	21.0***	24.7
Heart and liver	<i>liver</i>	5.5	2.4	5.0	4.2	4.0	4.1	4.5

* Significant at $p \leq 0.05$. P-value for comparison of preventive and recuperative areas; results from xtlogit for dichotomous variables.

*** Significant at $p \leq 0.001$. P-value for comparison of preventive and recuperative areas; results from xtlogit for dichotomous variables.

^a Difference between participants and nonparticipants (within program group) is significant at $p < 0.05$ (random effects regression analysis).

^b Difference between program groups is significant at $p < 0.05$ for participants (random effects regression analysis).

ANNEX 8.10 FREQUENCY OF CONSUMPTION OF NUTRIENT-RICH FOODS, INCLUDING RECIPES PROMOTED IN MOTHERS' CLUBS, BY INTERVENTION AREA AND PARTICIPATION (MOTHER EVER PARTICIPATED)

Consumption of nutrient-rich foods	Variable name	Prev + exp (n = 622)	Prev + not exp (n = 126)	All prev (n = 748)	Recp + exp (n = 626)	Recp + not exp (n = 126)	All recp (n = 752)	All children (n = 1,500)
(percent)								
Index children (12-41 mo)								
Foods/groups consumed 3 or more times in the last 7 days								
Eggs	<i>eggs3</i>	14.5	13.5	14.3	13.3	6.3	12.1	13.2
Flesh foods (meat, poultry, fish, seafood)	<i>flesh3</i>	74.7 ^{b, c}	61.9 ^b	72.6***	60.5 ^c	57.9	60.1***	66.3
Vitamin A-rich orange/red fruits/vegetables	<i>orange3</i>	46.5	50.8	47.3	48.4	38.9	46.8	47.0
Recipes promoted in Mothers' Clubs:								
Gruel with WSB	<i>grwsb3</i>	32.6 ^{b, c}	4.8 ^b	27.9***	17.4 ^{b, c}	2.4 ^b	14.9***	21.3
Gruel with eggs	<i>gregg3</i>	6.8	0.0	5.6***	2.4	0.0	2.0***	3.8
Gruel with milk	<i>grmilk3</i>	12.7 ^c	6.3	11.6*	8.6 ^c	6.3	8.2*	9.9
Mashed plantain w/pumpkin and fish sauce	<i>mplant3</i>	4.7 ^c	0.8	4.0***	1.3 ^c	0.0	1.1***	2.5
Mashed vegetables	<i>mvege3</i>	6.8 ^c	3.2	6.2*	3.5 ^c	2.4	3.3*	4.7
Gruel with dried fish	<i>grfish3</i>	3.4	0.8	2.9	2.4	0.0	2.0	2.5
Gruel with beans/groundnuts	<i>grbean3</i>	1.0	0.8	0.9 ^a	0.8	0.0	0.7 ^a	0.8
Other food prepared with WSB	<i>owsb3</i>	9.8 ^{b, c}	4.0 ^b	8.8**	6.1 ^c	1.6	5.3**	7.1
Any of the above promoted in Mothers' Clubs	<i>recfood3</i>	68.1 ^{b, c}	36.5 ^{b, d}	62.8***	44.4 ^{b, c}	24.6 ^{b, d}	41.1***	51.9
Foods/groups consumed 7 or more times in the last 7 days								
Eggs	<i>eggs7</i>	3.4 ^c	0.8	2.9	1.6 ^c	0.8	1.5	2.2
Flesh foods (meat, poultry, fish, seafood)	<i>flesh7</i>	34.6 ^{b, c}	17.5 ^b	31.7***	22.8 ^c	19.8	22.3***	27.0
Vitamin A-rich orange/red fruits/vegetables	<i>orange7</i>	26.7	36.5	28.4	31.5	23.8	30.2	29.3
Any of the recipes promoted in Mothers' Clubs	<i>recfood7</i>	35.9 ^{b, c}	7.9 ^{b, d}	31.2***	17.4 ^{b, c}	1.6 ^{b, d}	14.8***	22.9

* Significant at $p \leq 0.05$. P-value for comparison of preventive and recuperative areas; results from xtlogit for dichotomous variables.

** Significant at $p \leq 0.01$. P-value for comparison of preventive and recuperative areas; results from xtlogit for dichotomous variables.

*** Significant at $p \leq 0.001$. P-value for comparison of preventive and recuperative areas; results from xtlogit for dichotomous variables.

^a P-value for comparison of preventive and recuperative areas is not available because the maximization procedure failed to converge to solution.

^b Difference between participants and nonparticipants (within program group) is significant at $p < 0.05$ (random effects regression analysis).

^c Difference between program groups is significant at $p < 0.05$ for participants (random effects regression analysis).

^d Difference between program groups is significant at $p < 0.05$ for nonparticipants (random effects regression analysis).

**ANNEX 8.11 FREQUENCY OF CONSUMPTION OF NUTRIENT-RICH FOODS, INCLUDING RECIPES
PROMOTED IN MOTHERS' CLUBS, BY INTERVENTION AREA AND
CURRENT PARTICIPATION (CHILDREN 6-42 MONTHS)**

Consumption of nutrient-rich foods	Variable name	Prev + curr	Prev + not curr	All prev	Recp + curr	Recp + not curr	All recp	All children
		(n = 218)	(n = 467)	(n = 748)	(n = 106)	(n = 646)	(n = 752)	(n = 1,500)
(percent)								
Index children (12-41 mo)								
Foods/groups consumed 3 or more times in the last 7 days								
Eggs	<i>eggs3</i>	17.1	12.6	14.3	15.1	11.6	12.1	13.2
Flesh foods (meat, poultry, fish, seafood)	<i>flesh3</i>	79.0 ^{a, b}	68.7 ^{a, c}	72.6***	69.8 ^{a, b}	58.5 ^{a, b}	60.1***	66.3
Vitamin A-rich orange/red fruits/vegetables	<i>orange3</i>	44.8	48.6	47.3	57.6	45.1	46.8	47.0
Recipes promoted in Mothers' Clubs:								
Gruel with WSB	<i>grwsb3</i>	58.6 ^a	4 ^{a, b}	27.9***	58.5 ^a	7.7 ^a	14.9***	21.3
Gruel with eggs	<i>gregg3</i>	9.3 ^a	3.4 ^{a, c}	5.6***	8.5 ^a	0.9 ^{a, c}	2.0***	3.8
Gruel with milk	<i>grmilk3</i>	17.1 ^a	8.4 ^a	11.6*	18.9	6.5	8.2*	9.9
Mashed plantain w/pumpkin and fish sauce	<i>mplant3</i>	7.5 ^{a9.}	1.9 ^a	4.0***	2.8	0.8	1.1***	2.5
Mashed vegetables	<i>mvege3</i>	6.4	6.0 ^c	6.2*	7.6 ^a	2.6 ^{a, c}	3.3*	4.7
Gruel with dried fish	<i>grfish3</i>	4.3	2.1	2.9	3.8	1.7	2.0	2.5
Gruel with beans/groundnuts	<i>grbean3</i>	1.1	0.9	0.9	1.9	0.5	0.7	0.8
Other food prepared with WSB	<i>owsb3</i>	16.0 ^a	4.5 ^a	8.8**	17.9 ^a	3.3 ^a	5.3**	7.1
Any of the above promoted in Mothers' Clubs	<i>recfood3</i>	87.2 ^a	48.0 ^{a, c}	62.8***	85.9 ^a	33.8 ^{a, c}	41.1***	51.9
Foods/groups consumed 7 or more times in the last 7 days								
Eggs	<i>eggs7</i>	5.3 ^a	1.5 ^a	2.9	2.8	1.2	1.5	2.2
Flesh foods (meat, poultry, fish, seafood)	<i>flesh7</i>	37.7 ^{a, b}	28.1 ^{a, c}	31.7***	26.4 ^b	21.7 ^c	22.3***	27.0
Vitamin A-rich orange/red fruits/vegetables	<i>orange7</i>	22.8 ^a	31.7 ^a	28.4	35.9	29.3	30.2	29.3
Any of the recipes promoted in Mothers' Clubs	<i>recfood7</i>	54.5 ^a	17.1 ^{a, c}	31.2***	48.1 ^a	9.3 ^{a, c}	14.8***	22.9

* Significant at $p \leq 0.05$. P-value for comparison of preventive and recuperative areas; results from xtlogit for dichotomous variables.

** Significant at $p \leq 0.01$. P-value for comparison of preventive and recuperative areas; results from xtlogit for dichotomous variables.

*** Significant at $p \leq 0.001$. P-value for comparison of preventive and recuperative areas; results from xtlogit for dichotomous variables.

^a Difference between participants and nonparticipants (within program group) is significant at $p < 0.05$ (random effects regression analysis).

^b Difference between program groups is significant at $p < 0.05$ for participants (random effects regression analysis).

^c Difference between program groups is significant at $p < 0.05$ for nonparticipants (random effects regression analysis).

**ANNEX 8.12 MEAN DIETARY DIVERSITY, BY AGE GROUP, INTERVENTION AREA, AND PARTICIPATION
(CHILDREN 6-42 MONTHS)**

	Variable name	Prev + exp		Prev + not exp		All prev		Recp + exp		Recp + not exp		All recp		Total	
		Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Child age															
- 6-11 mo	dietdiv8	3.4	1.9	- ^a	- ^a	3.4	1.9	3.1	1.6	- ^a	^a	3.1	1.6	3.2	1.8
- 12-17 mo	dietdiv8	4.4 ^c	1.5	4.4	1.6	4.4*	1.5	4.0 ^c	1.5	4.0	1.4	4.0*	1.5	4.2	1.5
- 18-23 mo	dietdiv8	4.6 ^{b, c}	1.5	3.9 ^b	1.2	4.5	1.4	4.2 ^c	1.4	4.0	1.5	4.2	1.4	4.3	1.4
- 24-29 mo	dietdiv8	4.5	1.4	4.2	1.2	4.5	1.4	4.3 ^b	1.4	3.7 ^b	1.5	4.2	1.4	4.4	1.4
- 30-35 mo	dietdiv8	4.3	1.2	4.4	1.3	4.3	1.2	4.3	1.5	4.3	1.5	4.3	1.5	4.3	1.3
- 36 mo and above	dietdiv8	4.5	1.5	4.0	1.2	4.4	1.5	4.1	1.4	3.9	1.5	4.1	1.4	4.2	1.4
All children	dietdiv8	4.4 ^c	1.5	4.2	1.3	4.3***	1.5	4.1 ^c	1.5	3.9	1.5	4.1***	1.5	4.2	1.5

Note: Shaded area indicates n in cell size <= 20.

* Significant at $p \leq 0.05$. P-value for comparison of preventive and recuperative areas; results from xtlogit for dichotomous variables.

*** Significant at $p \leq 0.001$. P-value for comparison of preventive and recuperative areas; results from xtlogit for dichotomous variables.

^a The result is not presented due to small sample size.

^b Difference between participants and nonparticipants (within program group) is significant at $p < 0.05$ (random effects regression analysis).

^c Difference between program groups is significant at $p < 0.05$ for participants (random effects regression analysis).

ANNEX 8.13 RECEIPT OF VITAMIN AND MINERAL SUPPLEMENTS, BY INDEX CHILDREN (12-41 MONTHS) AND THEIR MOTHERS^a

	Variable name	Prev + exp (n = 622)	Prev + not exp (n = 126)	All prev (n = 748)	Recp + exp (n = 626)	Recp + not exp (n = 126)	All recp (n = 752)	All children (n = 1,500)
					(percent)			
Percent received vitamin A capsule in last 6 mo (children)	<i>q402a</i>	56.8 ^{b, c}	26.2 ^b	51.6*	47.9 ^{b, c}	34.9 ^b	45.7*	48.7
Percent of women who received prenatal iron supplements	<i>q309</i>	86.3 ^b	62.4 ^{b, d}	82.3	85.9 ^b	74.6 ^{b, d}	84.0	83.2
Percent of women who received postpartum vitamin A	<i>q316</i>	70.9 ^{b, c}	27.0 ^b	63.5***	58.6 ^{b, c}	31.7 ^b	54.1***	58.8

* Significant at $p \leq 0.05$. P-value for comparison of preventive and recuperative areas; results from xtlogit for dichotomous variables.

*** Significant at $p \leq 0.001$. P-value for comparison of preventive and recuperative areas; results from xtlogit for dichotomous variables.

^a Data were not available for younger siblings receipt of vitamin A capsules, nor for supplements received by their mothers during and after birth of younger sibling. Data were available for receipt of WSB and Sprinkles, but too few children received (35 for WSB and 24 for Sprinkles) to allow analysis.

^b Difference between participants and nonparticipants (within program group) is significant at $p < 0.05$ (random effects regression analysis).

^c Difference between program groups is significant at $p < 0.05$ for participants (random effects regression analysis).

^d Difference between program groups is significant at $p < 0.05$ for nonparticipants (random effects regression analysis).

ANNEX 8.14 FEEDING DURING AND AFTER DIARRHEA: INDEX CHILDREN^a

Changes in feeding	Variable name	Prev + exp	Prev + not exp	All prev	Recp + exp	Recp + not exp	All recp	All children
					(percent)			
Index children (age group 12-41 months)		(n = 226)	(n = 42)	(n = 268)	(n = 208)	(n = 40)	(n = 248)	(n = 516)
Liquids	<i>q409</i>							
- gives nothing to drink)		3.1	4.8	3.4	5.3	5.0	5.2	4.3
- gave much less		6.6	4.8	6.3	7.2	2.5	6.5	6.4
- gave a little less		21.7	26.2	22.4	25.5	20.0	24.6	23.6
- gave about the same		16.4	26.2	22.4	25.5	20.0	24.6	23.6
- gave more liquid		52.2	38.1	50.0	43.8	50.0	44.8	47.4
Semisolid and solid foods	<i>q410</i>							
- stopped giving foods		3.5	9.5	4.5	2.4	0.0	2.0	3.3
- gave much less		11.5	21.4	13.1	8.2	7.5	8.1	10.6
- gave a little less		26.1	26.2	26.1	35.6	35.0	35.5	30.6
- gave about the same		25.2	28.6	25.7	25.0	32.5	26.2	26.1
- gave more food		33.6 ^b	14.3 ^b	30.6	28.8	25.0	28.2	29.4
Among those who recovered		(n = 195)	(n = 39)	(n = 234)	(n = 182)	(n = 37)	(n = 219)	(n = 453)
Percent who gave an extra meal per day	<i>q415n</i>	72.8 ^{b, c}	35.9 ^b	66.7*	59.3 ^{b, c}	37.8 ^b	55.7*	61.4
Among those given extra meals		(n = 546)	(n = 1)	(n = 547)	(n = 212)	(n = 0)	(n = 212)	(n = 759)
Mean number of days extra meal given	<i>q416n</i>	6.2	6.0	6.2	6.8	6.6	6.8	6.5

Note: Shaded area indicates n in cell size <= 20.

* Significant at $p \leq 0.05$. P-value for comparison of preventive and recuperative areas; results from xtlogit for dichotomous variables.

^a These questions were only asked if the child had had diarrhea in the last 2 weeks (n = 517 for index children). Results are not reported for younger siblings, because most were exclusively breastfed; the sample size for children with diarrhea and not exclusively breastfed was too small (n = ?).

^b Difference between participants and nonparticipants (within program group) is significant at $p < 0.05$ (random effects regression analysis).

^c Difference between program groups is significant at $p < 0.05$ for participants (random effects regression analysis).

ANNEX 8.15 IMMUNIZATION STATUS, BY INTERVENTION GROUP AND BY PARTICIPATION (AMONG CHILDREN 12-41 MONTHS OF AGE)

Immunization status	Variable name	Prev + exp	Prev + not exp	All prev	Recp + exp	Recp + not exp	All recp	All children
		Percent						
		(n = 371)	(n = 55)	(n = 426)	(n = 400)	(n = 61)	(n = 461)	(n = 887)
Fully immunized (according to card)	<i>immcard</i>	31.0 ^a	14.6 ^a	28.9	24.8	26.2	25.0	26.8
		(n = 86)	(n = 39)	(n = 125)	(n = 74)	(n = 30)	(n = 104)	(n = 229)
Fully immunized (from recall)	<i>immrecal</i>	30.2	20.5	27.2	35.1	13.3	28.9	28.0
		(n = 457)	(n = 94)	(n = 551)	(n = 474)	(n = 91)	(n = 565)	(n = 1,116)
Fully immunized (card or recall)	<i>fimmuniz</i>	30.9 ^a	17.0 ^a	28.5	26.4	22.0	25.7	27.1

^a Difference between participants and nonparticipants (within program group) is significant at $p < 0.05$ (random effects regression analysis).

ANNEX 9.1 SCORING OF VARIABLES FOR THE FOOD INSECURITY SCALE

No.	Food insecurity-related experience	Scoring
1.	Bought a cereal that was less preferred due to lack of money	Yes = 1 No = 0 Like all cereals = 0
2.	Extent of gap between number of meals consumed during the day and ideal number of meals	No difference = 0 Gap of 1 meal = 0.5 Gap of 2 or more meals = 1
3.	Ate less frequently due to lack of food	Yes = 1 No = 0
4.	Ate less frequently due to lack of fuel (or money to buy fuel)	Yes = 1 No = 0
5.	Cooked with less beans than usual	Yes = 1 No = 0
6.	Cooked with no beans	Yes = 1 No = 0
7.	Cooking without even adding the head of a herring for flavor	Yes = 1 No = 0
8.	Cooked same food day after day	Yes = 1 No = 0
9.	Frequency of going to bed hungry in past 30 days	Never = 0 1-7 times = 0.5 More than 7 times = 1
10.	Children ever went to bed hungry in past 30 days	Yes = 1 No = 0
11.	Frequency of having worried about not having enough food in past 30 days	Never = 0 1-2 times = 0.25 At least once a week = 0.5 Almost every day = 1
	Total possible score	Maximum possible score = 11
	Terciles of food insecurity (based on distribution of food insecurity scale)	1 = low food insecurity 2 = moderate food insecurity 3 = severe food insecurity

ANNEX 10.1 SCORING OF WOMEN'S WELL-BEING SCALES

Measured symptom	Variable	Scoring
A. Mental stress scale		
Poor appetite	Q904a	No = 0 Yes = 1
Shaking/trembling hands	Q904b	
Being easily excited/irritable	Q904c	
Difficulty in enjoying daily life	Q904d	
Difficulty to do daily work	Q904e	
Getting easily tired	Q904f	
Total possible score	Q904ment	Minimum = 0; Maximum = 6
B. Frequency of stress symptoms scale		
Headaches	Q905a	Never = 0 Sometimes = 1 Often = 2
Get easily frightened	Q905b	
Sleep poorly	Q905c	
Suffer from poor digestion	Q905d	
Have trouble thinking clearly	Q905e	
Feel sad or unhappy	Q905f	
Lose interest in things	Q905g	
Feel tired all the time	Q905h	
Don't want to play with the children	Q905i	
Too tired to play with children	Q905j	
Total possible score	Q905htlh	Minimum = 0; Maximum = 20
C. Time stress scale		
Feel there is not enough time to care for house	Q905k	Never = 0 Sometimes = 1 Often = 2
Feel there is not enough time to care for children	Q905l	
Feel there is not enough time to care for self	Q905m	
Feel worried there is not enough time to do daily work	Q905n	
Total possible score	Q905time	Minimum = 0; Maximum = 8
D. Life satisfaction scale		
- Are satisfied with the way they live	Q903a	Yes = 2 Neither yes or no = 1 No = 0
- Have the important things they wanted in their life	Q903b	
- Would change their life over if they could	Q903c	
- Are happy with their last child	Q903d	
- Like their daily work	Q903e	
- Satisfied with husband/partner's help	Q903f	
- Satisfied with help from mother-in-law	Q903g	
- Satisfied with help from mother	Q903h	
- Satisfied with help received from other family members	Q903i	
- Satisfied with help from those outside their family	Q903j	
Total possible	Q903lsat	Minimum = 0; Maximum = 20
D. Life satisfaction scale 2 (not including support from spouse or mother-in-law)		
- Are satisfied with the way they live	Q903a	Yes = 2 Neither yes or no = 1 No = 0
- Have the important things they wanted in their life	Q903b	
- Would change their life over if they could	Q903c	
- Are happy with their last child	Q903d	
- Like their daily work	Q903e	
- Satisfied with husband/partner's help	Q903f	
- Satisfied with help received from other family members	Q903i	
- Satisfied with help from those outside their family	Q903j	
Total possible	Q903lsat2	Minimum = 0; Maximum = 16

ANNEX 12.1 SUMMARY OF EVALUATION RESULTS

Outcome	Baseline differences	Overall difference between program communities at final survey and improvements since baseline ^a	Difference by program participation ^a	
			CURRENT PARTICIPATION	EVER PARTICIPATED
ANTHROPOMETRIC OUTCOMES ^b				
Height for age Z-score (HAZ)	None P: -1.39 R: -1.38	P: -1.30 R: -1.43 Difference in favor of P: +0.13 Z-scores	<u>Overall:</u> N/A ^c <u>Within program group:</u> Inappropriate comparison because of age-confounding in preventive and targeting of malnourished in recuperative	<u>Overall:</u> N/A <u>Within program group:</u> <i>Preventive:</i> No differences by participation. <i>Recuperative:</i> Inappropriate comparison because program targets more malnourished children
Stunting prevalence	None P: 28.6% R: 28.5%	P: 23.2% R: 28.7% Difference in favor of P: +5.5 percentage points	<u>Overall:</u> N/A <u>Within program group:</u> Inappropriate comparison because of age-confounding in preventive and targeting of malnourished in recuperative	<u>Overall:</u> N/A <u>Within program group:</u> <i>Preventive:</i> 23.7% stunting among ever participated and 22.1% among nonparticipants (difference not statistically significant) <i>Recuperative:</i> Not valid comparison because program targets more malnourished children
Weight for age Z-score (WAZ)	None P:-1.28 R:-1.26	P: -1.26 R: -1.49 Difference in favor of P: +0.23 Z-scores	<u>Overall:</u> N/A <u>Within program group:</u> Inappropriate comparison because of age-confounding in preventive and targeting of malnourished in recuperative	<u>Overall:</u> N/A <u>Within program group:</u> <i>Preventive:</i> Ever participated is worse off by 0.08 Z-scores (difference not statistically significant).

^a Only differences that are statistically significant at $P < 0.05$ are reported in this summary table (except where noted).

^b Note that although differences by participation (current and ever) are presented in this table for anthropometric outcomes, they are not discussed in the text because of noncomparability of these indicators among participants between program approaches.

^c Participants are not comparable between groups due to different targeting, therefore overall participant-nonparticipant comparison is not valid.

Outcome	Baseline differences	Overall difference between program communities at final survey and improvements since baseline ^a	Difference by program participation ^a	
			CURRENT PARTICIPATION	EVER PARTICIPATED
Underweight prevalence	None P: 24.3% R: 25.4%	P: 22.9% R: 30.1% Difference in favor of P: +7.2 percentage points	<u>Overall:</u> N/A <u>Within program group:</u> Inappropriate comparison because of age-confounding in preventive and targeting of malnourished in recuperative	<u>Overall:</u> N/A <u>Within program group:</u> <i>Preventive:</i> 23.9% underweight among ever participated and 20.2% among nonparticipants (difference not statistically significant). <i>Recuperative:</i> Inappropriate comparison because program targets undernourished children
Weight for height Z-score (WHZ)	None P: -0.53 R: -0.52	P: -0.58 R: -0.78 Difference in favor of P: +0.20 Z-scores	<u>Overall:</u> N/A <u>Within program group:</u> Inappropriate comparison because of age-confounding in preventive and targeting of malnourished in recuperative	<u>Overall:</u> N/A <u>Within program group:</u> <i>Preventive:</i> ever participated is worse than never participated by 0.1 Z-scores (difference not statistically significant). <i>Recuperative:</i> Inappropriate comparison because program targets undernourished children
Wasting prevalence	None P: 6.0% R: 5.4%	P: 5.2% R: 9.2% Difference in favor of P: +4 percentage points	<u>Overall:</u> N/A <u>Within program group:</u> Inappropriate comparison because of age-confounding in preventive and targeting of malnourished in recuperative	<u>Overall:</u> N/A <u>Within program group:</u> <i>Preventive:</i> 5.6% wasting among ever participated versus 4.0% among nonparticipants (difference not statistically significant). <i>Recuperative:</i> Inappropriate comparison because program targets undernourished children
PROGRAM PARTICIPATION				
<i>Use of RPs</i>	Not assessed	No difference between program groups. High RP use in both groups (> 90% of children were taken to RP) but RP use patterns are age dependent. Fewer older children taken to preventive area RPs.	N/A	N/A

Outcome	Baseline differences	Overall difference between program communities at final survey and improvements since baseline ^a	Difference by program participation ^a	
			CURRENT PARTICIPATION	EVER PARTICIPATED
<i>Enrollment in food assistance and BCC package</i>	N/A: Program had not started at baseline	<ul style="list-style-type: none"> - Participation rates are same in program groups for pregnant and lactating women (58% and 63%). - Current enrollment of child beneficiaries is higher in preventive (37% vs. 14.1%). - More children in preventive areas were ever exposed to the program (73% vs. 28%) 	N/A	N/A
HOUSEHOLD RESOURCES FOR CARE				
<i>Household characteristics</i>	None	<p>Overall, no major differences in most characteristics. Household durable goods and productive asset ownership slightly higher in preventive group.</p> <p>Household asset ownership same as baseline in preventive, but lower than baseline in recuperative.</p>	<p><u>Overall</u>: No differences between households in construction, number of rooms, home ownership, sources of drinking water or household assets. Greater household size among current participants compared to nonparticipant.</p> <p><u>Within program group</u>: <i>Preventive</i>: Number of household assets higher among current participants in preventive. No difference for other household characteristics <i>Recuperative</i>: No differences in participant-nonparticipant comparisons.</p>	<p><u>Overall</u>: No differences between households in construction, number of rooms, home ownership, sources of drinking water. Household size larger among participants in both groups.</p> <p><u>Within program group</u>: No differences in participant-nonparticipant comparisons in either group.</p>
Food security				
<i>Food security experiences</i>	None	<p>Greater food insecurity among recuperative group</p> <p>Severe food insecurity higher among recuperative</p> <p>No improvements since baseline, and overall food insecurity is very high among both groups</p>	<p><u>Overall</u>: Lower food insecurity among current participants.</p> <p><u>Within program group</u>: Difference in food insecurity between current and nonparticipants larger in preventive versus recuperative.</p>	<p><u>Overall</u>: No difference.</p> <p><u>Within program group</u>: No differences in participant-nonparticipant comparisons between groups.</p>

Outcome	Baseline differences	Overall difference between program communities at final survey and improvements since baseline ^a	Difference by program participation ^a	
			CURRENT PARTICIPATION	EVER PARTICIPATED
Months of household food insufficiency	Not assessed at baseline	Preventive communities better off in terms of months of food insufficiency and severity of insufficiency. Preventive program somewhat protective of food insufficiency in most severely food insufficient months.	<u>Overall</u> : Fewer months of food insufficiency among current participants <u>Within program group</u> : Greater/more significant difference between participants and nonparticipants in preventive.	<u>Overall</u> : No difference. <u>Within program group</u> : No differences in participant-nonparticipant comparisons between groups.
Severity of months of household food insufficiency	Not assessed at baseline	Preventive communities better off in terms of months of food insufficiency and severity of insufficiency. Preventative program somewhat protective of food insufficiency in most severely food insufficient months.	<u>Overall</u> : Fewer months of food insufficiency among current participants. <u>Within program group</u> : Greater/more significant difference between participants and nonparticipants in preventive.	<u>Overall</u> : No difference. <u>Within program group</u> : No differences in participant-nonparticipant comparisons between groups.
CAREGIVER RESOURCES FOR CARE				
General characteristics	None	Overall, no differences between groups in caregiver education, partner's education, marital status, employment status, BMI, asset ownership by respondents, control over household purchases, involvement in decisionmaking, material and financial support. Work patterns slightly different between groups (but nonsignificant). Respondents in preventive communities work more outside home, spend more time outside.	<u>Overall</u> : Higher social support for participants, higher rates of employment among participants, and longer duration away from home. <u>Within program group</u> : Larger difference in social support between participants and nonparticipants in preventive. More pregnant women among current beneficiaries in recuperative compared to preventive or nonparticipant recuperative (27 % vs. 15%).	<u>Overall</u> : Higher rates of pregnancy among nonparticipants than participants. Better communication with spouse among participants. Slightly higher asset ownership among participant respondents. Slightly higher proportion of partnered women among participants. <u>Within program group</u> : No differences in participant-nonparticipant comparisons between groups.

Outcome	Baseline differences	Overall difference between program communities at final survey and improvements since baseline ^a	Difference by program participation ^a	
			CURRENT PARTICIPATION	EVER PARTICIPATED
<i>Caregiver knowledge</i>				
Baseline scales (introduction of foods, feeding frequency, overall knowledge)	None	Knowledge higher among P v. R. Knowledge scores improved overall from baseline to final.	<u>Overall</u> : Overall significant differences between participants and nonparticipants in both groups. <u>Within program group</u> : Differences between participants and nonparticipants greater in P v. R.	<u>Overall</u> : Greater improvements in nutrition knowledge from baseline among participants. Large difference between those who had ever been exposed to the program and those who had never been exposed. <u>Within program group</u> : Differences between participants and nonparticipants greater in P v. R.
BCC topic specific scales (overall, BF, complementary feeding, child illness, general health, malnutrition)	Not assessed at baseline	Slightly better in P communities v. R communities.	<u>Overall</u> : Differences between current participants and nonparticipants stat sig. for overall knowledge, BF, and complementary feeding. Marginally significant for childhood illness. <u>Within program group</u> : BF knowledge significant higher in P In R., sig. better overall knowledge, BF, and complementary feeding knowledge among participants.	<u>Overall</u> : Large differences between participants and nonparticipants <u>Within program group</u> : More significant differences among participants and nonparticipants in P vs. R, especially for overall knowledge score.
Mental and physical well-being	None	Preventive area caregivers better off on 4 women’s well-being measures. Slight improvement since baseline on mental stress and other well-being measures	<u>Overall</u> : Current participation associated with better self-rated health, lower mental stress and lower time stress. <u>Within program group</u> : Differences between participants and nonparticipants larger in preventive areas.	<u>Overall</u> : No differences between respondents who had ever been exposed to the program and those who had not.
IMPACT ON CARE PRACTICES				
Awareness, trial, and adoption of recommended practices	Not assessed at baseline	Awareness, trial, and adoption better in preventive compared to recuperative for complementary feeding related practices. No difference for breast-feeding related practices (as expected by design) Drop off from awareness to trial to adoption dependent on participation and type of behavior being recommended.	Not evaluated	<u>Overall</u> : Knowledge and adoption for all practices is higher in participants v. nonparticipants. Trial was higher among participants for all practices except adding a beaten egg to a child’s gruel trial. <u>Within program groups</u> : No difference between program groups in patterns of awareness, trial, and adoption by participation.

Outcome	Baseline differences	Overall difference between program communities at final survey and improvements since baseline ^a	Difference by program participation ^a	
			CURRENT PARTICIPATION	EVER PARTICIPATED
<i>Breastfeeding</i>				
Initiation of BF	None	Program groups are very similar. Improved practices since baseline. 69% initiate BF w/in 1 st hr; 88% gave colostrum; only 9% reported giving pre-lacteals. Younger siblings more likely BF in 1 st hr vs. index children (83% v. 65%), more likely to been given colostrum (95% v. 86%).	N/A	Overall: Sig. > percent of participants BF w/in 1hr, gave colostrum, and did not give pre-lacteals compared to nonparticipants. <u>Within program groups:</u> Differences between participants and nonparticipants were large for each practice and not different between program groups.
Exclusive BF	None	Program groups are very similar. EBF improved since baseline: among children < 6mo old at final survey--92% exclusive BF in last 24h, compared to 47% at baseline.	N/A; too few nonparticipants in this age subgroup (< 6 mo)	N/A; too few nonparticipants in this age subgroup (< 6 mo)
Duration of any BF	None	Similar BF pattern in P & R group at final. No differences since baseline. Most children breastfed to 18mo, sharp decline in second half of child’s second year.	Comparison between current participants and nonparticipants is confounded by large differences in age distribution; differences within 6-mo age groups NS with low statistical power, except more children 18-23 mo were BF among current partic in R (68%) than among nonpartic in R (38%).	Differences within 6-mo age groups are NS with low statistical power No evidence that participation (current or ever) compromised continued breastfeeding
<i>Complementary feeding</i>				
Introduction of foods	None	No difference between program groups. Large improvements since baseline; 86% delayed introduction of other liquids until at least 6 mo; 77% of women reported first giving semi-solid food to index children at 6-6.9 mo	N/A	<u>Overall:</u> Early introduction of liquids and semi-solids much more likely to be reported among nonparticipants v. participants.
Feeding frequency	None	No meaningful differences between P & R in frequency of feeding. Breastfed children 12-23 mo much more likely to receive minimum recommended number of meals (46% received 3+ meals) compared to non-breastfed (only 14% received recommended 4+ meals)	<u>Overall:</u> No significant differences by current participation status; for non-breastfed children 12-23 mo, frequency of feeding and likelihood of receiving minimum recommended number of meals higher among current participants (23%) vs. nonparticipants (8%).	<u>Overall:</u> No significant differences

Outcome	Baseline differences	Overall difference between program communities at final survey and improvements since baseline ^a	Difference by program participation ^a	
			CURRENT PARTICIPATION	EVER PARTICIPATED
Feeding patterns	None	Small difference between groups; children in P slightly more likely to have received evening meal yesterday than children in R. Not much improvement since baseline and only 11% index children received evening meal on day before survey. 19% of mothers reported they wake index child for evening meal.	<u>Overall</u> : No differences	<u>Overall</u> : No differences
Dietary diversity	None	Slightly higher in P than in R; lower than baseline in both areas.	<u>Overall</u> : Slightly higher among current participants in both areas. <u>Within program groups</u> : Much higher for currently participating children 6-11 mo in preventive group (4.2 vs. 2.5 food groups for nonparticipants)	Higher among participants in some age groups (18-23 in P and 24-29 in R)
Nutrient rich foods/new recipes	None	P more likely than R to have eaten Vit A rich fruits and veg, other fruit and veg, dairy, and other animal-source foods yesterday. Children in P more likely to have had eggs, beef and pork; 63% in P group vs. 41% in R group had recommended recipes ≥ 3 x/wk, 31% v. 15% had them 7x or more in last week (P v. R). Decline since baseline in consumption of animal source foods	No major differences except younger siblings (6-11 mo) of current participants in P more likely to have legumes (72% vs. 44% of nonparticipants)	<u>Overall</u> : Nonparticipants less likely to have eaten several nutrient-dense animal source food groups. <u>Within program groups</u> : <i>Preventive</i> : Index children who were participants much more likely to have had flesh foods (meat poultry fish) 7 or more times in the last week (36% of those ever participating vs. 8% of those never participating).

Outcome	Baseline differences	Overall difference between program communities at final survey and improvements since baseline ^a	Difference by program participation ^a	
			CURRENT PARTICIPATION	EVER PARTICIPATED
Use of fortified foods and vitamin-mineral supplements	None	P more likely to have ever received WSB and to be currently receiving WSB. Mean age at receiving WSB was 7.7mo in preventive vs. 13.6 mo in recuperative areas; P areas received rations for more months (11.7 v. 7.5) Vit A supplementation for index children in last 6 mo and for postpartum mothers higher in preventive group. Vit. A capsule coverage increased from 29% to 49%. Iron supplement coverage increased 65% to 83% at final survey. Postpartum Vit. A supplementation increased from 8% to 59% at final survey and higher in preventive group (71%).	<u>Overall:</u> Participants much more likely to have consumed WSB yesterday, by design. <u>Within program groups:</u> More participants in preventive group consumed WSB, compared to recuperative, because of greater participation rates.	N/A: Only currently participating children receive WSB Participants in both P & R much more likely to have received vitamin A (children and mothers) and somewhat more likely to receive prenatal iron (mothers).
Feeding during and after diarrhea	None	No difference by program group in reporting of giving more liquids, offering the same or more solid foods. After recovery, 61% reported able to give child an extra meal/d. P group more likely to give extra meal (67% v. 56%).	<u>Overall:</u> Current participants more likely to report giving an extra meal after diarrhea (76% vs. 54%)	<u>Overall:</u> Those who had ever participated were much more likely to report giving an extra meal (66% vs. 37%)
Responsive Feeding	None	Fewer children reported to feed themselves than at baseline, and slightly fewer in P than in R. At final, more women could name positive strategies for coping when child refuses food. 65% named 2+ strategies; this was slightly higher in P than in R. Fewer women reported negative strategies than at baseline, with no difference between P & R.	N/A	<u>Overall:</u> Mothers who had participated were more likely to report taking action when child refused to eat (both areas). <u>Within program groups:</u> Women in P who had participated were more likely to know 2+ positive strategies.
Immunization and care-seeking during illness	None	Immunization: increased coverage for full immunization since baseline, but coverage remains low (27%), with no difference between program areas. Decrease since baseline in care-seeking during illness, no major differences between P & R.	<u>Overall:</u> No differences in immunization or care-seeking related to current participation.	<u>Overall:</u> No major differences in immunization or care-seeking. <u>Within program group:</u> Higher rates of full immunization among participants in P, but not in R.